



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2026 – 08:35 AM UTC

PDB ID : 4FYX / pdb_00004fyx
Title : E. coli Aspartate Transcarbamoylase complexed with dCTP, UTP, and Mg²⁺
Authors : Cockrell, G.M.; Kantrowitz, E.R.
Deposited on : 2012-07-05
Resolution : 2.09 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	8172 (2.10-2.06)
Clashscore	190562	8714 (2.10-2.06)
Ramachandran outliers	187476	8641 (2.10-2.06)
Sidechain outliers	187428	8642 (2.10-2.06)
RSRZ outliers	180081	8177 (2.10-2.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	310	6% 91% 7%
1	C	310	7% 92% 6%
2	B	153	10% 91% 5%
2	D	153	3% 92% 5%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 7901 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aspartate carbamoyltransferase catalytic chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	310	Total	C	N	O	S	0	0	0
			2415	1527	423	456	9			
1	C	310	Total	C	N	O	S	0	0	0
			2415	1527	423	456	9			


- Molecule 2 is a protein called Aspartate carbamoyltransferase regulatory chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	148	Total	C	N	O	S	0	0	0
			1160	729	205	221	5			
2	D	148	Total	C	N	O	S	0	0	0
			1159	729	205	220	5			

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Zn	0	0
			1	1		
3	D	1	Total	Zn	0	0
			1	1		

- Molecule 4 is URIDINE 5'-TRIPHOSPHATE (CCD ID: UTP) (formula: C₉H₁₅N₂O₁₅P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total	C	N	O	P	0	0
			29	9	2	15	3		
4	D	1	Total	C	N	O	P	0	0
			29	9	2	15	3		

- Molecule 5 is 2'-DEOXYCYTIDINE-5'-TRIPHOSPHATE (CCD ID: DCP) (formula: $C_9H_{16}N_3O_{13}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	B	1	Total	C	N	O	P	0	0
			28	9	3	13	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	D	1	Total	C	N	O	P	0	0
			28	9	3	13	3		

- Molecule 6 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total	Mg	0	0
			1	1		
6	D	1	Total	Mg	0	0
			1	1		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	202	Total	O	0	0
			202	202		
7	B	98	Total	O	0	0
			98	98		
7	C	222	Total	O	0	0
			222	222		
7	D	112	Total	O	0	0
			112	112		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Aspartate carbamoyltransferase catalytic chain




- Molecule 1: Aspartate carbamoyltransferase catalytic chain



- Molecule 2: Aspartate carbamoyltransferase regulatory chain



- Molecule 2: Aspartate carbamoyltransferase regulatory chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 3 2 1	Depositor
Cell constants a, b, c, α , β , γ	121.21Å 121.21Å 141.76Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	49.22 – 2.09 49.22 – 2.09	Depositor EDS
% Data completeness (in resolution range)	100.0 (49.22-2.09) 99.7 (49.22-2.09)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.56 (at 2.08Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.165 , 0.212 0.174 , 0.216	Depositor DCC
R_{free} test set	3630 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	29.6	Xtriage
Anisotropy	0.057	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 41.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.021 for -h,-k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7901	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.65% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, UTP, MG, DCP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.47	0/2461	0.77	1/3339 (0.0%)
1	C	0.53	0/2461	0.82	3/3339 (0.1%)
2	B	0.41	0/1177	0.73	0/1590
2	D	0.41	0/1176	0.73	0/1590
All	All	0.47	0/7275	0.78	4/9858 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	80	SER	N-CA-C	10.12	120.13	108.49
1	C	269	ARG	N-CA-C	-5.42	100.87	109.76
1	C	222	VAL	N-CA-C	5.31	115.93	109.30
1	A	84	LYS	N-CA-C	5.19	116.63	110.19

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2415	0	2422	15	0
1	C	2415	0	2422	13	0
2	B	1160	0	1183	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	1159	0	1183	4	0
3	B	1	0	0	0	0
3	D	1	0	0	0	0
4	B	29	0	11	0	0
4	D	29	0	11	0	0
5	B	28	0	12	0	0
5	D	28	0	12	0	0
6	B	1	0	0	0	0
6	D	1	0	0	0	0
7	A	202	0	0	4	0
7	B	98	0	0	0	0
7	C	222	0	0	4	0
7	D	112	0	0	1	0
All	All	7901	0	7256	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:17:ARG:NH2	7:A:530:HOH:O	2.18	0.76
1:A:153:ASP:OD2	7:A:530:HOH:O	2.10	0.69
1:C:305:ASN:ND2	7:C:582:HOH:O	2.30	0.64
1:C:9:ILE:HG21	1:C:299:LEU:HD21	1.80	0.63
2:B:99:LEU:HD21	2:B:134:ILE:HD12	1.79	0.62
1:C:29:LYS:NZ	7:C:597:HOH:O	2.29	0.61
1:A:84:LYS:HB2	1:A:86:GLU:H	1.66	0.61
1:A:9:ILE:HG21	1:A:299:LEU:HD21	1.83	0.61
1:C:80:SER:OG	1:C:81:LEU:N	2.28	0.58
1:C:76:SER:OG	1:C:77:ALA:N	2.39	0.54
1:C:82:GLY:O	1:C:86:GLU:N	2.21	0.54
1:C:279:LYS:HG2	7:C:404:HOH:O	2.07	0.54
1:A:17:ARG:NE	7:A:447:HOH:O	2.41	0.53
1:A:17:ARG:NH1	7:A:405:HOH:O	2.41	0.53
2:D:10:GLU:OE1	7:D:399:HOH:O	2.18	0.52
1:C:79:THR:OG1	1:C:83:LYS:NZ	2.44	0.51
2:B:71:VAL:O	2:B:97:PRO:HB3	2.14	0.48
1:A:116:THR:HG22	1:A:124:VAL:HG13	1.95	0.47
2:D:22:PRO:HG2	2:D:25:ILE:HG13	1.96	0.47
1:C:218:VAL:O	1:C:222:VAL:HG23	2.15	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:269:ARG:HB2	7:C:557:HOH:O	2.13	0.47
1:A:189:PRO:HG2	1:A:192:LEU:HB2	1.96	0.47
2:D:81:ALA:O	2:D:96:ARG:HD2	2.15	0.46
1:A:244:LYS:HA	1:A:247:PHE:CZ	2.51	0.46
1:A:229:ARG:NH1	1:A:231:GLN:OE1	2.49	0.45
1:A:231:GLN:HB3	1:A:233:GLU:CD	2.42	0.44
1:A:36:PRO:HA	1:A:65:ARG:O	2.17	0.44
1:C:229:ARG:HA	1:C:272:GLU:CD	2.43	0.44
1:A:78:ASN:HB2	1:A:83:LYS:HB2	1.99	0.44
1:C:83:LYS:H	1:C:83:LYS:HG2	1.42	0.44
2:D:19:ASP:OD1	2:D:20:HIS:N	2.51	0.42
2:B:19:ASP:OD2	2:B:20:HIS:N	2.53	0.42
1:C:36:PRO:HA	1:C:65:ARG:O	2.19	0.42
1:A:229:ARG:HA	1:A:272:GLU:CD	2.46	0.41
1:A:75:ASP:CG	1:A:78:ASN:HD21	2.28	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	308/310 (99%)	297 (96%)	10 (3%)	1 (0%)	36	36
1	C	308/310 (99%)	293 (95%)	14 (4%)	1 (0%)	36	36
2	B	146/153 (95%)	140 (96%)	4 (3%)	2 (1%)	9	5
2	D	146/153 (95%)	143 (98%)	3 (2%)	0	100	100
All	All	908/926 (98%)	873 (96%)	31 (3%)	4 (0%)	30	28

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	76	SER
1	A	80	SER
2	B	131	ALA
2	B	132	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	261/261 (100%)	252 (97%)	9 (3%)	32	34
1	C	261/261 (100%)	250 (96%)	11 (4%)	26	26
2	B	132/137 (96%)	130 (98%)	2 (2%)	57	64
2	D	132/137 (96%)	130 (98%)	2 (2%)	57	64
All	All	786/796 (99%)	762 (97%)	24 (3%)	35	38

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	59	PHE
1	A	79	THR
1	A	81	LEU
1	A	84	LYS
1	A	124	VAL
1	A	229	ARG
1	A	231	GLN
1	A	285	TYR
1	A	309	VAL
2	B	7	LEU
2	B	99	LEU
1	C	59	PHE
1	C	80	SER
1	C	81	LEU
1	C	83	LYS
1	C	86	GLU
1	C	236	ASP
1	C	270	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	285	TYR
1	C	305	ASN
1	C	308	LEU
1	C	310	LEU
2	D	96	ARG
2	D	153	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	78	ASN
1	A	174	GLN
2	B	80	GLN
1	C	154	ASN
2	D	47	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	UTP	D	202	6	29,30,30	1.91	9 (31%)	43,47,47	1.40	6 (13%)
4	UTP	B	202	6	29,30,30	1.84	9 (31%)	43,47,47	1.41	8 (18%)
5	DCP	D	203	6	28,29,29	1.89	7 (25%)	41,45,45	0.97	2 (4%)
5	DCP	B	203	6	28,29,29	1.96	7 (25%)	41,45,45	1.01	3 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	UTP	D	202	6	-	0/22/38/38	0/2/2/2
4	UTP	B	202	6	-	0/22/38/38	0/2/2/2
5	DCP	D	203	6	-	2/22/34/34	0/2/2/2
5	DCP	B	203	6	-	3/22/34/34	0/2/2/2

All (32) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	202	UTP	C3'-C2'	-4.88	1.40	1.53
4	B	202	UTP	C3'-C2'	-4.81	1.40	1.53
5	B	203	DCP	C2'-C3'	-4.73	1.40	1.52
5	D	203	DCP	C4-N4	4.67	1.45	1.33
5	D	203	DCP	C2'-C3'	-4.67	1.41	1.52
5	B	203	DCP	C4-N4	4.59	1.45	1.33
4	D	202	UTP	PB-O3A	-4.55	1.54	1.59
5	B	203	DCP	PB-O3A	-4.02	1.55	1.59
4	B	202	UTP	PB-O3A	-3.93	1.55	1.59
4	D	202	UTP	PA-O3A	-3.46	1.55	1.59
4	B	202	UTP	PB-O3B	-3.08	1.56	1.59
5	D	203	DCP	PA-O3A	3.03	1.62	1.59
5	B	203	DCP	PA-O3A	2.80	1.62	1.59
5	D	203	DCP	PB-O3A	-2.72	1.56	1.59
5	D	203	DCP	C2-N1	-2.66	1.34	1.40
4	D	202	UTP	PB-O3B	-2.66	1.56	1.59
4	B	202	UTP	PA-O3A	-2.54	1.56	1.59
5	B	203	DCP	C2-N1	-2.46	1.34	1.40
5	D	203	DCP	C6-C5	2.34	1.40	1.35
5	D	203	DCP	O4'-C4'	-2.32	1.39	1.45
4	B	202	UTP	C2-N3	-2.32	1.33	1.38
4	D	202	UTP	C2-N3	-2.26	1.34	1.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	202	UTP	O4'-C4'	-2.26	1.40	1.45
5	B	203	DCP	C6-C5	2.25	1.40	1.35
4	B	202	UTP	O4'-C4'	-2.24	1.40	1.45
5	B	203	DCP	O4'-C4'	-2.15	1.40	1.45
4	B	202	UTP	C2-N1	2.14	1.41	1.38
4	B	202	UTP	C4-N3	-2.11	1.35	1.38
4	D	202	UTP	C4-N3	-2.11	1.35	1.38
4	D	202	UTP	C6-C5	2.10	1.40	1.35
4	D	202	UTP	C2-N1	2.09	1.41	1.38
4	B	202	UTP	C6-C5	2.06	1.39	1.35

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	202	UTP	C4-N3-C2	-4.01	121.63	126.61
4	B	202	UTP	C4-N3-C2	-3.81	121.88	126.61
4	B	202	UTP	N3-C2-N1	3.60	119.58	114.89
4	D	202	UTP	C5-C4-N3	3.59	119.83	114.80
4	D	202	UTP	N3-C2-N1	3.53	119.48	114.89
4	B	202	UTP	C5-C4-N3	3.22	119.31	114.80
4	D	202	UTP	O4-C4-C5	-2.84	120.27	125.16
4	B	202	UTP	O4-C4-C5	-2.52	120.81	125.16
5	B	203	DCP	O2-C2-N3	-2.37	118.60	122.33
4	D	202	UTP	O3G-PG-O3B	2.30	112.34	104.64
5	D	203	DCP	O5'-C5'-C4'	2.24	116.62	108.99
4	B	202	UTP	C1'-N1-C2	2.18	121.51	117.59
4	D	202	UTP	O5'-C5'-C4'	2.12	116.20	108.99
4	B	202	UTP	C3'-C2'-C1'	2.08	105.40	101.46
4	B	202	UTP	O5'-C5'-C4'	2.07	116.05	108.99
4	B	202	UTP	O1B-PB-O3B	2.06	112.83	107.27
5	B	203	DCP	O5'-C5'-C4'	2.04	115.94	108.99
5	B	203	DCP	O3G-PG-O3B	2.04	111.46	104.64
5	D	203	DCP	O3G-PG-O3B	2.00	111.36	104.64

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	203	DCP	PG-O3B-PB-O2B
5	B	203	DCP	C5'-O5'-PA-O1A
5	D	203	DCP	PG-O3B-PB-O2B
5	D	203	DCP	PG-O3B-PB-O1B

Continued on next page...


Continued from previous page...

Mol	Chain	Res	Type	Atoms
5	B	203	DCP	PB-O3A-PA-O2A


There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




Ligand UTP B 202



Ligand DCP D 203





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	310/310 (100%)	0.01	20 (6%)	25 26	19, 30, 70, 203	0
1	C	310/310 (100%)	-0.10	23 (7%)	20 22	17, 26, 80, 173	0
2	B	148/153 (96%)	0.44	15 (10%)	12 13	23, 41, 76, 137	0
2	D	148/153 (96%)	0.10	5 (3%)	48 50	22, 38, 65, 117	0
All	All	916/926 (98%)	0.06	63 (6%)	23 24	17, 32, 76, 203	0

All (63) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	81	LEU	8.0
1	A	81	LEU	7.2
1	C	243	VAL	6.4
1	A	77	ALA	6.4
1	C	79	THR	5.7
2	B	131	ALA	5.7
1	A	79	THR	5.3
1	A	76	SER	5.2
2	D	6	LYS	5.1
1	A	310	LEU	5.0
1	C	77	ALA	4.9
1	C	83	LYS	4.9
1	A	80	SER	4.7
1	C	308	LEU	4.6
1	C	310	LEU	4.5
1	C	245	ALA	4.5
1	C	76	SER	4.4
1	C	80	SER	4.4
1	A	84	LYS	4.3
1	C	244	LYS	4.1
1	C	82	GLY	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	247	PHE	4.0
1	C	84	LYS	3.8
1	C	78	ASN	3.7
1	A	85	GLY	3.6
1	A	243	VAL	3.6
1	A	83	LYS	3.6
1	A	78	ASN	3.6
1	A	247	PHE	3.5
1	C	85	GLY	3.5
1	A	309	VAL	3.5
1	C	309	VAL	3.3
1	C	231	GLN	3.3
1	A	82	GLY	3.1
2	B	130	ARG	3.1
2	D	153	ASN	3.0
1	C	246	GLN	2.9
1	C	235	LEU	2.8
2	D	152	ALA	2.8
2	B	132	ASN	2.8
2	B	8	GLN	2.7
1	A	233	GLU	2.7
2	D	7	LEU	2.6
2	B	89	TYR	2.6
2	B	92	VAL	2.6
2	B	153	ASN	2.6
2	B	7	LEU	2.6
2	B	134	ILE	2.5
2	B	99	LEU	2.5
1	C	234	ARG	2.3
2	B	6	LYS	2.3
2	B	12	ILE	2.3
1	C	233	GLU	2.2
1	A	246	GLN	2.2
1	C	270	VAL	2.2
2	B	86	ILE	2.1
1	A	271	ASP	2.1
1	A	270	VAL	2.1
2	B	96	ARG	2.1
1	A	308	LEU	2.1
2	D	55	ARG	2.1
1	A	1	ALA	2.1
2	B	9	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	UTP	B	202	29/29	0.93	0.08	36,42,57,145	0
5	DCP	B	203	28/28	0.93	0.08	32,45,57,60	0
4	UTP	D	202	29/29	0.94	0.07	31,41,52,53	0
5	DCP	D	203	28/28	0.94	0.07	30,39,51,52	0
6	MG	B	204	1/1	0.96	0.05	47,47,47,47	0
6	MG	D	204	1/1	0.98	0.03	43,43,43,43	0
3	ZN	B	201	1/1	0.99	0.02	24,24,24,24	0
3	ZN	D	201	1/1	1.00	0.01	22,22,22,22	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around UTP B 202:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around DCP B 203:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around UTP D 202:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around DCP D 203:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.