



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 02:49 AM UTC

PDB ID : 4FZS / pdb_00004fzs
Title : Structure of human SNX1 BAR domain
Authors : van Weering, J.R.T.; Sessions, R.B.; Traer, C.J.; Kloer, D.P.; Bhatia, V.K.; Stamou, D.; Hurley, J.H.; Cullen, P.J.
Deposited on : 2012-07-07
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	224	
1	B	224	

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 3341 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Sorting nexin-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	209	Total	C	N	O	S	0	0	0
			1757	1111	311	332	3			
1	B	189	Total	C	N	O	S	0	0	0
			1584	1007	274	300	3			


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	299	GLY	-	expression tag	UNP Q13596
A	300	ALA	-	expression tag	UNP Q13596
B	299	GLY	-	expression tag	UNP Q13596
B	300	ALA	-	expression tag	UNP Q13596


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Sorting nexin-1



• Molecule 1: Sorting nexin-1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	88.99Å 118.09Å 61.54Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.76 – 2.80 19.76 – 2.80	Depositor EDS
% Data completeness (in resolution range)	(Not available) (19.76-2.80) 92.0 (19.76-2.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.68 (at 2.79Å)	Xtriage
Refinement program	BUSTER 2.8.0	Depositor
R, R_{free}	0.228 , 0.258 0.248 , 0.289	Depositor DCC
R_{free} test set	719 reflections (4.71%)	wwPDB-VP
Wilson B-factor (Å ²)	67.5	Xtriage
Anisotropy	0.627	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 50.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3341	wwPDB-VP
Average B, all atoms (Å ²)	85.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.87	0/1786	1.55	14/2403 (0.6%)
1	B	0.87	0/1608	1.56	11/2160 (0.5%)
All	All	0.87	0/3394	1.56	25/4563 (0.5%)

There are no bond length outliers.

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	489	PHE	CA-CB-CG	8.43	122.23	113.80
1	A	441	ASN	N-CA-C	8.04	118.56	108.19
1	B	514	PHE	N-CA-C	6.32	121.77	113.30
1	A	306	ASP	N-CA-C	-6.13	103.89	113.02
1	A	481	PHE	CA-CB-CG	6.02	119.82	113.80
1	B	329	VAL	N-CA-CB	6.00	118.25	110.57
1	B	308	TRP	CA-C-N	5.88	128.44	120.38
1	B	308	TRP	C-N-CA	5.88	128.44	120.38
1	A	457	GLU	CA-C-N	5.72	127.94	120.28
1	A	457	GLU	C-N-CA	5.72	127.94	120.28
1	B	488	ASP	N-CA-C	-5.51	105.35	112.68
1	B	489	PHE	CA-C-N	5.51	127.66	120.28
1	B	489	PHE	C-N-CA	5.51	127.66	120.28
1	B	310	GLU	CA-C-N	5.46	127.86	120.44
1	B	310	GLU	C-N-CA	5.46	127.86	120.44
1	A	307	ILE	N-CA-C	-5.42	107.15	111.81
1	A	480	ARG	CA-C-N	5.33	127.36	120.44
1	A	480	ARG	C-N-CA	5.33	127.36	120.44
1	A	375	GLU	N-CA-C	-5.31	105.14	111.03
1	A	315	GLU	CA-C-N	5.27	127.20	120.56
1	A	315	GLU	C-N-CA	5.27	127.20	120.56
1	B	512	GLU	CB-CG-CD	5.17	121.38	112.60
1	A	346	GLN	CA-C-N	5.03	126.98	120.44
1	A	346	GLN	C-N-CA	5.03	126.98	120.44
1	A	344	THR	CB-CA-C	5.02	119.38	110.85

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1757	0	1754	26	0
1	B	1584	0	1581	16	0
All	All	3341	0	3335	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:384:GLN:O	1:A:387:ASN:HB2	1.81	0.81
1:B:323:LEU:HD21	1:B:486:SER:HB2	1.66	0.76
1:A:387:ASN:ND2	1:A:388:ASP:H	1.88	0.71
1:B:323:LEU:HD21	1:B:486:SER:CB	2.21	0.71
1:A:385:ALA:C	1:A:387:ASN:H	1.99	0.68
1:B:337:ARG:HG2	1:B:390:PHE:HZ	1.59	0.68
1:B:337:ARG:HG2	1:B:390:PHE:CZ	2.31	0.65
1:A:370:LEU:HD11	1:B:500:LEU:HD13	1.80	0.64
1:A:388:ASP:N	1:A:388:ASP:OD1	2.37	0.57
1:B:406:VAL:HG22	1:B:477:GLU:HB3	1.88	0.55
1:A:387:ASN:C	1:A:389:PHE:H	2.14	0.55
1:B:390:PHE:CD1	1:B:390:PHE:N	2.76	0.54
1:B:495:LYS:O	1:B:499:THR:HG23	2.09	0.52
1:A:495:LYS:O	1:A:499:THR:HG23	2.09	0.51
1:A:384:GLN:O	1:A:387:ASN:CB	2.56	0.51
1:A:329:VAL:HG13	1:B:367:LEU:HD21	1.93	0.50
1:B:323:LEU:CD2	1:B:486:SER:HB2	2.40	0.50
1:A:385:ALA:C	1:A:387:ASN:N	2.66	0.50
1:A:387:ASN:ND2	1:A:388:ASP:N	2.60	0.48
1:A:387:ASN:C	1:A:389:PHE:N	2.73	0.46
1:B:515:LEU:HD22	1:B:516:PRO:HD2	1.97	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:385:ALA:O	1:A:387:ASN:N	2.47	0.46
1:A:329:VAL:CG1	1:B:367:LEU:HD21	2.46	0.46
1:B:358:GLU:HG2	1:B:363:LEU:HB3	1.98	0.45
1:A:385:ALA:O	1:A:390:PHE:HB3	2.17	0.45
1:A:406:VAL:HG12	1:A:477:GLU:HB3	1.98	0.45
1:A:399:TYR:CD1	1:A:484:GLU:HB3	2.52	0.45
1:A:316:VAL:HG11	1:A:406:VAL:HG23	1.99	0.44
1:A:476:LYS:HE2	1:A:480:ARG:HH21	1.82	0.44
1:A:403:LEU:HA	1:A:406:VAL:HG22	1.99	0.43
1:A:328:ALA:O	1:A:332:THR:HG23	2.19	0.43
1:B:430:LYS:HD2	1:B:452:GLU:HB3	1.99	0.43
1:A:443:PRO:HA	1:A:446:LEU:HB2	2.01	0.42
1:A:358:GLU:HG2	1:A:363:LEU:HB3	2.02	0.42
1:A:430:LYS:HD2	1:A:452:GLU:HB3	2.01	0.42
1:B:328:ALA:O	1:B:332:THR:HG23	2.20	0.41
1:A:457:GLU:O	1:A:461:THR:HG23	2.20	0.40
1:A:500:LEU:HD13	1:B:370:LEU:HD11	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	207/224 (92%)	197 (95%)	8 (4%)	2 (1%)	12	38
1	B	181/224 (81%)	180 (99%)	1 (1%)	0	100	100
All	All	388/448 (87%)	377 (97%)	9 (2%)	2 (0%)	24	55

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	386	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	438	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	187/198 (94%)	158 (84%)	29 (16%)	2	9
1	B	169/198 (85%)	147 (87%)	22 (13%)	4	14
All	All	356/396 (90%)	305 (86%)	51 (14%)	3	11

All (51) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	306	ASP
1	A	307	ILE
1	A	308	TRP
1	A	342	LEU
1	A	363	LEU
1	A	370	LEU
1	A	379	GLN
1	A	382	GLN
1	A	383	GLU
1	A	388	ASP
1	A	391	LEU
1	A	400	ILE
1	A	412	GLN
1	A	428	GLN
1	A	429	LYS
1	A	441	ASN
1	A	442	LYS
1	A	445	LYS
1	A	446	LEU
1	A	455	GLU
1	A	461	THR
1	A	477	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	481	PHE
1	A	486	SER
1	A	490	LYS
1	A	499	THR
1	A	500	LEU
1	A	501	LEU
1	A	507	LEU
1	B	307	ILE
1	B	310	GLU
1	B	329	VAL
1	B	337	ARG
1	B	342	LEU
1	B	363	LEU
1	B	365	ARG
1	B	370	LEU
1	B	378	GLU
1	B	379	GLN
1	B	388	ASP
1	B	400	ILE
1	B	427	LEU
1	B	428	GLN
1	B	429	LYS
1	B	444	ASP
1	B	455	GLU
1	B	465	ARG
1	B	477	GLU
1	B	499	THR
1	B	500	LEU
1	B	515	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	336	HIS
1	A	384	GLN
1	A	387	ASN
1	A	418	GLN
1	A	421	GLN
1	A	424	GLN
1	A	441	ASN
1	A	504	GLN
1	B	336	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	346	GLN
1	B	448	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2			OWAB(Å ²)	Q < 0.9
1	A	209/224 (93%)	0.33	6 (2%)	53	43	52, 82, 137, 150	0
1	B	189/224 (84%)	0.20	7 (3%)	45	36	53, 76, 120, 144	0
All	All	398/448 (88%)	0.26	13 (3%)	49	39	52, 80, 128, 150	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	307	ILE	3.2
1	B	380	LEU	3.0
1	A	390	PHE	2.9
1	B	443	PRO	2.8
1	B	435	ALA	2.7
1	A	510	TYR	2.6
1	A	344	THR	2.5
1	A	442	LYS	2.4
1	A	481	PHE	2.3
1	B	477	GLU	2.2
1	B	392	LEU	2.1
1	A	341	ALA	2.1
1	B	311	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.