



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 11:13 PM UTC

PDB ID : 5G0X / pdb_00005g0x
Title : Pseudomonas aeruginosa HDAH bound to acetate.
Authors : Kraemer, A.; Meyer-Almes, F.J.; Yildiz, O.
Deposited on : 2016-03-23
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5551 (1.70-1.70)
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)
RSRZ outliers	180081	5554 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	379	88% 8% ..
1	C	379	92% 7% .

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6307 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HDAH.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	370	Total	C	N	O	S	0	0	0
			2819	1772	519	516	12			
1	C	376	Total	C	N	O	S	0	0	0
			2864	1801	527	524	12			

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	C	1	Total	Zn	0	0
			1	1		

- Molecule 3 is POTASSIUM ION (CCD ID: K) (formula: K).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	K	0	0
			2	2		
3	C	2	Total	K	0	0
			2	2		

- Molecule 4 is ACETATE ION (CCD ID: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (CCD ID: EPE) (formula: $C_8H_{18}N_2O_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	S	0	0
			15	8	2	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	S	0	0
			15	8	2	4	1		
5	A	1	Total	C	N	O	S	0	0
			15	8	2	4	1		
5	C	1	Total	C	N	O	S	0	0
			15	8	2	4	1		
5	C	1	Total	C	N	O	S	0	0
			15	8	2	4	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	268	Total	O	0	0
			268	268		
6	C	267	Total	O	0	0
			267	267		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: HDAH



• Molecule 1: HDAH



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	81.39Å 81.39Å 205.85Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	75.69 – 1.70 75.69 – 1.70	Depositor EDS
% Data completeness (in resolution range)	99.9 (75.69-1.70) 99.9 (75.69-1.70)	Depositor EDS
R_{merge}	0.28	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.25 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5.8.0103	Depositor
R, R_{free}	0.134 , 0.184 0.149 , 0.191	Depositor DCC
R_{free} test set	3852 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	14.0	Xtriage
Anisotropy	0.401	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 40.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6307	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 19.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EPE, K, ZN, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.79	0/2888	0.83	0/3922
1	C	0.79	0/2935	0.84	0/3988
All	All	0.79	0/5823	0.83	0/7910

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2819	0	2739	29	0
1	C	2864	0	2785	19	0
2	A	1	0	0	0	0
2	C	1	0	0	0	0
3	A	2	0	0	0	0
3	C	2	0	0	0	0
4	A	4	0	3	0	0
4	C	4	0	3	0	0
5	A	45	0	54	0	0
5	C	30	0	36	2	0
6	A	268	0	0	8	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	C	267	0	0	6	0
All	All	6307	0	5620	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:294:ARG:NH1	1:A:328:GLU:OE2	2.08	0.85
1:C:205:GLN:HE22	1:C:278:ARG:H	1.28	0.78
1:A:300:HIS:HD2	6:A:2225:HOH:O	1.68	0.76
1:A:203:LEU:HD12	1:A:289:MET:HE1	1.70	0.72
1:A:44:GLU:HG2	6:A:2038:HOH:O	1.92	0.68
1:A:100:GLN:O	6:A:2094:HOH:O	2.14	0.66
1:A:340:LEU:HD13	1:C:25:LEU:HD21	1.82	0.62
1:A:367:GLN:NE2	6:A:2265:HOH:O	2.33	0.60
1:C:101:ASP:OD1	5:C:1383:EPE:H51	2.02	0.60
1:A:300:HIS:CD2	6:A:2225:HOH:O	2.47	0.58
1:A:39:HIS:CD2	1:A:41:GLU:H	2.22	0.57
1:C:226:GLY:HA2	6:C:2179:HOH:O	2.03	0.57
1:C:374:GLN:NE2	6:C:2265:HOH:O	2.37	0.57
1:C:124:ASP:OD1	1:C:170:HIS:HE1	1.88	0.56
1:C:205:GLN:NE2	1:C:278:ARG:H	2.02	0.55
1:A:59:THR:HG23	1:A:64:LEU:HD21	1.90	0.54
1:A:239:HIS:HD2	1:A:285:SER:OG	1.91	0.53
1:A:100:GLN:HG3	6:A:2096:HOH:O	2.10	0.51
1:C:251:LEU:O	1:C:255:GLU:HG2	2.12	0.50
1:A:79:HIS:CE1	1:A:157:ALA:H	2.31	0.49
1:A:209:PHE:HA	1:A:210:PRO:C	2.37	0.49
1:C:101:ASP:OD2	5:C:1383:EPE:H71	2.13	0.49
1:A:135:TYR:OH	1:A:309:HIS:HD2	1.97	0.48
1:A:135:TYR:OH	1:A:309:HIS:CD2	2.67	0.47
1:C:255:GLU:HG3	6:C:2264:HOH:O	2.14	0.47
1:A:167:LYS:HE3	1:A:197:ASP:OD2	2.14	0.47
1:A:217:GLU:H	1:A:217:GLU:CD	2.22	0.47
1:C:223:ARG:NH2	6:C:2156:HOH:O	2.47	0.47
1:A:243:MET:SD	1:A:292:MET:HE1	2.55	0.46
1:C:209:PHE:HA	1:C:210:PRO:C	2.41	0.45
1:A:63:GLN:HE21	1:A:65:ARG:HD3	1.81	0.45
1:C:58:LEU:HD11	1:C:327:GLU:CG	2.47	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:79:HIS:HD2	1:A:147:PRO:O	2.00	0.45
1:A:4:ARG:NE	1:A:63:GLN:OE1	2.51	0.44
1:A:203:LEU:CD1	1:A:289:MET:HE1	2.44	0.43
1:C:282:HIS:HE1	6:C:2245:HOH:O	2.02	0.43
1:A:100:GLN:HA	1:A:100:GLN:OE1	2.19	0.43
1:C:205:GLN:HE22	1:C:278:ARG:N	2.06	0.42
1:C:243:MET:SD	1:C:292:MET:HE1	2.60	0.42
1:A:247:GLN:HG2	6:A:2191:HOH:O	2.19	0.41
1:A:175:VAL:O	1:A:198:VAL:HA	2.21	0.41
1:A:79:HIS:HE1	1:A:157:ALA:H	1.66	0.41
1:C:3:ARG:HD3	6:C:2224:HOH:O	2.19	0.41
1:A:309:HIS:HE1	6:A:2040:HOH:O	2.03	0.40
1:C:175:VAL:O	1:C:198:VAL:HA	2.21	0.40
1:A:36:ALA:HB1	1:C:36:ALA:HB1	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	368/379 (97%)	356 (97%)	11 (3%)	1 (0%)	36	22
1	C	374/379 (99%)	361 (96%)	11 (3%)	2 (0%)	24	12
All	All	742/758 (98%)	717 (97%)	22 (3%)	3 (0%)	30	16

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	37	ALA
1	C	37	ALA
1	C	3	ARG

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	282/288 (98%)	278 (99%)	4 (1%)	59	46
1	C	287/288 (100%)	281 (98%)	6 (2%)	47	30
All	All	569/576 (99%)	559 (98%)	10 (2%)	51	36

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	41	GLU
1	A	147	PRO
1	A	340	LEU
1	A	367	GLN
1	C	41	GLU
1	C	61	ARG
1	C	147	PRO
1	C	153	PHE
1	C	169	ARG
1	C	377	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	32	GLN
1	A	39	HIS
1	A	63	GLN
1	A	79	HIS
1	A	239	HIS
1	A	309	HIS
1	C	32	GLN
1	C	158	ASN
1	C	170	HIS
1	C	205	GLN
1	C	228	ASN
1	C	230	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	280	GLN
1	C	282	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 13 ligands modelled in this entry, 6 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	ACT	A	1375	2	3,3,3	0.92	0	3,3,3	1.14	0
5	EPE	A	1377	-	15,15,15	2.04	1 (6%)	19,20,20	0.98	1 (5%)
4	ACT	C	1381	2	3,3,3	0.89	0	3,3,3	1.67	1 (33%)
5	EPE	A	1376	-	15,15,15	1.73	1 (6%)	19,20,20	1.32	1 (5%)
5	EPE	C	1383	-	15,15,15	1.76	1 (6%)	19,20,20	1.36	3 (15%)
5	EPE	A	1378	-	15,15,15	1.93	1 (6%)	19,20,20	1.06	0
5	EPE	C	1382	-	15,15,15	1.77	1 (6%)	19,20,20	1.29	2 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EPE	A	1377	-	-	1/9/19/19	0/1/1/1
5	EPE	A	1376	-	-	5/9/19/19	0/1/1/1
5	EPE	C	1383	-	-	4/9/19/19	0/1/1/1
5	EPE	A	1378	-	-	7/9/19/19	0/1/1/1
5	EPE	C	1382	-	-	1/9/19/19	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	1377	EPE	C10-S	-7.54	1.67	1.77
5	A	1378	EPE	C10-S	-6.97	1.67	1.77
5	C	1382	EPE	C10-S	-6.28	1.68	1.77
5	C	1383	EPE	C10-S	-6.11	1.69	1.77
5	A	1376	EPE	C10-S	-5.96	1.69	1.77

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	1376	EPE	O2S-S-C10	4.59	113.67	106.73
5	C	1382	EPE	O2S-S-C10	3.21	111.57	106.73
5	C	1383	EPE	O3S-S-C10	2.80	111.47	106.00
5	C	1382	EPE	O3S-S-C10	2.34	110.59	106.00
5	C	1383	EPE	C2-C3-N4	2.30	115.28	110.65
5	A	1377	EPE	O1S-S-C10	2.20	110.05	106.73
4	C	1381	ACT	OXT-C-CH3	2.16	124.11	115.05
5	C	1383	EPE	C5-N4-C3	2.11	113.39	108.84

There are no chirality outliers.

All (18) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	1376	EPE	C9-C10-S-O1S
5	A	1376	EPE	C9-C10-S-O2S
5	A	1378	EPE	C10-C9-N1-C2
5	A	1378	EPE	C10-C9-N1-C6
5	A	1378	EPE	C9-C10-S-O2S
5	C	1383	EPE	C9-C10-S-O1S
5	A	1378	EPE	N4-C7-C8-O8
5	A	1378	EPE	C9-C10-S-O3S

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
5	C	1383	EPE	C9-C10-S-O3S
5	C	1382	EPE	N4-C7-C8-O8
5	A	1376	EPE	C9-C10-S-O3S
5	A	1378	EPE	S-C10-C9-N1
5	C	1383	EPE	C8-C7-N4-C5
5	A	1378	EPE	C9-C10-S-O1S
5	C	1383	EPE	C9-C10-S-O2S
5	A	1377	EPE	N4-C7-C8-O8
5	A	1376	EPE	C10-C9-N1-C2
5	A	1376	EPE	C10-C9-N1-C6

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	C	1383	EPE	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	370/379 (97%)	-0.45	3 (0%) 82 85	9, 14, 29, 57	0
1	C	376/379 (99%)	-0.46	3 (0%) 82 85	8, 14, 30, 59	0
All	All	746/758 (98%)	-0.45	6 (0%) 82 85	8, 14, 30, 59	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	37	ALA	4.1
1	C	37	ALA	3.9
1	C	2	THR	3.5
1	A	371	CYS	3.3
1	C	38	GLY	2.8
1	A	2	THR	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	EPE	C	1383	15/15	0.69	0.17	36,41,82,85	0
5	EPE	A	1378	15/15	0.78	0.14	33,39,56,59	0
5	EPE	A	1377	15/15	0.86	0.11	34,42,70,74	0
5	EPE	C	1382	15/15	0.92	0.12	20,36,47,48	0
4	ACT	A	1375	4/4	0.93	0.10	17,19,23,23	0
5	EPE	A	1376	15/15	0.93	0.13	20,34,47,50	0
4	ACT	C	1381	4/4	0.95	0.10	17,19,21,23	0
2	ZN	C	1378	1/1	1.00	0.02	10,10,10,10	1
3	K	A	1373	1/1	1.00	0.02	10,10,10,10	0
3	K	A	1374	1/1	1.00	0.01	11,11,11,11	0
3	K	C	1379	1/1	1.00	0.01	11,11,11,11	0
3	K	C	1380	1/1	1.00	0.04	10,10,10,10	0
2	ZN	A	1372	1/1	1.00	0.01	10,10,10,10	1

6.5 Other polymers [i](#)

There are no such residues in this entry.