



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 05:49 PM UTC

PDB ID : 8G0M / pdb_00008g0m
Title : Structure of complex between TV6.6 and CD98hc ECD
Authors : Kariolis, M.S.; Lexa, K.; Liao, N.P.D.; Srivastava, D.; Tran, H.; Wells, R.C.
Deposited on : 2023-01-31
Resolution : 2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1898 (2.26-2.26)
Clashscore	190562	2005 (2.26-2.26)
Ramachandran outliers	187476	1965 (2.26-2.26)
Sidechain outliers	187428	1966 (2.26-2.26)
RSRZ outliers	180081	1898 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	425	11% 87% 10% .
2	B	227	2% 85% 7% 9%
3	C	8	62% 38%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 10175 atoms, of which 4995 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 4F2 cell-surface antigen heavy chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	413	Total	C	H	N	O	S	86	0	0
			6412	2049	3194	549	615	5			


There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	631	GLU	-	expression tag	UNP P08195
A	632	ASN	-	expression tag	UNP P08195
A	633	LEU	-	expression tag	UNP P08195
A	634	TYR	-	expression tag	UNP P08195
A	635	PHE	-	expression tag	UNP P08195
A	636	GLN	-	expression tag	UNP P08195

- Molecule 2 is a protein called TV 6.6 Fc fragment.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	207	Total	C	H	N	O	S	52	0	0
			3339	1078	1663	282	311	5			

- Molecule 3 is an oligosaccharide called beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	C	8	Total	C	H	N	O		21	0	0
			186	54	90	3	39				

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	1	0
			10	2	6	2		
4	A	1	Total	C	H	O	1	0
			10	2	6	2		
4	B	1	Total	C	H	O	1	0
			10	2	6	2		
4	B	1	Total	C	H	O	1	0
			10	2	6	2		
4	B	1	Total	C	H	O	1	0
			10	2	6	2		

- Molecule 5 is TETRAETHYLENE GLYCOL (CCD ID: PG4) (formula: $C_8H_{18}O_5$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	H	O	1	0
			31	8	18	5		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	49	Total	O	0	0
			49	49		
6	B	108	Total	O	0	0
			108	108		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 4F2 cell-surface antigen heavy chain



- Molecule 2: TV 6.6 Fc fragment



- Molecule 3: beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	136.15Å 167.13Å 83.68Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.68 – 2.25 44.68 – 2.25	Depositor EDS
% Data completeness (in resolution range)	100.0 (44.68-2.25) 100.0 (44.68-2.25)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.15 (at 2.24Å)	Xtriage
Refinement program	REFMAC 5.8.0403	Depositor
R, R_{free}	0.204 , 0.245 0.205 , 0.244	Depositor DCC
R_{free} test set	2335 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å ²)	52.3	Xtriage
Anisotropy	0.088	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 37.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	10175	wwPDB-VP
Average B, all atoms (Å ²)	66.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.44% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, NAG, BMA, MAN, GAL, FUC, PG4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.55	0/3290	0.95	0/4461
2	B	0.58	0/1724	0.94	0/2350
All	All	0.56	0/5014	0.95	0/6811

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3218	3194	3184	24	0
2	B	1676	1663	1656	8	0
3	C	96	90	82	1	0
4	A	8	12	12	0	0
4	B	12	18	18	0	0
5	A	13	18	18	0	0
6	A	49	0	0	0	0
6	B	108	0	0	1	0
All	All	5180	4995	4970	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:309:THR:HG22	1:A:317:SER:HB3	1.79	0.65
2:B:434:ASN:ND2	6:B:602:HOH:O	2.41	0.53
1:A:268:GLN:O	1:A:269:LYS:C	2.53	0.52
2:B:259:VAL:HG23	2:B:308:VAL:HG11	1.93	0.51
1:A:251:LEU:HD13	1:A:259:LEU:HD21	1.92	0.50
1:A:597:LEU:C	1:A:597:LEU:HD12	2.38	0.49
1:A:411:THR:HB	1:A:414:HIS:CB	2.43	0.48
1:A:628:TYR:O	1:A:629:ALA:HB2	2.13	0.47
1:A:595:LEU:HA	1:A:624:LEU:HD13	1.95	0.47
1:A:348:ARG:HB3	1:A:437:GLN:OE1	2.14	0.47
1:A:218:TRP:CE3	1:A:303:ARG:HG3	2.50	0.47
1:A:496:PHE:CG	1:A:497:PRO:HD2	2.50	0.46
1:A:250:TYR:OH	1:A:471:GLU:OE2	2.27	0.46
1:A:225:TYR:CZ	1:A:468:TYR:HA	2.50	0.46
1:A:411:THR:HB	1:A:414:HIS:HB2	1.97	0.45
2:B:276:ASN:HB2	2:B:322:LYS:HB3	1.98	0.45
2:B:406:LEU:HD12	2:B:406:LEU:C	2.42	0.45
1:A:318:TRP:CD2	1:A:353:LEU:HD13	2.53	0.44
1:A:250:TYR:O	1:A:253:SER:OG	2.30	0.44
1:A:251:LEU:HD13	1:A:259:LEU:CD2	2.47	0.44
1:A:265:HIS:CE1	1:A:310:PRO:HA	2.52	0.44
1:A:230:LEU:HD21	1:A:261:LEU:HD11	2.01	0.43
2:B:243:PHE:CG	3:C:5:NAG:H5	2.54	0.42
1:A:458:PHE:O	1:A:530:ARG:HD3	2.19	0.42
1:A:322:GLN:O	1:A:326:VAL:HG23	2.20	0.42
1:A:564:LEU:HB3	1:A:624:LEU:HB2	2.02	0.41
1:A:335:GLU:O	1:A:339:GLN:HG2	2.20	0.41
2:B:365:LEU:HB3	2:B:441:LEU:HD23	2.03	0.41
1:A:305:ILE:HD11	1:A:375:LEU:HD23	2.03	0.40
2:B:252:MET:HB2	2:B:255:ARG:HD3	2.03	0.40
2:B:263:VAL:HB	2:B:302:VAL:CG1	2.51	0.40
1:A:294:LEU:HD23	1:A:304:VAL:HG21	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	409/425 (96%)	391 (96%)	17 (4%)	1 (0%)	43	50
2	B	205/227 (90%)	204 (100%)	1 (0%)	0	100	100
All	All	614/652 (94%)	595 (97%)	18 (3%)	1 (0%)	43	50

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	310	PRO

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	349/359 (97%)	348 (100%)	1 (0%)	86	90
2	B	193/209 (92%)	191 (99%)	2 (1%)	68	77
All	All	542/568 (95%)	539 (99%)	3 (1%)	78	84

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	234	GLN
2	B	340	LYS
2	B	400	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	236	HIS
1	A	346	GLN
1	A	386	GLN
1	A	448	GLN
1	A	542	HIS
2	B	347	GLN
2	B	362	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

8 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	C	1	3,2	14,14,15	0.41	0	17,19,21	0.64	0
3	NAG	C	2	3	14,14,15	0.41	0	17,19,21	0.37	0
3	BMA	C	3	3	11,11,12	0.56	0	15,15,17	0.61	0
3	MAN	C	4	3	11,11,12	0.81	1 (9%)	15,15,17	0.84	0
3	NAG	C	5	3	14,14,15	0.34	0	17,19,21	0.74	0
3	GAL	C	6	3	11,11,12	0.37	0	15,15,17	0.71	1 (6%)
3	MAN	C	7	3	11,11,12	0.31	0	15,15,17	0.54	0
3	FUC	C	8	3	10,10,11	0.25	0	14,14,16	0.49	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	1	3,2	-	2/6/23/26	0/1/1/1
3	NAG	C	2	3	-	0/6/23/26	0/1/1/1
3	BMA	C	3	3	-	0/2/19/22	0/1/1/1
3	MAN	C	4	3	-	0/2/19/22	0/1/1/1
3	NAG	C	5	3	-	0/6/23/26	0/1/1/1
3	GAL	C	6	3	-	0/2/19/22	0/1/1/1
3	MAN	C	7	3	-	0/2/19/22	0/1/1/1
3	FUC	C	8	3	-	-	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	4	MAN	C2-C3	-2.33	1.48	1.52

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	6	GAL	C1-O5-C5	2.02	114.90	112.19

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	C	1	NAG	C8-C7-N2-C2
3	C	1	NAG	O7-C7-N2-C2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	5	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	EDO	A	701	-	3,3,3	0.06	0	2,2,2	0.06	0
4	EDO	B	503	-	3,3,3	0.18	0	2,2,2	0.15	0
4	EDO	B	501	-	3,3,3	0.25	0	2,2,2	0.39	0
4	EDO	B	502	-	3,3,3	0.19	0	2,2,2	0.23	0
4	EDO	A	703	-	3,3,3	0.26	0	2,2,2	0.39	0
5	PG4	A	702	-	12,12,12	0.41	0	11,11,11	0.23	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	A	701	-	-	1/1/1/1	-
4	EDO	B	503	-	-	1/1/1/1	-
4	EDO	B	501	-	-	0/1/1/1	-
4	EDO	B	502	-	-	0/1/1/1	-
4	EDO	A	703	-	-	1/1/1/1	-
5	PG4	A	702	-	-	6/10/10/10	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	702	PG4	O3-C5-C6-O4
5	A	702	PG4	O4-C7-C8-O5
4	A	701	EDO	O1-C1-C2-O2
4	A	703	EDO	O1-C1-C2-O2
5	A	702	PG4	C6-C5-O3-C4
5	A	702	PG4	O1-C1-C2-O2
5	A	702	PG4	C5-C6-O4-C7
4	B	503	EDO	O1-C1-C2-O2
5	A	702	PG4	O2-C3-C4-O3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	413/425 (97%)	0.68	45 (10%) 10 10	40, 67, 120, 149	0
2	B	207/227 (91%)	-0.21	5 (2%) 59 61	30, 48, 81, 112	0
All	All	620/652 (95%)	0.38	50 (8%) 18 16	30, 60, 111, 149	0

All (50) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	253	ILE	5.3
1	A	323	VAL	4.6
1	A	587	ALA	4.4
1	A	629	ALA	4.3
1	A	580	ALA	4.2
1	A	406	SER	4.0
1	A	410	SER	3.8
2	B	444	SER	3.6
1	A	548	PRO	3.5
1	A	285	GLY	3.4
1	A	259	LEU	3.4
1	A	395	ASN	3.3
1	A	214	PRO	3.3
1	A	574	LEU	3.2
1	A	586	SER	3.2
1	A	391	LEU	3.2
1	A	585	ALA	3.1
2	B	296	TYR	3.1
1	A	264	ILE	3.1
1	A	363	TRP	3.0
1	A	326	VAL	2.9
1	A	321	THR	2.9
1	A	589	LEU	2.8
1	A	261	LEU	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	588	SER	2.7
1	A	275	THR	2.7
1	A	290	PHE	2.6
1	A	305	ILE	2.6
1	A	412	GLY	2.5
1	A	583	LEU	2.5
1	A	327	ALA	2.5
1	A	357	SER	2.4
1	A	544	PHE	2.4
2	B	238	PRO	2.4
1	A	278	LEU	2.4
1	A	241	LEU	2.3
1	A	272	VAL	2.3
1	A	545	SER	2.3
1	A	306	LEU	2.2
2	B	254	SER	2.2
1	A	582	ASP	2.2
1	A	414	HIS	2.2
1	A	350	ILE	2.2
1	A	336	PHE	2.2
1	A	356	ALA	2.2
1	A	329	LYS	2.1
1	A	270	ASP	2.1
1	A	320	SER	2.1
1	A	280	ILE	2.0
1	A	269	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MAN	C	7	11/12	0.76	0.15	59,76,91,91	4
3	FUC	C	8	10/11	0.87	0.13	67,77,91,91	3

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NAG	C	1	14/15	0.90	0.11	54,64,80,91	1
3	GAL	C	6	11/12	0.90	0.12	52,72,91,91	4
3	NAG	C	2	14/15	0.92	0.10	49,56,64,91	2
3	NAG	C	5	14/15	0.93	0.08	61,68,78,91	2
3	MAN	C	4	11/12	0.93	0.08	53,58,91,91	3
3	BMA	C	3	11/12	0.96	0.06	50,54,91,91	2

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	EDO	A	701	4/4	0.76	0.17	92,97,102,103	1
4	EDO	A	703	4/4	0.84	0.21	71,80,83,84	1
4	EDO	B	503	4/4	0.85	0.20	76,85,89,92	1
4	EDO	B	502	4/4	0.86	0.18	78,90,94,95	1
5	PG4	A	702	13/13	0.87	0.16	64,77,83,84	1
4	EDO	B	501	4/4	0.92	0.12	59,67,69,74	1

6.5 Other polymers [i](#)

There are no such residues in this entry.