



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 12:33 PM UTC

PDB ID : 5G1U / pdb_00005g1u
Title : Linalool Dehydratase Isomerase in complex with Geraniol
Authors : Chambers, S.; Hau, A.; Man, H.; Omar, M.; Turkenburg, J.P.; Grogan, G.
Deposited on : 2016-03-30
Resolution : 2.57 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.57 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4770 (2.60-2.56)
Clashscore	190562	5124 (2.60-2.56)
Ramachandran outliers	187476	5046 (2.60-2.56)
Sidechain outliers	187428	5046 (2.60-2.56)
RSRZ outliers	180081	4770 (2.60-2.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	372	% 87% 8% ..
1	B	372	 88% 8% ..
1	C	372	% 87% 8% ..
1	D	372	 87% 8% ..
1	E	372	% 87% 9% ..

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 15138 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called LINALOOL DEHYDRATASE/ISOMERASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	362	Total	C	N	O	S	Se	0	0	0
			2857	1850	473	521	4	9			
1	B	364	Total	C	N	O	S	Se	0	0	0
			2874	1858	479	524	4	9			
1	C	363	Total	C	N	O	S	Se	0	0	0
			2862	1853	472	524	4	9			
1	D	362	Total	C	N	O	S	Se	0	0	0
			2867	1854	479	521	4	9			
1	E	362	Total	C	N	O	S	Se	0	0	0
			2862	1853	476	520	4	9			

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	-	expression tag	UNP E1XUJ2
B	1	MSE	-	expression tag	UNP E1XUJ2
C	1	MSE	-	expression tag	UNP E1XUJ2
D	1	MSE	-	expression tag	UNP E1XUJ2
E	1	MSE	-	expression tag	UNP E1XUJ2

- Molecule 2 is Geraniol (CCD ID: 64Z) (formula: C₁₀H₁₈O).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			11	10	1		
2	B	1	Total	C	O	0	0
			11	10	1		
2	C	1	Total	C	O	0	0
			11	10	1		
2	D	1	Total	C	O	0	0
			11	10	1		
2	E	1	Total	C	O	0	0
			11	10	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	152	Total	O	0	0
			152	152		
3	B	180	Total	O	0	0
			180	180		
3	C	160	Total	O	0	0
			160	160		
3	D	146	Total	O	0	0
			146	146		
3	E	123	Total	O	0	0
			123	123		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: LINALOOL DEHYDRATASE/ISOMERASE



- Molecule 1: LINALOOL DEHYDRATASE/ISOMERASE




- Molecule 1: LINALOOL DEHYDRATASE/ISOMERASE



- Molecule 1: LINALOOL DEHYDRATASE/ISOMERASE





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	99.80Å 106.04Å 220.51Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	90.92 – 2.57 90.92 – 2.57	Depositor EDS
% Data completeness (in resolution range)	100.0 (90.92-2.57) 100.0 (90.92-2.57)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.58 (at 2.27Å)	Xtriage
Refinement program	REFMAC 5.8.0131	Depositor
R, R_{free}	0.161 , 0.198 (Not available) , 0.177	Depositor DCC
R_{free} test set	5571 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	36.1	Xtriage
Anisotropy	0.597	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 27.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	15138	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 64Z

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.85	0/2934	0.99	6/3983 (0.2%)
1	B	0.87	1/2951 (0.0%)	1.03	8/4005 (0.2%)
1	C	0.88	2/2939 (0.1%)	0.98	7/3990 (0.2%)
1	D	0.83	0/2944	0.97	7/3995 (0.2%)
1	E	0.83	0/2939	0.98	7/3988 (0.2%)
All	All	0.85	3/14707 (0.0%)	0.99	35/19961 (0.2%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	363	PRO	CA-C	5.50	1.54	1.51
1	C	121	LYS	CA-C	-5.50	1.45	1.52
1	C	363	PRO	CA-C	5.07	1.54	1.51

The worst 5 of 35 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	212	LYS	N-CA-C	-12.09	96.65	111.40
1	B	213	ASP	N-CA-C	8.83	124.75	113.88
1	A	121	LYS	N-CA-C	-8.15	95.20	109.06
1	E	276	ARG	CG-CD-NE	7.23	127.92	112.00
1	C	121	LYS	N-CA-C	-7.18	95.50	110.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2857	0	2722	23	0
1	B	2874	0	2746	16	0
1	C	2862	0	2724	21	0
1	D	2867	0	2740	18	0
1	E	2862	0	2733	14	0
2	A	11	0	0	0	0
2	B	11	0	0	0	0
2	C	11	0	0	0	0
2	D	11	0	0	0	0
2	E	11	0	0	0	0
3	A	152	0	0	5	0
3	B	180	0	0	3	0
3	C	160	0	0	5	0
3	D	146	0	0	0	0
3	E	123	0	0	1	0
All	All	15138	0	13665	89	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 89 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:193:ARG:NH2	1:A:362:MSE:O	2.23	0.72
1:B:125:MSE:CE	3:B:2048:HOH:O	2.41	0.69
1:D:193:ARG:NH2	1:D:362:MSE:O	2.26	0.68
1:E:193:ARG:NH2	1:E:362:MSE:O	2.27	0.68
1:A:124:ILE:HG21	1:A:169:ILE:HD13	1.74	0.68

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	360/372 (97%)	354 (98%)	4 (1%)	2 (1%)	21	39
1	B	362/372 (97%)	354 (98%)	7 (2%)	1 (0%)	36	56
1	C	361/372 (97%)	352 (98%)	7 (2%)	2 (1%)	21	39
1	D	360/372 (97%)	353 (98%)	6 (2%)	1 (0%)	36	56
1	E	360/372 (97%)	352 (98%)	7 (2%)	1 (0%)	36	56
All	All	1803/1860 (97%)	1765 (98%)	31 (2%)	7 (0%)	30	49

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	120	GLU
1	A	212	LYS
1	B	3	GLU
1	C	120	GLU
1	C	212	LYS

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	287/293 (98%)	276 (96%)	11 (4%)	29	54
1	B	290/293 (99%)	278 (96%)	12 (4%)	27	51
1	C	288/293 (98%)	277 (96%)	11 (4%)	29	54
1	D	289/293 (99%)	278 (96%)	11 (4%)	29	54

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	E	287/293 (98%)	278 (97%)	9 (3%)	35 60
All	All	1441/1465 (98%)	1387 (96%)	54 (4%)	30 55

5 of 54 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	119	ILE
1	D	47	ARG
1	E	103	LYS
1	C	124	ILE
1	C	214	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 31 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	36	ASN
1	E	36	ASN
1	C	250	HIS
1	E	227	HIS
1	D	250	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	64Z	A	1366	-	10,10,10	3.82	3 (30%)	11,11,11	1.34	1 (9%)
2	64Z	C	1366	-	10,10,10	3.98	4 (40%)	11,11,11	1.07	1 (9%)
2	64Z	E	1366	-	10,10,10	4.16	5 (50%)	11,11,11	1.22	2 (18%)
2	64Z	D	1366	-	10,10,10	4.02	4 (40%)	11,11,11	1.19	1 (9%)
2	64Z	B	1366	-	10,10,10	3.80	3 (30%)	11,11,11	1.28	2 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	64Z	A	1366	-	-	4/9/9/9	-
2	64Z	C	1366	-	-	2/9/9/9	-
2	64Z	E	1366	-	-	4/9/9/9	-
2	64Z	D	1366	-	-	4/9/9/9	-
2	64Z	B	1366	-	-	2/9/9/9	-

The worst 5 of 19 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	1366	64Z	C6-C5	9.84	1.55	1.33
2	E	1366	64Z	C6-C5	9.68	1.55	1.33
2	D	1366	64Z	C6-C5	9.50	1.54	1.33
2	B	1366	64Z	C6-C5	9.36	1.54	1.33
2	A	1366	64Z	C6-C5	9.06	1.53	1.33

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1366	64Z	C-C1-C9	2.75	120.91	114.59
2	B	1366	64Z	C-C1-C9	2.47	120.28	114.59
2	D	1366	64Z	C-C1-C9	2.38	120.07	114.59
2	C	1366	64Z	C-C1-C9	2.28	119.84	114.59

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	1366	64Z	C-C1-C9	2.26	119.79	114.59

There are no chirality outliers.

5 of 16 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	1366	64Z	C2-C3-C4-C5
2	E	1366	64Z	C2-C3-C4-C5
2	C	1366	64Z	C3-C4-C5-C6
2	C	1366	64Z	C3-C4-C5-C8
2	A	1366	64Z	C2-C3-C4-C5

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	353/372 (94%)	-0.37	2 (0%) 85 84	29, 40, 59, 75	0
1	B	355/372 (95%)	-0.42	1 (0%) 90 89	27, 37, 58, 87	0
1	C	354/372 (95%)	-0.37	5 (1%) 73 70	29, 38, 61, 89	0
1	D	353/372 (94%)	-0.40	1 (0%) 90 89	28, 39, 59, 71	0
1	E	353/372 (94%)	-0.36	3 (0%) 82 81	29, 40, 59, 79	0
All	All	1768/1860 (95%)	-0.38	12 (0%) 84 82	27, 39, 59, 89	0

The worst 5 of 12 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	2	ALA	4.7
1	C	120	GLU	3.1
1	A	4	LEU	3.0
1	A	365	PRO	3.0
1	C	3	GLU	2.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	64Z	D	1366	11/11	0.91	0.20	55,64,71,71	0
2	64Z	E	1366	11/11	0.91	0.18	53,58,61,61	0
2	64Z	B	1366	11/11	0.92	0.17	58,63,67,69	0
2	64Z	A	1366	11/11	0.93	0.17	58,63,67,70	0
2	64Z	C	1366	11/11	0.94	0.13	52,56,61,61	0

6.5 Other polymers [i](#)

There are no such residues in this entry.