



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 24, 2026 – 03:08 PM UTC

PDB ID : 8G2S / pdb_00008g2s
Title : Horse liver alcohol dehydrogenase His-51-Gln form complexed with NAD⁺ and capric acid
Authors : Plapp, B.V.
Deposited on : 2023-02-06
Resolution : 1.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1756 (1.46-1.46)
Clashscore	190562	1795 (1.46-1.46)
Ramachandran outliers	187476	1776 (1.46-1.46)
Sidechain outliers	187428	1776 (1.46-1.46)
RSRZ outliers	180081	1756 (1.46-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	374	95% 5%
1	B	374	97% .

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 6737 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alcohol dehydrogenase E chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	374	Total	C	N	O	S	0	26	0
			2891	1839	482	543	27			
1	B	374	Total	C	N	O	S	0	31	0
			2917	1853	486	551	27			


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	51	GLN	HIS	engineered mutation	UNP P00327
B	51	GLN	HIS	engineered mutation	UNP P00327

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Zn	0	0
			2	2		
2	B	2	Total	Zn	0	0
			2	2		

- Molecule 3 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (ACIDIC FORM) (CCD ID: NAJ) (formula: C₂₁H₂₇N₇O₁₄P₂) (labeled as "Ligand of Interest" by depositor).



- # DKA
-
- The image displays the chemical structure of DKA (Decanoic acid). It features a zigzag carbon chain with 10 carbon atoms, labeled C1 through C10. The carboxylic acid group at the right end is highlighted in red, with the carbonyl oxygen labeled O1 and the hydroxyl oxygen labeled O2.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	1
			24	20	4		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	1
			24	20	4		

- Molecule 5 is (4S)-2-METHYL-2,4-PENTANEDIOL (CCD ID: MPD) (formula: C₆H₁₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			8	6	2		
5	A	1	Total	C	O	0	1
			8	6	2		

- Molecule 6 is (4R)-2-METHYLPENTANE-2,4-DIOL (CCD ID: MRD) (formula: C₆H₁₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	1
			8	6	2		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	399	Total	O	0	18
			407	407		
7	B	352	Total	O	0	19
			358	358		

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Alcohol dehydrogenase E chain

Chain A:  95% 5%



- Molecule 1: Alcohol dehydrogenase E chain

Chain B:  97% .



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	44.24Å 51.25Å 92.46Å 92.01° 102.87° 109.91°	Depositor
Resolution (Å)	19.98 – 1.45 19.98 – 1.45	Depositor EDS
% Data completeness (in resolution range)	87.4 (19.98-1.45) 87.4 (19.98-1.45)	Depositor EDS
R_{merge}	0.03	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.23 (at 1.45Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.156 , 0.189 0.156 , 0.189	Depositor DCC
R_{free} test set	2272 reflections (1.73%)	wwPDB-VP
Wilson B-factor (Å ²)	14.9	Xtriage
Anisotropy	0.472	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 37.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.36$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6737	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.08% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MPD, ZN, DKA, NAJ, MRD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.08	2/3053 (0.1%)	1.14	1/4117 (0.0%)
1	B	1.04	1/3089 (0.0%)	1.15	0/4164
All	All	1.06	3/6142 (0.0%)	1.15	1/8281 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	201	GLY	C-O	5.38	1.28	1.23
1	A	139	HIS	CE1-NE2	5.35	1.37	1.32
1	B	32	LYS	C-O	5.34	1.29	1.23

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	326	ASP	CA-CB-CG	5.20	117.80	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2891	0	2955	10	0
1	B	2917	0	2977	7	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	44	0	26	0	0
3	B	44	0	26	0	0
4	A	24	0	38	0	0
4	B	24	0	38	1	0
5	A	16	0	28	1	0
6	A	8	0	14	1	0
7	A	407	0	0	0	0
7	B	358	0	0	1	0
All	All	6737	0	6102	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:256[A]:GLU:OE1	7:B:501:HOH:O	2.10	0.70
5:A:405:MPD:H52	5:A:405:MPD:HM2	1.85	0.57
1:B:40[A]:MET:HE1	1:B:150:THR:HG22	1.94	0.49
1:A:40[B]:MET:HE1	1:A:150:THR:HG22	1.95	0.48
1:B:40[A]:MET:HE1	1:B:150:THR:CG2	2.46	0.45
1:A:118[B]:MET:HE2	1:A:118[B]:MET:HB2	1.68	0.45
1:A:70:ALA:HB1	1:A:166:LEU:HD22	1.98	0.44
1:B:350:LEU:O	1:B:372:LEU:HA	2.17	0.44
1:A:282[B]:CYS:HB3	1:A:287:GLY:HA3	2.00	0.43
1:A:306:MET:SD	4:B:404[B]:DKA:H82	2.59	0.42
1:A:350:LEU:O	1:A:372:LEU:HA	2.19	0.42
1:A:218:ARG:CZ	6:A:407[B]:MRD:H5C3	2.49	0.42
1:A:100:CYS:HB2	1:A:112:LEU:HD12	2.02	0.41
1:B:282[B]:CYS:HB3	1:B:287:GLY:HA3	2.02	0.41
1:A:40[B]:MET:HE1	1:A:150:THR:CG2	2.50	0.41
1:A:40[B]:MET:HE2	1:A:40[B]:MET:HB2	1.89	0.41
1:B:290:VAL:HA	1:B:315[A]:LYS:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	398/374 (106%)	388 (98%)	10 (2%)	0	100	100
1	B	403/374 (108%)	394 (98%)	9 (2%)	0	100	100
All	All	801/748 (107%)	782 (98%)	19 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	333/308 (108%)	331 (99%)	2 (1%)	78	58
1	B	338/308 (110%)	337 (100%)	1 (0%)	86	74
All	All	671/616 (109%)	668 (100%)	3 (0%)	86	70

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	239[A]	GLU
1	A	239[B]	GLU
1	B	268	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	34	HIS
1	A	300	ASN
1	B	34	HIS
1	B	138	HIS
1	B	259	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 4 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	NAJ	B	403	-	46,48,48	1.85	8 (17%)	64,73,73	2.75	15 (23%)
5	MPD	A	405	-	7,7,7	0.32	0	9,10,10	0.68	0
4	DKA	A	404[A]	2	11,11,11	0.95	1 (9%)	11,11,11	1.16	1 (9%)
4	DKA	A	404[B]	2	11,11,11	0.56	0	11,11,11	1.39	1 (9%)
4	DKA	B	404[B]	2	11,11,11	0.56	0	11,11,11	1.06	1 (9%)
4	DKA	B	404[A]	2	11,11,11	0.86	1 (9%)	11,11,11	1.00	0
3	NAJ	A	403	-	46,48,48	1.62	8 (17%)	64,73,73	1.95	12 (18%)
5	MPD	A	406[A]	-	7,7,7	0.19	0	9,10,10	0.69	0
6	MRD	A	407[B]	-	7,7,7	0.15	0	9,10,10	0.46	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAJ	B	403	-	-	3/30/62/62	0/5/5/5
5	MPD	A	405	-	-	2/5/5/5	-
4	DKA	A	404[A]	2	-	4/9/9/9	-
4	DKA	A	404[B]	2	-	5/9/9/9	-
4	DKA	B	404[B]	2	-	6/9/9/9	-
4	DKA	B	404[A]	2	-	0/9/9/9	-
3	NAJ	A	403	-	-	5/30/62/62	0/5/5/5
5	MPD	A	406[A]	-	-	1/5/5/5	-
6	MRD	A	407[B]	-	-	0/5/5/5	-

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	403	NAJ	C2N-N1N	5.64	1.41	1.35
3	B	403	NAJ	C6A-N1A	-5.34	1.24	1.35
3	A	403	NAJ	C2A-N3A	5.11	1.45	1.33
3	B	403	NAJ	C2N-C3N	-5.08	1.30	1.39
3	B	403	NAJ	C2A-N3A	3.83	1.42	1.33
3	A	403	NAJ	C2N-C3N	3.83	1.45	1.39
3	A	403	NAJ	C6A-N1A	-3.75	1.27	1.35
3	A	403	NAJ	PN-O3	3.32	1.63	1.59
3	B	403	NAJ	C6N-N1N	2.95	1.42	1.35
3	A	403	NAJ	C6N-N1N	2.60	1.41	1.35
3	A	403	NAJ	PA-O2A	-2.31	1.44	1.55
3	B	403	NAJ	C3N-C7N	2.30	1.54	1.50
3	A	403	NAJ	O2D-C2D	2.28	1.48	1.43
4	A	404[A]	DKA	O2-C1	-2.25	1.23	1.30
3	A	403	NAJ	C8A-N9A	2.21	1.42	1.37
3	B	403	NAJ	C6N-C5N	-2.16	1.34	1.38
4	B	404[A]	DKA	O2-C1	-2.13	1.23	1.30
3	B	403	NAJ	C4N-C3N	2.05	1.42	1.39

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	403	NAJ	C2N-C3N-C4N	10.48	130.44	118.26
3	B	403	NAJ	C5N-C4N-C3N	-9.75	110.78	120.36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	403	NAJ	C6N-N1N-C2N	-9.34	113.93	121.88
3	A	403	NAJ	N1A-C2A-N3A	-6.89	115.91	128.84
3	A	403	NAJ	C2A-N1A-C6A	6.60	128.15	118.55
3	B	403	NAJ	N1A-C2A-N3A	-5.89	117.79	128.84
3	A	403	NAJ	C8A-N7A-C5A	4.70	112.63	104.26
3	A	403	NAJ	N9A-C8A-N7A	-4.39	105.25	113.40
3	B	403	NAJ	C5A-C4A-N3A	-4.05	121.31	127.27
3	B	403	NAJ	C2A-N3A-C4A	4.02	118.06	111.14
3	B	403	NAJ	C2A-N1A-C6A	3.99	124.35	118.55
3	B	403	NAJ	C5N-C6N-N1N	3.79	125.55	120.38
3	A	403	NAJ	N6A-C6A-N1A	3.55	123.65	118.72
3	A	403	NAJ	C6A-C5A-C4A	-3.45	112.96	117.03
3	B	403	NAJ	C4N-C3N-C7N	-3.40	111.81	121.06
4	A	404[B]	DKA	C3-C2-C1	-3.30	105.89	114.51
3	B	403	NAJ	N9A-C8A-N7A	-3.29	107.30	113.40
3	B	403	NAJ	O7N-C7N-C3N	3.25	123.58	119.60
3	B	403	NAJ	C6N-N1N-C1D	2.85	125.33	119.73
3	A	403	NAJ	C8A-N9A-C4A	2.72	111.13	106.03
4	A	404[A]	DKA	C3-C2-C1	-2.72	107.41	114.51
3	A	403	NAJ	C2D-C3D-C4D	-2.62	97.55	102.61
3	B	403	NAJ	O7N-C7N-N7N	-2.52	118.97	122.62
3	A	403	NAJ	O2N-PN-O1N	2.48	123.99	112.44
3	B	403	NAJ	O4B-C1B-C2B	-2.40	101.47	106.62
3	B	403	NAJ	O2N-PN-O1N	2.39	123.57	112.44
3	A	403	NAJ	O7N-C7N-C3N	-2.36	116.71	119.60
3	A	403	NAJ	C6N-N1N-C2N	-2.27	119.94	121.88
4	B	404[B]	DKA	C3-C2-C1	-2.25	108.64	114.51
3	A	403	NAJ	O3-PA-O1A	-2.08	104.44	110.70

There are no chirality outliers.

All (26) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	403	NAJ	O4D-C1D-N1N-C2N
3	A	403	NAJ	O4D-C1D-N1N-C6N
3	A	403	NAJ	C2D-C1D-N1N-C2N
3	A	403	NAJ	C2D-C1D-N1N-C6N
3	B	403	NAJ	O4D-C1D-N1N-C2N
3	B	403	NAJ	O4D-C1D-N1N-C6N
3	B	403	NAJ	C2D-C1D-N1N-C2N
4	B	404[B]	DKA	C4-C5-C6-C7
4	A	404[B]	DKA	C1-C2-C3-C4

Continued on next page...

Continued from previous page...



Mol	Chain	Res	Type	Atoms
4	A	404[B]	DKA	C2-C3-C4-C5
4	B	404[B]	DKA	C3-C4-C5-C6
4	B	404[B]	DKA	C1-C2-C3-C4
4	A	404[A]	DKA	C2-C3-C4-C5
4	A	404[A]	DKA	C3-C4-C5-C6
5	A	405	MPD	O2-C2-C3-C4
5	A	406[A]	MPD	O2-C2-C3-C4
5	A	405	MPD	C2-C3-C4-C5
4	B	404[B]	DKA	O2-C1-C2-C3
4	A	404[B]	DKA	O1-C1-C2-C3
4	A	404[B]	DKA	O2-C1-C2-C3
4	B	404[B]	DKA	C5-C6-C7-C8
4	A	404[A]	DKA	C7-C8-C9-C10
4	B	404[B]	DKA	O1-C1-C2-C3
4	A	404[B]	DKA	C6-C7-C8-C9
4	A	404[A]	DKA	C6-C7-C8-C9
3	A	403	NAJ	O4B-C4B-C5B-O5B




There are no ring outliers.


3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	405	MPD	1	0
4	B	404[B]	DKA	1	0
6	A	407[B]	MRD	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	374/374 (100%)	-0.44	1 (0%) 90 92	6, 12, 20, 47	26 (6%)
1	B	374/374 (100%)	-0.32	1 (0%) 90 92	5, 14, 23, 39	28 (7%)
All	All	748/748 (100%)	-0.38	2 (0%) 90 92	5, 13, 22, 47	54 (7%)

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1	SER	3.2
1	B	1	SER	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	MPD	A	405	8/8	0.82	0.18	26,29,33,33	8
5	MPD	A	406[A]	8/8	0.83	0.13	28,30,31,31	8
6	MRD	A	407[B]	8/8	0.86	0.11	18,19,19,19	8

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	DKA	A	404[B]	12/12	0.92	0.10	10,13,14,15	12
4	DKA	A	404[A]	12/12	0.92	0.10	13,19,28,28	12
4	DKA	B	404[B]	12/12	0.94	0.08	13,17,22,22	12
4	DKA	B	404[A]	12/12	0.94	0.08	14,17,19,22	12
3	NAJ	A	403	44/44	0.98	0.04	7,9,14,21	0
2	ZN	B	402	1/1	0.98	0.03	13,13,13,13	0
3	NAJ	B	403	44/44	0.99	0.04	9,12,18,22	0
2	ZN	A	402	1/1	1.00	0.01	11,11,11,11	0
2	ZN	B	401	1/1	1.00	0.04	13,13,13,13	0
2	ZN	A	401	1/1	1.00	0.05	11,11,11,11	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around DKA A 404 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around DKA B 404 (B):**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around DKA B 404 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around NAJ A 403:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.