



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 07:28 AM UTC

PDB ID : 6G3B / pdb_00006g3b
Title : AvaII restriction endonuclease in complex with an RNA/DNA hybrid
Authors : Kisiala, M.; Kowalska, M.; Czapinska, H.; Bochtler, M.
Deposited on : 2018-03-24
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)
RNA backbone	3983	1011 (2.20-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	238	3% 89% 5% 5%
1	B	238	3% 90% 5% 5%
2	C	11	9% 82% 18%
3	D	11	9% 64% 36%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4906 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Type II site-specific deoxyribonuclease.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	225	Total	C	N	O	S	0	9	0
			1867	1188	319	354	6			
1	B	225	Total	C	N	O	S	0	7	0
			1854	1179	316	353	6			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP Q8YYB7
A	2	GLY	-	expression tag	UNP Q8YYB7
A	231	LEU	-	expression tag	UNP Q8YYB7
A	232	GLU	-	expression tag	UNP Q8YYB7
A	233	HIS	-	expression tag	UNP Q8YYB7
A	234	HIS	-	expression tag	UNP Q8YYB7
A	235	HIS	-	expression tag	UNP Q8YYB7
A	236	HIS	-	expression tag	UNP Q8YYB7
A	237	HIS	-	expression tag	UNP Q8YYB7
A	238	HIS	-	expression tag	UNP Q8YYB7
B	1	MET	-	initiating methionine	UNP Q8YYB7
B	2	GLY	-	expression tag	UNP Q8YYB7
B	231	LEU	-	expression tag	UNP Q8YYB7
B	232	GLU	-	expression tag	UNP Q8YYB7
B	233	HIS	-	expression tag	UNP Q8YYB7
B	234	HIS	-	expression tag	UNP Q8YYB7
B	235	HIS	-	expression tag	UNP Q8YYB7
B	236	HIS	-	expression tag	UNP Q8YYB7
B	237	HIS	-	expression tag	UNP Q8YYB7
B	238	HIS	-	expression tag	UNP Q8YYB7

- Molecule 2 is a RNA chain called RNA (5'-R(P*GP*UP*AP*GP*GP*AP*CP*CP*AP*UP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	11	Total 470	C 212	N 90	O 148	P 20	0	11	0

- Molecule 3 is a DNA chain called DNA (5'-D(*CP*CP*AP*TP*GP*GP*TP*CP*CP*TP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	11	Total 438	C 212	N 76	O 130	P 20	0	11	0

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	126	Total 126	O 126	0	1
4	B	143	Total 143	O 143	0	4
4	C	3	Total 3	O 3	0	0
4	D	5	Total 5	O 5	0	1

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Type II site-specific deoxyribonuclease

Chain A: 



- Molecule 1: Type II site-specific deoxyribonuclease

Chain B: 



- Molecule 2: RNA (5'-R(P*GP*UP*AP*GP*GP*AP*CP*CP*AP*UP*G)-3')

Chain C: 



- Molecule 3: DNA (5'-D(*CP*CP*AP*TP*GP*GP*TP*CP*CP*TP*A)-3')

Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	37.68Å 104.05Å 78.29Å 90.00° 93.27° 90.00°	Depositor
Resolution (Å)	52.02 – 1.80 52.02 – 1.80	Depositor EDS
% Data completeness (in resolution range)	95.4 (52.02-1.80) 95.4 (52.02-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.45 (at 1.79Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.168 , 0.202 0.176 , 0.208	Depositor DCC
R_{free} test set	2665 reflections (4.80%)	wwPDB-VP
Wilson B-factor (Å ²)	33.8	Xtriage
Anisotropy	0.132	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 56.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4906	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.20% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.83	0/1910	0.90	1/2576 (0.0%)
1	B	0.89	0/1894	0.95	2/2559 (0.1%)
2	C	0.09	0/526	0.22	0/818
3	D	0.14	0/488	0.37	0/748
All	All	0.77	0/4818	0.82	3/6701 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	140	GLY	N-CA-C	-5.97	101.46	112.62
1	B	81	ILE	CA-C-N	-5.43	114.65	120.03
1	B	81	ILE	C-N-CA	-5.43	114.65	120.03

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1867	0	1860	7	0
1	B	1854	0	1836	5	0
2	C	470	0	212	3	0
3	D	438	0	239	2	0
4	A	126	0	0	2	0
4	B	143	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	C	3	0	0	0	0
4	D	5	0	0	0	0
All	All	4906	0	4147	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:75:PRO:HG3	1:A:117:ILE:HD13	1.87	0.56
2:C:2[B]:U:O2'	2:C:3[B]:A:OP2	2.31	0.49
1:A:68:LYS:NZ	4:A:305:HOH:O	2.45	0.48
2:C:2[A]:U:O2'	2:C:3[A]:A:OP1	2.31	0.48
3:D:5[B]:DG:H2'	3:D:6[B]:DG:C8	2.50	0.45
1:A:207:GLU:H	1:A:207:GLU:CD	2.25	0.45
1:B:135:TRP:CD2	1:B:150:TYR:HB2	2.53	0.44
1:A:73:ARG:N	4:A:301:HOH:O	2.44	0.44
1:B:76:LEU:HD12	1:B:77:PRO:HD2	2.00	0.44
1:A:166:ASP:HB3	1:A:186:PHE:CE1	2.53	0.43
1:B:204:GLU:H	1:B:204:GLU:CD	2.25	0.43
1:B:142:MET:HG3	1:B:143:VAL:HG23	2.00	0.43
2:C:2[A]:U:O2'	2:C:3[A]:A:P	2.77	0.43
1:A:74:THR:HG22	1:A:229[B]:LYS:HE3	2.01	0.42
1:B:64:THR:HG23	1:B:68:LYS:HE2	2.02	0.42
1:A:135:TRP:CD2	1:A:150:TYR:HB2	2.54	0.42
3:D:3[B]:DA:H2'	3:D:4[B]:DT:C6	2.55	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	230/238 (97%)	226 (98%)	4 (2%)	0	100	100
1	B	228/238 (96%)	220 (96%)	8 (4%)	0	100	100
All	All	458/476 (96%)	446 (97%)	12 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	205/208 (99%)	205 (100%)	0	100	100
1	B	203/208 (98%)	202 (100%)	1 (0%)	81	80
All	All	408/416 (98%)	407 (100%)	1 (0%)	86	87

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	165	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	7	GLN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	C	0/11	-	-

There are no RNA backbone outliers to report.

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2			OWAB(Å ²)	Q < 0.9
1	A	225/238 (94%)	0.32	8 (3%)	46	46	15, 42, 77, 112	9 (4%)
1	B	225/238 (94%)	0.23	6 (2%)	56	56	16, 40, 75, 106	7 (3%)
2	C	11/11 (100%)	1.58	1 (9%)	15	13	46, 51, 62, 63	11 (100%)
3	D	11/11 (100%)	1.27	1 (9%)	15	13	46, 51, 62, 63	11 (100%)
All	All	472/498 (94%)	0.33	16 (3%)	48	48	15, 42, 77, 112	38 (8%)

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	D	1[A]	DC	6.2
2	C	11[A]	G	6.1
1	B	174	ALA	4.0
1	B	175	ALA	2.9
1	B	169	GLU	2.7
1	B	45	LEU	2.7
1	B	176	ILE	2.5
1	A	138	CYS	2.4
1	A	169	GLU	2.4
1	B	172	SER	2.3
1	A	76	LEU	2.3
1	A	170	ASN	2.2
1	A	229[A]	LYS	2.2
1	A	2	GLY	2.1
1	A	175	ALA	2.1
1	A	231	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.