



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 19, 2026 – 02:52 AM UTC

PDB ID : 6G3X / pdb_00006g3x
Title : Native Structure of the mouse 8-oxoguanine DNA Glycosylase mOGG1
Authors : Masuyer, G.; Helleday, T.; Stenmark, P.
Deposited on : 2018-03-26
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	318	 7% 92% 6% •
1	B	318	 12% 94% 5% •
1	C	318	 11% 93% 5% •

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 7824 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called N-glycosylase/DNA lyase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	313	Total	C	N	O	S	5	0	0
			2493	1586	450	446	11			
1	B	313	Total	C	N	O	S	8	0	0
			2487	1582	449	445	11			
1	C	312	Total	C	N	O	S	4	0	0
			2473	1575	444	443	11			

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	8	GLY	-	expression tag	UNP O08760
A	9	SER	-	expression tag	UNP O08760
A	10	HIS	-	expression tag	UNP O08760
B	8	GLY	-	expression tag	UNP O08760
B	9	SER	-	expression tag	UNP O08760
B	10	HIS	-	expression tag	UNP O08760
C	8	GLY	-	expression tag	UNP O08760
C	9	SER	-	expression tag	UNP O08760
C	10	HIS	-	expression tag	UNP O08760

- Molecule 2 is NICKEL (II) ION (CCD ID: NI) (formula: Ni).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Ni	0	0
			1	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	139	Total	O	0	0
			139	139		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	122	Total 122	O 122	0	0
3	C	109	Total 109	O 109	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: N-glycosylase/DNA lyase



- Molecule 1: N-glycosylase/DNA lyase



- Molecule 1: N-glycosylase/DNA lyase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	80.77Å 81.21Å 167.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	83.77 – 2.10 83.77 – 2.10	Depositor EDS
% Data completeness (in resolution range)	98.9 (83.77-2.10) 98.9 (83.77-2.10)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.20 (at 2.07Å)	Xtriage
Refinement program	REFMAC 5.8.0218	Depositor
R, R_{free}	0.221 , 0.274 0.228 , 0.275	Depositor DCC
R_{free} test set	3351 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å ²)	36.1	Xtriage
Anisotropy	0.234	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 34.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.015 for k,h,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7824	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.78% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.52	0/2563	0.75	0/3488
1	B	0.50	0/2557	0.74	0/3480
1	C	0.49	0/2541	0.73	0/3456
All	All	0.50	0/7661	0.74	0/10424

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2493	0	2436	12	0
1	B	2487	0	2428	9	0
1	C	2473	0	2412	9	0
2	A	1	0	0	0	0
3	A	139	0	0	1	0
3	B	122	0	0	0	0
3	C	109	0	0	0	0
All	All	7824	0	7276	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:266:PRO:HD2	1:A:315:GLN:HE22	1.40	0.84
1:B:266:PRO:HD2	1:B:315:GLN:HE22	1.42	0.81
1:A:249:LYS:HD3	1:A:268:ASP:HB3	1.69	0.75
1:C:82:ASP:HA	1:C:85:VAL:HG13	1.71	0.71
1:C:266:PRO:HD2	1:C:315:GLN:HE22	1.59	0.67
1:B:266:PRO:HD2	1:B:315:GLN:NE2	2.15	0.61
1:C:147:SER:HB3	1:C:155:ILE:HG12	1.86	0.58
1:A:266:PRO:HD2	1:A:315:GLN:NE2	2.16	0.53
1:A:59:LEU:HD21	1:A:99:TYR:CZ	2.45	0.51
1:C:11:MET:SD	1:C:81:ASP:HA	2.51	0.50
1:C:151:ASN:HD22	1:C:154:ARG:HB2	1.76	0.50
1:B:14:ARG:NH2	1:B:85:VAL:HG13	2.27	0.49
1:B:46:ARG:HG3	1:B:139:GLU:OE2	2.12	0.49
1:B:148:SER:O	1:B:204:ARG:NH1	2.45	0.49
1:A:103:ASP:HB2	3:A:614:HOH:O	2.14	0.47
1:A:283:PRO:HB2	1:A:285:THR:HG22	1.96	0.46
1:A:148:SER:O	1:A:204:ARG:NH1	2.49	0.46
1:B:34:ARG:HA	1:B:68:GLN:HE22	1.81	0.46
1:A:320:SER:O	1:A:324:ARG:HG2	2.16	0.45
1:C:214:ALA:O	1:C:218:GLU:HB3	2.16	0.45
1:C:148:SER:O	1:C:150:ASN:N	2.51	0.43
1:B:34:ARG:NH2	1:B:36:ASP:OD2	2.50	0.43
1:A:106:LEU:HG	1:A:110:TYR:CE1	2.55	0.42
1:A:35:LEU:H	1:A:68:GLN:HE22	1.67	0.42
1:C:266:PRO:HD2	1:C:315:GLN:NE2	2.29	0.42
1:B:27:PRO:HA	1:B:72:GLN:OE1	2.20	0.41
1:A:150:ASN:OD1	1:A:154:ARG:HB3	2.20	0.41
1:B:51:SER:O	1:B:52:PRO:C	2.64	0.41
1:C:92:GLU:O	1:C:95:THR:HB	2.21	0.41
1:A:35:LEU:H	1:A:68:GLN:NE2	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	309/318 (97%)	295 (96%)	14 (4%)	0	100	100
1	B	309/318 (97%)	299 (97%)	10 (3%)	0	100	100
1	C	304/318 (96%)	295 (97%)	8 (3%)	1 (0%)	36	36
All	All	922/954 (97%)	889 (96%)	32 (4%)	1 (0%)	48	50

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	81	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	263/266 (99%)	262 (100%)	1 (0%)	84	89
1	B	262/266 (98%)	262 (100%)	0	100	100
1	C	260/266 (98%)	259 (100%)	1 (0%)	84	89
All	All	785/798 (98%)	783 (100%)	2 (0%)	86	91

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	287	GLN
1	C	96	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	68	GLN
1	A	72	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	119	HIS
1	A	227	GLN
1	A	315	GLN
1	B	68	GLN
1	B	315	GLN
1	C	68	GLN
1	C	72	GLN
1	C	150	ASN
1	C	151	ASN
1	C	164	GLN
1	C	179	HIS
1	C	315	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	313/318 (98%)	0.40	21 (6%) 24 25	24, 37, 64, 112	1 (0%)
1	B	313/318 (98%)	0.79	39 (12%) 8 8	27, 42, 71, 90	2 (0%)
1	C	312/318 (98%)	0.90	36 (11%) 9 10	31, 46, 76, 102	1 (0%)
All	All	938/954 (98%)	0.70	96 (10%) 12 12	24, 41, 72, 112	4 (0%)

All (96) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	288	ALA	6.0
1	B	93	LEU	5.2
1	B	16	LEU	5.0
1	C	323	LEU	5.0
1	A	285	THR	4.8
1	B	90	LEU	4.6
1	B	22	LEU	4.6
1	B	21	ALA	4.2
1	B	323	LEU	4.2
1	B	290	GLY	4.0
1	A	82	ASP	4.0
1	C	85	VAL	3.8
1	B	80	GLY	3.8
1	B	87	ARG	3.8
1	B	285	THR	3.7
1	B	24	ALA	3.7
1	B	23	TRP	3.5
1	B	88	PRO	3.4
1	C	245	GLY	3.4
1	C	81	ASP	3.4
1	B	15	THR	3.4
1	A	85	VAL	3.4
1	B	82	ASP	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	86	SER	3.2
1	A	284	LYS	3.1
1	B	91	GLU	3.1
1	C	80	GLY	3.0
1	A	288	ALA	3.0
1	B	89	THR	3.0
1	C	218	GLU	3.0
1	A	11	MET	3.0
1	B	8	GLY	2.9
1	A	10	HIS	2.9
1	C	199	LEU	2.8
1	A	287	GLN	2.8
1	C	90	LEU	2.8
1	C	195	HIS	2.8
1	B	27	PRO	2.8
1	C	190	PRO	2.8
1	B	74	TYR	2.7
1	A	291	PRO	2.7
1	C	142	PHE	2.7
1	B	85	VAL	2.7
1	B	123	VAL	2.7
1	C	221	GLY	2.7
1	A	274	ILE	2.7
1	A	80	GLY	2.6
1	C	185	HIS	2.6
1	B	20	PRO	2.6
1	B	19	SER	2.6
1	B	97	HIS	2.6
1	B	76	THR	2.6
1	C	224	TRP	2.6
1	C	285	THR	2.6
1	B	196	LEU	2.6
1	B	319	PHE	2.6
1	B	75	CYS	2.6
1	B	25	SER	2.6
1	A	81	ASP	2.5
1	A	41	SER	2.5
1	A	270	HIS	2.5
1	C	319	PHE	2.5
1	A	273	GLN	2.4
1	B	194	THR	2.4
1	C	18	SER	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	17	SER	2.4
1	C	214	ALA	2.4
1	A	185	HIS	2.4
1	B	324	ARG	2.4
1	C	286	SER	2.4
1	A	20	PRO	2.4
1	A	40	ALA	2.3
1	C	222	PRO	2.3
1	C	216	LEU	2.3
1	A	122	ARG	2.3
1	A	126	LYS	2.3
1	B	71	ASP	2.3
1	C	291	PRO	2.2
1	C	215	ILE	2.2
1	C	219	GLN	2.2
1	C	82	ASP	2.2
1	C	146	CYS	2.2
1	C	270	HIS	2.1
1	B	103	ASP	2.1
1	C	71	ASP	2.1
1	B	291	PRO	2.1
1	B	77	VAL	2.1
1	C	184	LEU	2.1
1	C	217	GLU	2.1
1	B	122	ARG	2.1
1	C	234	GLU	2.0
1	A	294	LEU	2.0
1	C	20	PRO	2.0
1	C	147	SER	2.0
1	C	186	ALA	2.0
1	C	196	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NI	A	401	1/1	1.00	0.01	31,31,31,31	0

6.5 Other polymers

There are no such residues in this entry.