



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 18, 2026 – 08:35 AM UTC

PDB ID : 7G3W / pdb_00007g3w
Title : Crystal Structure of rat Autotaxin in complex with N-[(2R)-5-chloro-2,3-dihydro-1H-inden-2-yl]-5-(3-methyl-1,2,4-oxadiazol-5-yl)pyrimidin-2-amine, i.e. SMILES c1c(cnc(n1)N[C@H]1Cc2cc(Cl)ccc2C1)C1=NC(=NO1)C with IC50=0.186827 microM
Authors : Stihle, M.; Benz, J.; Hunziker, D.; Pinard, E.; Rudolph, M.G.
Deposited on : 2023-06-05
Resolution : 1.51 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.51 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5890 (1.54-1.50)
Clashscore	190562	6116 (1.54-1.50)
Ramachandran outliers	187476	6002 (1.54-1.50)
Sidechain outliers	187428	5999 (1.54-1.50)
RSRZ outliers	180081	5891 (1.54-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	846	 6% 85% 10% • 5%
2	B	8	 100%

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 7481 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Isoform 2 of Ectonucleotide pyrophosphatase/phosphodiesterase family member 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	805	Total	C	N	O	S	0	23	0
			6630	4214	1137	1226	53			

There are 14 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	53	ALA	ASN	engineered mutation	UNP Q64610
A	410	ALA	ASN	engineered mutation	UNP Q64610
A	591	THR	ARG	engineered mutation	UNP Q64610
A	863	GLY	-	expression tag	UNP Q64610
A	864	GLY	-	expression tag	UNP Q64610
A	865	ARG	-	expression tag	UNP Q64610
A	866	HIS	-	expression tag	UNP Q64610
A	867	HIS	-	expression tag	UNP Q64610
A	868	HIS	-	expression tag	UNP Q64610
A	869	HIS	-	expression tag	UNP Q64610
A	870	HIS	-	expression tag	UNP Q64610
A	871	HIS	-	expression tag	UNP Q64610
A	872	HIS	-	expression tag	UNP Q64610
A	873	HIS	-	expression tag	UNP Q64610

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	8	Total	C	N	O	0	0	0
			94	52	2	40			

- Molecule 3 is N-[(2S)-5-chloro-2,3-dihydro-1H-inden-2-yl]-5-(3-methyl-1,2,4-oxadiazol-5-yl)pyrimidin-2-amine (CCD ID: Y9V) (formula: C₁₆H₁₄ClN₅O) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	Cl	N	O	0	0
			23	16	1	5	1		

- Molecule 4 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	2	Total Ca	0	0
			2 2		

- Molecule 5 is POTASSIUM ION (CCD ID: K) (formula: K).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total K	0	0
			1 1		

- Molecule 6 is ACETATE ION (CCD ID: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 7 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	Zn	0	0
			1	1		

- Molecule 8 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	1	Total	Na	0	0
			1	1		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	719	Total	O	0	2
			721	721		


3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Isoform 2 of Ectonucleotide pyrophosphatase/phosphodiesterase family member 2



- Molecule 2: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	83.92Å 91.92Å 119.61Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	72.89 – 1.51 72.89 – 1.51	Depositor EDS
% Data completeness (in resolution range)	81.9 (72.89-1.51) 69.2 (72.89-1.51)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.38 (at 1.52Å)	Xtriage
Refinement program	REFMAC 5.8.0232	Depositor
R, R_{free}	0.171 , 0.207 (Not available) , 0.203	Depositor DCC
R_{free} test set	6013 reflections (4.17%)	wwPDB-VP
Wilson B-factor (Å ²)	18.0	Xtriage
Anisotropy	0.007	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 27.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7481	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.42% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, ZN, BMA, MAN, ACT, NAG, Y9V, CA, K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.19	18/6874 (0.3%)	1.30	19/9315 (0.2%)

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	595	HIS	CE1-NE2	7.81	1.40	1.32
1	A	546	GLU	CD-OE1	7.75	1.40	1.25
1	A	539	PHE	C-O	7.45	1.33	1.24
1	A	487	PHE	C-O	6.60	1.31	1.23
1	A	427	HIS	CE1-NE2	6.55	1.39	1.32
1	A	515	LEU	C-O	5.88	1.31	1.24
1	A	374	ASN	C-O	5.69	1.31	1.24
1	A	412	THR	C-O	5.65	1.30	1.24
1	A	343	LYS	C-O	5.59	1.30	1.24
1	A	330	GLU	CD-OE1	5.50	1.35	1.25
1	A	169	SER	CA-CB	-5.46	1.45	1.53
1	A	538	THR	C-O	5.36	1.31	1.23
1	A	390	GLY	CA-C	-5.32	1.47	1.52
1	A	433	HIS	CE1-NE2	5.27	1.37	1.32
1	A	554	GLY	C-O	-5.26	1.19	1.24
1	A	561	GLU	C-O	-5.13	1.17	1.24
1	A	236	VAL	C-O	5.13	1.29	1.24
1	A	534	LEU	C-O	5.10	1.30	1.24

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	827	ARG	CG-CD-NE	-9.72	90.62	112.00
1	A	546	GLU	CB-CG-CD	7.53	125.40	112.60
1	A	377	THR	CB-CA-C	6.39	121.40	110.86

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	50	PRO	N-CA-CB	6.25	109.87	103.00
1	A	540	ARG	CB-CA-C	6.15	117.05	110.65
1	A	272	THR	CB-CA-C	5.90	119.33	109.89
1	A	377	THR	CA-C-N	5.84	132.70	121.54
1	A	377	THR	C-N-CA	5.84	132.70	121.54
1	A	364	VAL	CA-C-O	-5.80	114.11	120.43
1	A	681	PHE	CA-CB-CG	-5.73	108.07	113.80
1	A	237[A]	PHE	CA-CB-CG	5.73	119.53	113.80
1	A	237[B]	PHE	CA-CB-CG	5.73	119.53	113.80
1	A	273	PHE	CA-CB-CG	5.54	119.34	113.80
1	A	822	HIS	CA-CB-CG	-5.40	108.40	113.80
1	A	515	LEU	CA-C-O	-5.35	114.06	120.10
1	A	202	ARG	CA-C-O	5.26	123.98	119.66
1	A	749	GLU	CB-CA-C	-5.14	101.13	110.63
1	A	271	GLY	N-CA-C	-5.08	106.75	112.33
1	A	659	SER	N-CA-CB	5.06	117.67	110.03

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6630	0	6427	40	0
2	B	94	0	79	0	0
3	A	23	0	0	0	0
4	A	2	0	0	0	0
5	A	1	0	0	0	0
6	A	8	0	6	1	0
7	A	1	0	0	0	0
8	A	1	0	0	0	0
9	A	721	0	0	7	0
All	All	7481	0	6512	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:479[A]:LYS:NZ	9:A:1005:HOH:O	2.32	0.61
1:A:594:ARG:HH12	1:A:636:SER:HB3	1.64	0.61
1:A:539:PHE:O	1:A:541:PRO:HD3	2.01	0.60
6:A:905:ACT:H1	9:A:1045:HOH:O	2.04	0.56
1:A:600[B]:ARG:HH21	1:A:600[B]:ARG:HG3	1.72	0.54
1:A:371:PHE:CE1	1:A:457:PRO:HA	2.44	0.53
1:A:222:PRO:HA	1:A:225:HIS:CE1	2.45	0.52
1:A:230:ASN:ND2	9:A:1001:HOH:O	2.18	0.50
1:A:463:LYS:C	9:A:1526:HOH:O	2.55	0.49
1:A:401:LYS:HA	1:A:401:LYS:HE3	1.95	0.49
1:A:171[A]:ASP:OD2	1:A:306:TYR:OH	2.28	0.48
1:A:201:MET:HE1	1:A:356:VAL:CG2	2.44	0.47
1:A:576:ASN:HD22	1:A:579:GLU:CD	2.23	0.47
1:A:594:ARG:NH1	1:A:636:SER:HB3	2.30	0.47
1:A:163:PRO:HB3	1:A:350:CYS:O	2.15	0.47
1:A:365:THR:OG1	1:A:367:ASP:OD1	2.27	0.46
1:A:856:HIS:CD2	1:A:858[A]:TYR:CE1	3.03	0.46
1:A:508[A]:LEU:HD23	1:A:508[A]:LEU:HA	1.79	0.46
1:A:166:ILE:HD12	1:A:351:VAL:HG11	1.98	0.45
1:A:470:PHE:C	1:A:470:PHE:CD1	2.94	0.45
1:A:600[B]:ARG:CG	1:A:600[B]:ARG:NH2	2.77	0.45
1:A:269:ARG:HB3	1:A:300:PRO:HA	1.97	0.45
1:A:468:CYS:SG	1:A:469:PHE:N	2.90	0.45
1:A:600[B]:ARG:HG3	1:A:600[B]:ARG:NH2	2.31	0.45
1:A:183:LYS:HD3	9:A:1363:HOH:O	2.18	0.44
1:A:269:ARG:CZ	1:A:269:ARG:HB2	2.47	0.44
1:A:366:CYS:HB2	1:A:468:CYS:HB2	1.88	0.44
1:A:584:ARG:O	1:A:588:LYS:HG2	2.18	0.44
1:A:600[B]:ARG:HD2	1:A:611:ILE:HD11	1.99	0.43
1:A:148[A]:CYS:SG	1:A:190:LYS:HE3	2.59	0.43
1:A:205:TYR:CD1	1:A:206:PRO:HA	2.53	0.43
1:A:237[B]:PHE:CD2	1:A:252:ARG:NH1	2.87	0.43
1:A:450[A]:ARG:NH1	9:A:1004:HOH:O	2.32	0.43
1:A:550:PRO:HB2	1:A:611:ILE:HG12	2.01	0.42
1:A:199:PRO:HG3	1:A:499[B]:LYS:HE3	2.01	0.42
1:A:272:THR:HB	9:A:1225:HOH:O	2.19	0.42
1:A:600[B]:ARG:HH21	1:A:600[B]:ARG:CG	2.32	0.42
1:A:403:ASP:HA	1:A:404:PRO:HD2	1.95	0.41
1:A:414:LYS:HA	1:A:414:LYS:HD3	1.83	0.41
1:A:449:ARG:O	1:A:450[A]:ARG:HB2	2.21	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:458:LEU:C	1:A:458:LEU:HD12	2.46	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	822/846 (97%)	800 (97%)	21 (3%)	1 (0%)	48 25

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	378	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	750/765 (98%)	742 (99%)	8 (1%)	65 40

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	379	VAL
1	A	389	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	397	ILE
1	A	401	LYS
1	A	469	PHE
1	A	549	ARG
1	A	580	GLU
1	A	594	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	398	ASN
1	A	436	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

8 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	B	1	2,1	14,14,15	1.36	1 (7%)	17,19,21	1.19	2 (11%)
2	NAG	B	2	2	14,14,15	1.45	3 (21%)	17,19,21	3.00	6 (35%)
2	BMA	B	3	2	11,11,12	1.21	2 (18%)	15,15,17	2.10	5 (33%)
2	MAN	B	4	2	11,11,12	1.33	1 (9%)	15,15,17	2.08	3 (20%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	MAN	B	5	2	11,11,12	0.97	0	15,15,17	1.48	1 (6%)
2	MAN	B	6	2	11,11,12	1.31	2 (18%)	15,15,17	1.06	1 (6%)
2	MAN	B	7	2	11,11,12	1.16	2 (18%)	15,15,17	1.65	4 (26%)
2	MAN	B	8	2	11,11,12	1.61	3 (27%)	15,15,17	1.87	3 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	1/6/23/26	0/1/1/1
2	BMA	B	3	2	-	0/2/19/22	0/1/1/1
2	MAN	B	4	2	-	0/2/19/22	0/1/1/1
2	MAN	B	5	2	-	1/2/19/22	0/1/1/1
2	MAN	B	6	2	-	0/2/19/22	0/1/1/1
2	MAN	B	7	2	-	1/2/19/22	0/1/1/1
2	MAN	B	8	2	-	0/2/19/22	0/1/1/1

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	8	MAN	C2-C3	3.43	1.57	1.52
2	B	2	NAG	O6-C6	3.28	1.56	1.42
2	B	1	NAG	O5-C1	2.92	1.48	1.43
2	B	8	MAN	O3-C3	2.57	1.49	1.43
2	B	3	BMA	O3-C3	2.47	1.49	1.43
2	B	2	NAG	C8-C7	2.45	1.55	1.50
2	B	7	MAN	C1-C2	2.43	1.58	1.52
2	B	2	NAG	C6-C5	2.37	1.59	1.51
2	B	3	BMA	C2-C3	2.34	1.56	1.52
2	B	6	MAN	O5-C1	2.31	1.47	1.43
2	B	7	MAN	O2-C2	2.20	1.48	1.43
2	B	4	MAN	C2-C3	2.17	1.55	1.52
2	B	8	MAN	C1-C2	2.15	1.57	1.52
2	B	6	MAN	C4-C5	2.02	1.57	1.53

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2	NAG	C1-O5-C5	8.56	123.66	112.19
2	B	4	MAN	C1-O5-C5	6.49	120.88	112.19
2	B	8	MAN	O3-C3-C2	5.57	121.42	110.05
2	B	3	BMA	O2-C2-C1	-4.63	98.63	109.22
2	B	2	NAG	C3-C4-C5	4.45	118.30	110.23
2	B	2	NAG	O7-C7-C8	-4.17	114.62	122.05
2	B	2	NAG	O7-C7-N2	3.29	127.79	121.98
2	B	3	BMA	O2-C2-C3	3.26	116.90	110.15
2	B	4	MAN	O3-C3-C2	3.10	116.38	110.05
2	B	2	NAG	C2-N2-C7	3.05	126.99	122.90
2	B	7	MAN	C1-O5-C5	2.97	116.17	112.19
2	B	1	NAG	O5-C1-C2	-2.88	106.84	111.29
2	B	3	BMA	C1-C2-C3	2.86	113.81	109.64
2	B	7	MAN	O2-C2-C1	2.83	115.70	109.22
2	B	3	BMA	O4-C4-C5	-2.81	102.41	109.32
2	B	1	NAG	O4-C4-C5	-2.47	103.23	109.32
2	B	4	MAN	O5-C5-C6	2.46	112.45	107.66
2	B	8	MAN	O2-C2-C3	2.33	114.98	110.15
2	B	8	MAN	C2-C3-C4	-2.32	106.79	110.86
2	B	5	MAN	C3-C4-C5	2.28	114.36	110.23
2	B	3	BMA	O3-C3-C4	2.27	115.73	110.38
2	B	7	MAN	O2-C2-C3	-2.19	105.61	110.15
2	B	7	MAN	O4-C4-C3	-2.11	105.39	110.38
2	B	6	MAN	O5-C5-C6	2.09	111.74	107.66
2	B	2	NAG	O4-C4-C3	-2.01	105.64	110.38

There are no chirality outliers.


All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	7	MAN	O5-C5-C6-O6
2	B	2	NAG	O5-C5-C6-O6
2	B	5	MAN	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 5 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	ACT	A	904	-	3,3,3	1.48	0	3,3,3	0.62	0
6	ACT	A	905	-	3,3,3	1.09	0	3,3,3	0.68	0
3	Y9V	A	901	-	26,26,26	3.40	14 (53%)	36,37,37	2.34	14 (38%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	Y9V	A	901	-	-	0/8/16/16	0/4/4/4

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	901	Y9V	C21-N22	8.14	1.51	1.37
3	A	901	Y9V	C5-N4	7.75	1.45	1.34
3	A	901	Y9V	C11-C10	-5.36	1.31	1.39
3	A	901	Y9V	C5-N6	4.50	1.40	1.34
3	A	901	Y9V	C14-C12	-4.37	1.30	1.38
3	A	901	Y9V	C17-C8	4.28	1.60	1.53
3	A	901	Y9V	O19-C18	4.27	1.38	1.34
3	A	901	Y9V	C3-C2	3.80	1.45	1.39
3	A	901	Y9V	C3-N4	-3.30	1.27	1.34
3	A	901	Y9V	C11-C12	2.90	1.43	1.38
3	A	901	Y9V	C1-C2	-2.41	1.35	1.39
3	A	901	Y9V	C21-N20	2.40	1.35	1.29
3	A	901	Y9V	C18-N22	2.31	1.33	1.30
3	A	901	Y9V	C17-C16	-2.19	1.46	1.50

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	901	Y9V	C18-O19-N20	6.96	110.62	106.08
3	A	901	Y9V	N22-C21-N20	-5.49	109.89	114.91
3	A	901	Y9V	C16-C17-C8	-3.83	99.11	102.89
3	A	901	Y9V	O19-C18-N22	-3.29	110.52	113.29
3	A	901	Y9V	C21-N22-C18	3.20	104.21	102.90
3	A	901	Y9V	C17-C16-C10	3.04	112.54	110.50
3	A	901	Y9V	C3-N4-C5	-2.68	112.14	115.81
3	A	901	Y9V	C9-C10-C16	2.64	112.27	110.50
3	A	901	Y9V	C10-C9-C8	-2.62	100.31	102.89
3	A	901	Y9V	C23-C21-N20	2.40	125.74	121.71
3	A	901	Y9V	C3-C2-C1	2.39	118.27	115.36
3	A	901	Y9V	C2-C1-N6	-2.22	120.32	123.59
3	A	901	Y9V	N6-C5-N4	-2.22	124.18	126.43
3	A	901	Y9V	C2-C3-N4	2.19	126.82	123.59

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	905	ACT	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	805/846 (95%)	0.30	52 (6%)	25 29	8, 20, 48, 107	23 (2%)

All (52) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	469	PHE	7.8
1	A	572	VAL	6.7
1	A	578	LEU	6.3
1	A	68	VAL	5.5
1	A	581	LEU	5.1
1	A	468	CYS	4.9
1	A	71	PRO	4.5
1	A	397	ILE	4.2
1	A	51	TRP	4.1
1	A	50	PRO	3.9
1	A	243	LEU	3.8
1	A	860	SER	3.6
1	A	249	PHE	3.3
1	A	463	LYS	3.2
1	A	70	PRO	3.2
1	A	777	PHE	3.1
1	A	573	GLU	3.0
1	A	571	LYS	2.9
1	A	110	VAL	2.8
1	A	148[A]	CYS	2.8
1	A	210	PHE	2.8
1	A	576	ASN	2.8
1	A	577	LYS	2.7
1	A	416	PRO	2.7
1	A	366	CYS	2.7
1	A	585	LEU	2.6
1	A	377	THR	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	246	ARG	2.6
1	A	104	LYS	2.6
1	A	53	ALA	2.6
1	A	460	VAL	2.6
1	A	242	HIS	2.5
1	A	470	PHE	2.5
1	A	462	LYS	2.4
1	A	594	ARG	2.4
1	A	161	VAL	2.3
1	A	55	SER	2.3
1	A	582	ASN	2.3
1	A	251	HIS	2.3
1	A	461	TYR	2.3
1	A	537	ASN	2.3
1	A	667	LEU	2.3
1	A	54	THR	2.2
1	A	245	GLY	2.2
1	A	274	PHE	2.2
1	A	244	ARG	2.1
1	A	69	GLY	2.1
1	A	402	TYR	2.1
1	A	83	SER	2.1
1	A	583	LYS	2.0
1	A	400	SER	2.0
1	A	52	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	MAN	B	7	11/12	0.55	0.18	81,89,97,108	0
2	BMA	B	3	11/12	0.66	0.16	48,57,63,66	0
2	MAN	B	8	11/12	0.78	0.17	32,69,82,87	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	MAN	B	4	11/12	0.79	0.15	39,48,52,54	0
2	MAN	B	5	11/12	0.84	0.17	32,44,53,65	0
2	NAG	B	2	14/15	0.90	0.12	21,28,38,50	0
2	MAN	B	6	11/12	0.92	0.14	28,37,42,42	0
2	NAG	B	1	14/15	0.96	0.06	12,14,17,20	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	ACT	A	904	4/4	0.92	0.13	22,25,29,31	0
3	Y9V	A	901	23/23	0.95	0.10	18,24,60,62	0
6	ACT	A	905	4/4	0.97	0.08	23,25,31,38	0
5	K	A	903	1/1	0.98	0.05	21,21,21,21	0
8	NA	A	907	1/1	0.98	0.08	25,25,25,25	0
4	CA	A	902	1/1	0.99	0.07	21,21,21,21	0
7	ZN	A	906	1/1	1.00	0.02	15,15,15,15	0
4	CA	A	908	1/1	1.00	0.02	13,13,13,13	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.