



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 07:45 AM UTC

PDB ID : 5G4Q / pdb_00005g4q
Title : H.pylori Beta clamp in complex with 5-chloroisatin
Authors : Pandey, P.; Gourinath, S.
Deposited on : 2016-05-16
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	374	4% 84% 12% ..
1	B	374	8% 83% 15% ..

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 5643 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA POLYMERASE III SUBUNIT BETA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	364	Total	C	N	O	S	0	0	0
			2763	1779	438	535	11			
1	B	369	Total	C	N	O	S	0	0	0
			2811	1810	445	545	11			

- Molecule 2 is 5-chloro-1H-indole-2,3-dione (CCD ID: 2HQ) (formula: C₈H₄ClNO₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	Cl	N	O	0	0
			12	8	1	1	2		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	23	Total	O	0	0
			23	23		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	34	Total	O	0	0
			34	34		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: DNA POLYMERASE III SUBUNIT BETA



• Molecule 1: DNA POLYMERASE III SUBUNIT BETA



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	82.31Å 66.37Å 89.12Å 90.00° 115.96° 90.00°	Depositor
Resolution (Å)	80.00 – 2.30 80.00 – 2.30	Depositor EDS
% Data completeness (in resolution range)	91.1 (80.00-2.30) 91.1 (80.00-2.30)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.03 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5	Depositor
R, R_{free}	0.227 , 0.285 0.257 , 0.296	Depositor DCC
R_{free} test set	1896 reflections (4.77%)	wwPDB-VP
Wilson B-factor (Å ²)	47.0	Xtriage
Anisotropy	0.401	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 28.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.000 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5643	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 86.02 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.0628e-07. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 2HQ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.05	0/2809	1.07	4/3800 (0.1%)
1	B	1.06	2/2858 (0.1%)	1.08	6/3866 (0.2%)
All	All	1.06	2/5667 (0.0%)	1.08	10/7666 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	329	LEU	C-O	-6.25	1.16	1.24
1	B	338	VAL	N-CA	-5.29	1.39	1.46

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	220	ASP	N-CA-C	-7.36	101.06	110.19
1	B	10	LEU	N-CA-C	-6.35	104.28	111.14
1	A	231	HIS	N-CA-C	6.17	118.55	109.24
1	B	362	ASN	N-CA-C	6.05	118.90	109.52
1	B	156	GLY	N-CA-C	5.68	119.58	111.12
1	B	231	HIS	N-CA-C	5.61	118.36	109.50
1	B	346	SER	CA-C-N	5.48	126.69	119.84
1	B	346	SER	C-N-CA	5.48	126.69	119.84
1	A	362	ASN	N-CA-C	5.44	117.67	109.41
1	A	315	ALA	N-CA-C	5.13	117.91	110.42

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2763	0	2712	22	0
1	B	2811	0	2774	23	0
2	B	12	0	4	0	0
3	A	23	0	0	0	0
3	B	34	0	0	0	0
All	All	5643	0	5490	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:350:ILE:HD12	1:B:369:MET:HE1	1.61	0.83
1:A:350:ILE:HD12	1:A:369:MET:HE1	1.64	0.79
1:A:70:ILE:HD11	1:A:107:LEU:HD12	1.69	0.74
1:A:70:ILE:HD12	1:A:96:LEU:HD23	1.70	0.72
1:B:70:ILE:HD11	1:B:107:LEU:HD12	1.80	0.64
1:B:341:CYS:SG	1:B:342:ASN:N	2.70	0.64
1:A:158:LEU:HD22	1:A:242:TYR:CD2	2.37	0.59
1:B:70:ILE:HD12	1:B:96:LEU:HD23	1.85	0.57
1:B:263:GLU:CD	1:B:304:VAL:HG21	2.29	0.57
1:A:120:VAL:HG21	1:B:120:VAL:CG2	2.35	0.56
1:A:138:PHE:HB3	1:A:204:LEU:HD11	1.89	0.53
1:B:270:LEU:C	1:B:270:LEU:HD23	2.33	0.53
1:A:281:THR:HB	1:A:288:LEU:HD23	1.91	0.52
1:A:120:VAL:HG11	1:B:120:VAL:HG21	1.91	0.52
1:B:256:SER:OG	1:B:340:ARG:NH1	2.42	0.51
1:A:219:SER:OG	1:A:220:ASP:O	2.21	0.51
1:A:350:ILE:CD1	1:A:369:MET:HE1	2.40	0.50
1:A:120:VAL:HG21	1:B:120:VAL:HG21	1.94	0.50
1:B:75:PHE:O	1:B:79:ILE:HG12	2.11	0.50
1:B:157:ILE:HG22	1:B:199:LEU:HB2	1.95	0.49
1:A:270:LEU:C	1:A:270:LEU:HD23	2.38	0.49
1:A:1:MET:CE	1:A:67:VAL:C	2.87	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:5:VAL:HG21	1:A:10:LEU:HD22	1.95	0.48
1:A:3:ILE:HG22	1:A:62:SER:HA	1.96	0.47
1:B:350:ILE:HD12	1:B:369:MET:CE	2.39	0.47
1:A:278:ILE:HD12	1:A:322:ALA:HB2	1.97	0.47
1:A:158:LEU:O	1:A:170:VAL:HA	2.15	0.46
1:B:63:ASP:N	1:B:63:ASP:OD1	2.49	0.46
1:B:3:ILE:HG22	1:B:62:SER:HA	1.97	0.46
1:B:70:ILE:HD12	1:B:96:LEU:CD2	2.48	0.44
1:A:120:VAL:HG21	1:B:120:VAL:HG22	1.99	0.44
1:B:163:GLN:O	1:B:164:LYS:C	2.61	0.44
1:B:142:ALA:HB3	1:B:143:PRO:HD3	2.01	0.43
1:B:158:LEU:O	1:B:170:VAL:HA	2.20	0.42
1:B:138:PHE:HB3	1:B:204:LEU:CD1	2.51	0.41
1:A:79:ILE:HD12	1:A:87:ILE:HD13	2.02	0.41
1:A:32:HIS:CE1	1:A:110:PHE:CD2	3.09	0.41
1:A:126:SER:HB2	1:A:218:LYS:HG2	2.01	0.41
1:B:296:HIS:O	1:B:297:SER:CB	2.69	0.41
1:B:47:SER:OG	1:B:48:ASP:N	2.54	0.41
1:A:268:ILE:HG23	1:A:278:ILE:CD1	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	358/374 (96%)	332 (93%)	21 (6%)	5 (1%)	9	9
1	B	365/374 (98%)	344 (94%)	21 (6%)	0	100	100
All	All	723/748 (97%)	676 (94%)	42 (6%)	5 (1%)	18	23

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	85	SER
1	A	93	ASP
1	A	240	GLY
1	A	63	ASP
1	A	92	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	300/342 (88%)	289 (96%)	11 (4%)	30	45
1	B	307/342 (90%)	287 (94%)	20 (6%)	15	22
All	All	607/684 (89%)	576 (95%)	31 (5%)	21	32

All (31) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	62	SER
1	A	64	LYS
1	A	94	ASP
1	A	125	VAL
1	A	144	VAL
1	A	194	ASP
1	A	195	ILE
1	A	199	LEU
1	A	246	GLN
1	A	288	LEU
1	A	299	THR
1	B	38	ILE
1	B	44	LEU
1	B	55	SER
1	B	63	ASP
1	B	84	ASP
1	B	95	SER
1	B	104	SER
1	B	106	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	126	SER
1	B	139	LYS
1	B	153	GLU
1	B	182	GLN
1	B	194	ASP
1	B	199	LEU
1	B	230	MET
1	B	255	SER
1	B	280	LEU
1	B	312	ILE
1	B	323	LYS
1	B	356	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	86	ASN
1	A	100	GLN
1	A	189	HIS
1	A	246	GLN
1	A	286	ASN
1	B	12	ASN
1	B	165	HIS
1	B	166	GLN
1	B	182	GLN
1	B	317	HIS
1	B	362	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	2HQ	B	1375	-	13,13,13	0.70	0	19,19,19	1.05	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	2HQ	B	1375	-	-	-	0/2/2/2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	364/374 (97%)	0.55	14 (3%) 44 46	31, 51, 77, 107	0
1	B	369/374 (98%)	0.87	30 (8%) 18 19	29, 50, 74, 105	0
All	All	733/748 (97%)	0.71	44 (6%) 27 29	29, 50, 76, 107	0

All (44) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	374	LEU	4.2
1	A	374	LEU	3.6
1	A	239	ASP	3.5
1	B	360	HIS	3.5
1	B	341	CYS	3.3
1	B	361	LEU	3.1
1	A	84	ASP	3.0
1	A	361	LEU	2.9
1	A	165	HIS	2.8
1	A	373	THR	2.7
1	B	147	GLN	2.7
1	B	311	ASP	2.7
1	A	16	TYR	2.6
1	A	230	MET	2.6
1	B	354	LEU	2.6
1	A	43	PHE	2.5
1	B	300	ALA	2.5
1	B	146	GLU	2.5
1	B	153	GLU	2.5
1	B	373	THR	2.4
1	A	154	LEU	2.4
1	B	239	ASP	2.4
1	B	230	MET	2.4
1	A	297	SER	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	38	ILE	2.3
1	A	245	TYR	2.3
1	B	353	SER	2.2
1	B	278	ILE	2.2
1	B	43	PHE	2.2
1	B	359	SER	2.2
1	B	304	VAL	2.2
1	B	309	GLY	2.2
1	B	175	THR	2.2
1	A	186	ILE	2.1
1	B	339	LEU	2.1
1	B	314	LYS	2.1
1	B	193	GLU	2.1
1	B	245	TYR	2.1
1	B	296	HIS	2.1
1	B	1	MET	2.1
1	A	38	ILE	2.1
1	B	254	ILE	2.0
1	B	268	ILE	2.0
1	B	284	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	2HQ	B	1375	12/12	0.66	0.18	98,108,116,119	0

6.5 Other polymers [i](#)

There are no such residues in this entry.