



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 14, 2026 – 04:58 PM UTC

PDB ID : 6G4N / pdb_00006g4n
Title : Torpedo californica acetylcholinesterase bound to uncharged hybrid reactivator 2
Authors : Santoni, G.; De la Mora, E.; de Souza, J.; Silman, I.; Sussman, J.; Baati, R.; Weik, M.; Nachon, F.
Deposited on : 2018-03-28
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2481 (2.90-2.90)
Clashscore	190562	2690 (2.90-2.90)
Ramachandran outliers	187476	2623 (2.90-2.90)
Sidechain outliers	187428	2625 (2.90-2.90)
RSRZ outliers	180081	2481 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	537	% 72% 27% ..
1	B	537	% 70% 28% ..

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 8965 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Acetylcholinesterase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	532	Total	C	N	O	S	0	0	0
			4244	2721	720	781	22			
1	B	532	Total	C	N	O	S	0	1	0
			4249	2724	721	782	22			

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is 6-[4-[(7-chloranyl-1,2,3,4-tetrahydroacridin-9-yl)amino]butyl]-2-[(oxidanylamino)methyl]pyridin-3-ol (CCD ID: E2W) (formula: $C_{23}H_{27}ClN_4O_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	Cl	N	O	0	0
			30	23	1	4	2		
3	B	1	Total	C	Cl	N	O	0	0
			30	23	1	4	2		

- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Cl	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	172	Total	O	0	0
			172	172		
5	B	183	Total	O	0	0
			183	183		

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Chain A:

Residue	Category
R517	Green
V518	Green
Q519	Green
M520	Green
G521	Green
V522	Green
K530	Green
T535	Green
ALA	Green
CYS	Green
R429	Red
L430	Green
P433	Green
M436	Green
G437	Green
V438	Green
I439	Green
Y442	Green
E443	Green
I444	Green
E445	Green
F446	Green
P451	Green
L452	Green
V453	Green
L456	Green
A457	Green
Y458	Green
T459	Green
K460	Green
E461	Green
E462	Green
S466	Green
R467	Green
K478	Green
N483	Green
E484	Green
P485	Green
H486	Green
S487	Green
Q488	Red
E489	Green
H492	Green
P493	Green
L494	Green
E499	Green
Q500	Green
K501	Green
F502	Green
I503	Green
D504	Green
L505	Green
N506	Green
M510	Green
H513	Green
Q514	Green
R426	Green
A427	Green
E428	Green
P294	Green
V295	Green
L296	Green
L305	Green
G312	Green
N313	Green
F314	Green
Q318	Green
T319	Green
L320	Green
N324	Green
K325	Green
G328	Green
F339	Green
E344	Green
S345	Green
K346	Green
T347	Green
S348	Green
R349	Green
M353	Green
L358	Green
T372	Green
L373	Green
Q374	Green
V375	Green
T376	Green
D377	Green
K378	Green
N383	Green
N387	Green
R388	Green
V395	Green
G396	Green
D397	Green
K501	Green
F502	Green
I503	Green
D504	Green
L505	Green
N506	Green
M510	Green
H513	Green
Q514	Green
R426	Green
A427	Green
E428	Green
P294	Green
V295	Green
L296	Green
L305	Green
G312	Green
N313	Green
F314	Green
Q318	Green
T319	Green
L320	Green
N324	Green
K325	Green
G328	Green
F339	Green
E344	Green
S345	Green
K346	Green
T347	Green
S348	Green
R349	Green
M353	Green
L358	Green
T372	Green
L373	Green
Q374	Green
V375	Green
T376	Green
D377	Green
K378	Green
N383	Green
N387	Green
R388	Green
V395	Green
G396	Green
D397	Green
K501	Green
F502	Green
I503	Green
D504	Green
L505	Green
N506	Green
M510	Green
H513	Green
Q514	Green
R426	Green
A427	Green
E428	Green
P294	Green
V295	Green
L296	Green
L305	Green
G312	Green
N313	Green
F314	Green
Q318	Green
T319	Green
L320	Green
N324	Green
K325	Green
G328	Green
F339	Green
E344	Green
S345	Green
K346	Green
T347	Green
S348	Green
R349	Green
M353	Green
L358	Green
T372	Green
L373	Green
Q374	Green
V375	Green
T376	Green
D377	Green
K378	Green
N383	Green
N387	Green
R388	Green
V395	Green
G396	Green
D397	Green
K501	Green
F502	Green
I503	Green
D504	Green
L505	Green
N506	Green
M510	Green
H513	Green
Q514	Green
R426	Green
A427	Green
E428	Green
P294	Green
V295	Green
L296	Green
L305	Green
G312	Green
N313	Green
F314	Green
Q318	Green
T319	Green
L320	Green
N324	Green
K325	Green
G328	Green
F339	Green
E344	Green
S345	Green
K346	Green
T347	Green
S348	Green
R349	Green
M35	

Chain B:

Row	Col 1	Col 2	Col 3	Col 4	Col 5	Col 6	Col 7	Col 8	Col 9	Col 10	Col 11	Col 12
1	V395	G396	D397	H398	N399	V400	I401	M405	H406	F407	V408	L420
2	L421	F422	F423	N424	H425	A426	L430	V431	W432	F433	A434	A435
3	A436	G437	V438	I439	H440	E443	I444	E445	V453	K454	E455	Y458
4	E463	S466	R467	R468	I469	M470	K478	M481	E484	P485	H486	S487



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	91.81Å 107.21Å 150.45Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.29 – 2.90 46.29 – 2.90	Depositor EDS
% Data completeness (in resolution range)	99.7 (46.29-2.90) 99.9 (46.29-2.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.25 (at 2.91Å)	Xtriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, R_{free}	0.187 , 0.271 0.193 , 0.277	Depositor DCC
R_{free} test set	1007 reflections (3.00%)	wwPDB-VP
Wilson B-factor (Å ²)	36.4	Xtriage
Anisotropy	1.212	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 54.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8965	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 37.41 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.3091e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, CL, E2W

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.39	1/4367 (0.0%)	0.64	2/5929 (0.0%)
1	B	0.35	0/4375	0.61	0/5940
All	All	0.37	1/8742 (0.0%)	0.62	2/11869 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	38	PRO	C-O	-5.05	1.18	1.24

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	37	GLU	CA-C-N	-5.87	114.33	120.38
1	A	37	GLU	C-N-CA	-5.87	114.33	120.38

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4244	0	4096	114	0
1	B	4249	0	4102	115	0
2	A	28	0	26	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	28	0	26	0	0
3	A	30	0	0	1	0
3	B	30	0	0	1	0
4	B	1	0	0	0	0
5	A	172	0	0	7	0
5	B	183	0	0	13	0
All	All	8965	0	8250	229	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 229 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:459:THR:HG22	1:A:462:GLU:HG3	1.24	1.14
1:B:347:ILE:HG12	1:B:388:ARG:HB2	1.52	0.91
1:A:167:ASN:HD21	1:A:295:VAL:H	1.16	0.90
1:B:375:TYR:CZ	1:B:520:MET:HE2	2.08	0.89
1:B:43:MET:SD	1:B:46:ARG:NH2	2.50	0.84

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	530/537 (99%)	495 (93%)	31 (6%)	4 (1%)	16	44
1	B	531/537 (99%)	494 (93%)	33 (6%)	4 (1%)	16	44
All	All	1061/1074 (99%)	989 (93%)	64 (6%)	8 (1%)	16	44

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	23	LEU
1	A	488	GLN
1	B	380	ASP
1	A	42	ASN
1	B	515	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	465/469 (99%)	456 (98%)	9 (2%)	50	79
1	B	466/469 (99%)	459 (98%)	7 (2%)	57	84
All	All	931/938 (99%)	915 (98%)	16 (2%)	53	82

5 of 16 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	490	SER
1	B	358	LEU
1	A	530	LYS
1	B	356	VAL
1	A	494	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 18 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	398	HIS
1	B	525	ASN
1	B	513	HIS
1	A	514	GLN
1	B	272	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 1 is monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	NAG	A	602	1	14,14,15	0.70	0	17,19,21	0.80	1 (5%)
2	NAG	A	601	1	14,14,15	0.48	0	17,19,21	0.71	1 (5%)
2	NAG	B	602	1	14,14,15	0.83	1 (7%)	17,19,21	0.81	1 (5%)
3	E2W	A	603	-	32,33,33	1.32	3 (9%)	38,45,45	1.83	6 (15%)
3	E2W	B	603	-	32,33,33	1.05	1 (3%)	38,45,45	1.89	6 (15%)
2	NAG	B	601	1	14,14,15	0.50	0	17,19,21	0.77	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	602	1	-	2/6/23/26	0/1/1/1
2	NAG	A	601	1	-	2/6/23/26	0/1/1/1
2	NAG	B	602	1	-	2/6/23/26	0/1/1/1
3	E2W	A	603	-	-	6/10/18/18	0/4/4/4
3	E2W	B	603	-	-	5/10/18/18	0/4/4/4
2	NAG	B	601	1	-	2/6/23/26	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	603	E2W	C9-N10	2.87	1.35	1.32
3	A	603	E2W	C15-N11	2.63	1.51	1.45
3	B	603	E2W	C8-C9	2.57	1.44	1.40
3	A	603	E2W	C8-C9	2.46	1.44	1.40
2	B	602	NAG	C1-C2	2.46	1.55	1.52

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	603	E2W	C23-C10-N25	8.84	135.71	113.05
3	A	603	E2W	C23-C10-N25	7.89	133.27	113.05
3	B	603	E2W	C11-C8-C9	-3.35	117.75	120.90
3	B	603	E2W	C12-C11-C8	3.31	119.56	112.84
3	A	603	E2W	C12-C11-C8	3.22	119.38	112.84

There are no chirality outliers.

5 of 19 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	603	E2W	N25-C10-C23-C22
3	A	603	E2W	C15-C16-C17-C18
2	A	601	NAG	C4-C5-C6-O6
3	B	603	E2W	C15-C16-C17-C18
2	A	602	NAG	O5-C5-C6-O6


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	603	E2W	1	0
3	B	603	E2W	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	532/537 (99%)	0.02	6 (1%) 78 71	17, 33, 58, 125	0
1	B	532/537 (99%)	0.04	5 (0%) 81 75	19, 35, 59, 96	1 (0%)
All	All	1064/1074 (99%)	0.03	11 (1%) 79 73	17, 34, 59, 125	1 (0%)

The worst 5 of 11 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	486	HIS	2.7
1	B	486	HIS	2.7
1	B	260	GLU	2.6
1	B	534	ALA	2.6
1	A	453	VAL	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	NAG	A	601	14/15	0.78	0.13	58,67,76,82	0
2	NAG	B	601	14/15	0.78	0.13	43,63,72,75	0
2	NAG	A	602	14/15	0.80	0.12	31,45,65,65	0
2	NAG	B	602	14/15	0.83	0.12	19,41,63,64	0
3	E2W	A	603	30/30	0.87	0.14	17,52,66,87	0
3	E2W	B	603	30/30	0.89	0.15	36,55,87,88	0
4	CL	B	604	1/1	0.90	0.14	63,63,63,63	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around E2W B 603:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.