



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 01:19 PM UTC

PDB ID : 6G4R / pdb_00006g4r
Title : Corynebacterium glutamicum OxyR C206S mutant, H2O2-bound
Authors : Young, D.R.; Pedre, B.P.; Messens, J.M.
Deposited on : 2018-03-28
Resolution : 2.62 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.62 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4951 (2.64-2.60)
Clashscore	190562	5303 (2.64-2.60)
Ramachandran outliers	187476	5217 (2.64-2.60)
Sidechain outliers	187428	5217 (2.64-2.60)
RSRZ outliers	180081	4950 (2.64-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	327	3% 89% 8% • •
2	A	327	8% 86% 9% • 5%
2	G	327	16% 81% 8% • 10%
3	E	327	3% 88% 7% 5%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 9353 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hydrogen peroxide-inducible genes activator.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	322	Total	C	N	O	S	0	4	0
			2436	1540	408	480	8			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	206	SER	CYS	engineered mutation	UNP A0A2H5I9R9

- Molecule 2 is a protein called Hydrogen peroxide-inducible genes activator.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	310	Total	C	N	O	S	0	6	0
			2308	1467	383	450	8			
2	G	293	Total	C	N	O	S	0	5	0
			2162	1376	358	421	7			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	206	SER	CYS	engineered mutation	UNP A0A2H5I9R9
G	206	SER	CYS	engineered mutation	UNP A0A2H5I9R9


- Molecule 3 is a protein called Hydrogen peroxide-inducible genes activator.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	310	Total	C	N	O	S	0	2	0
			2327	1473	390	456	8			

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
E	206	SER	CYS	engineered mutation	UNP A0A2H5I9R9

- Molecule 4 is HYDROGEN PEROXIDE (CCD ID: PEO) (formula: H_2O_2).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	O	0	0
			2	2		
4	E	1	Total	O	0	0
			2	2		
4	E	1	Total	O	0	0
			2	2		

- Molecule 5 is SULFATE ION (CCD ID: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 6 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			4	2	2		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	B	32	Total	O	0	0
			32	32		
7	A	35	Total	O	0	0
			35	35		
7	E	20	Total	O	0	0
			20	20		
7	G	18	Total	O	0	0
			18	18		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Hydrogen peroxide-inducible genes activator





- Molecule 2: Hydrogen peroxide-inducible genes activator



- Molecule 2: Hydrogen peroxide-inducible genes activator



- Molecule 3: Hydrogen peroxide-inducible genes activator



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	71.22Å 56.99Å 154.18Å 90.00° 98.02° 90.00°	Depositor
Resolution (Å)	76.34 – 2.62 76.34 – 2.62	Depositor EDS
% Data completeness (in resolution range)	99.8 (76.34-2.62) 98.5 (76.34-2.62)	Depositor EDS
R_{merge}	0.25	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.55 (at 2.62Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	(Not available) , (Not available) (Not available) , 0.295	Depositor DCC
R_{free} test set	1919 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å ²)	43.8	Xtriage
Anisotropy	0.902	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 65.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	9353	wwPDB-VP
Average B, all atoms (Å ²)	74.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.95% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CSD, CSO, EDO, OCS, SO4, PEO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	B	0.09	0/2465	0.27	0/3360
2	A	0.10	0/2347	0.33	0/3207
2	G	0.13	0/2199	0.35	0/3005
3	E	0.10	0/2358	0.31	0/3215
All	All	0.10	0/9369	0.32	0/12787

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	G	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	G	200	LEU	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	2436	0	2461	16	0
2	A	2308	0	2306	12	0
2	G	2162	0	2114	10	0
3	E	2327	0	2334	10	0
4	B	2	0	0	0	0
4	E	4	0	0	0	0
5	B	5	0	0	0	0
6	A	4	0	6	0	0
7	A	35	0	0	1	0
7	B	32	0	0	0	0
7	E	20	0	0	0	0
7	G	18	0	0	0	0
All	All	9353	0	9221	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:193:LEU:HD12	2:A:198:LEU:HD11	1.73	0.69
1:B:202:ASP:OD1	1:B:205:HIS:ND1	2.26	0.67
2:A:169:ASP:OD2	7:A:501:HOH:O	2.15	0.64
3:E:157:SER:O	3:E:159:ALA:N	2.33	0.61
2:G:193:LEU:HD12	2:G:198:LEU:HD11	1.82	0.60
2:A:117:LEU:HD21	2:A:131[B]:ILE:HD11	1.84	0.59
2:G:14:ARG:HA	2:G:79:LEU:HD11	1.85	0.57
1:B:86:LEU:HD11	1:B:90:LYS:HE3	1.88	0.56
3:E:150:VAL:HG23	3:E:284:TYR:HB3	1.89	0.54
2:A:74:PHE:HZ	2:G:99:PRO:HG2	1.72	0.54
2:G:88:HIS:CD2	2:G:99:PRO:HD3	2.43	0.54
3:E:209:ASP:HA	3:E:212:VAL:HG22	1.88	0.53
3:E:23:LYS:O	3:E:62:VAL:HG22	2.10	0.51
1:B:56:GLU:HB3	1:B:63:ILE:HG12	1.93	0.51
3:E:53:GLN:HG3	3:E:65:THR:HG21	1.93	0.51
1:B:223:ILE:HD11	2:A:146:GLY:HA3	1.94	0.49
2:A:155:LEU:HD21	2:A:281:GLY:HA3	1.94	0.48
1:B:143:LEU:HD23	1:B:162:MET:HE3	1.95	0.48
2:G:150:VAL:HG23	2:G:284:TYR:HB3	1.96	0.48
1:B:201:LEU:O	1:B:208:HIS:ND1	2.46	0.47
1:B:150:VAL:HG23	1:B:284:TYR:HB3	1.96	0.46
2:G:239:MET:O	2:G:243:VAL:HG23	2.16	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:133:GLU:HB3	3:E:232:ALA:HA	1.98	0.45
3:E:157:SER:C	3:E:159:ALA:H	2.25	0.45
2:A:53:GLN:HG3	2:A:65:THR:HG21	1.99	0.45
2:G:208:HIS:CE1	2:G:213:ASP:HA	2.51	0.45
1:B:112:ILE:O	1:B:116:MET:HB3	2.17	0.45
1:B:90:LYS:HB3	1:B:95:SER:HB2	1.99	0.44
1:B:53[B]:GLN:HG3	1:B:65:THR:HG21	2.00	0.44
1:B:160:PRO:HG3	2:A:202:ASP:OD1	2.19	0.43
3:E:116:MET:HG3	3:E:301:LEU:HD23	1.99	0.42
3:E:165:ILE:HB	3:E:282:LEU:HB3	2.01	0.42
2:G:243:VAL:HG13	2:G:264:GLY:HA3	2.01	0.42
1:B:194:GLU:HG2	1:B:220:ILE:HG23	2.02	0.42
2:A:53:GLN:NE2	2:A:55:ILE:O	2.53	0.42
2:G:296:GLN:HA	2:G:299:LEU:HD22	2.02	0.42
1:B:53[A]:GLN:HG2	1:B:65:THR:HG21	2.02	0.41
3:E:257:PRO:HG3	3:E:312:ALA:HA	2.02	0.41
1:B:219:ASP:N	1:B:219:ASP:OD1	2.54	0.41
1:B:290:ARG:HD2	1:B:293:GLU:OE1	2.21	0.41
2:G:90:LYS:C	2:G:92:ALA:H	2.29	0.41
2:A:172:PHE:CZ	2:A:205:HIS:HB3	2.56	0.41
1:B:165:ILE:HB	1:B:282:LEU:HB3	2.04	0.40
2:A:236:THR:O	2:A:240:GLN:HG2	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	322/327 (98%)	315 (98%)	7 (2%)	0	100	100
2	A	309/327 (94%)	296 (96%)	9 (3%)	4 (1%)	9	19
2	G	289/327 (88%)	276 (96%)	9 (3%)	4 (1%)	9	17

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	E	307/327 (94%)	297 (97%)	8 (3%)	2 (1%)	18	35
All	All	1227/1308 (94%)	1184 (96%)	33 (3%)	10 (1%)	18	31

All (10) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	E	158	GLU
2	A	90	LYS
2	A	131[A]	ILE
2	A	131[B]	ILE
3	E	96	LEU
2	G	91	GLY
2	G	95	SER
2	G	213	ASP
2	A	91	GLY
2	G	211	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	265/269 (98%)	256 (97%)	9 (3%)	32	58
2	A	246/270 (91%)	235 (96%)	11 (4%)	24	48
2	G	222/270 (82%)	211 (95%)	11 (5%)	22	44
3	E	250/270 (93%)	244 (98%)	6 (2%)	43	69
All	All	983/1079 (91%)	946 (96%)	37 (4%)	30	54

All (37) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	36	GLN
1	B	56	GLU
1	B	59	THR
1	B	62	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	63	ILE
1	B	116	MET
1	B	199	LEU
1	B	219	ASP
1	B	223	ILE
2	A	44	VAL
2	A	61	LYS
2	A	62	VAL
2	A	96	LEU
2	A	102	VAL
2	A	134	ASP
2	A	155	LEU
2	A	246	LEU
2	A	296	GLN
2	A	301	LEU
2	A	319	LEU
3	E	96	LEU
3	E	200	LEU
3	E	246	LEU
3	E	283	VAL
3	E	296[A]	GLN
3	E	296[B]	GLN
2	G	18	THR
2	G	44	VAL
2	G	96	LEU
2	G	102	VAL
2	G	117	LEU
2	G	134	ASP
2	G	139	LEU
2	G	205	HIS
2	G	243	VAL
2	G	296	GLN
2	G	299	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
2	G	208	HIS
2	G	269	ASN
2	G	296	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

5 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	CSD	G	215	2	4,7,8	0.95	0	1,8,10	0.88	0
1	OCS	B	22	1	6,8,9	0.86	0	7,11,13	1.34	0
2	CSD	A	215	2	4,7,8	1.04	0	1,8,10	0.16	0
1	CSO	B	215	1	3,6,7	0.63	0	1,6,8	0.51	0
3	CSD	E	22	3	4,7,8	1.04	0	1,8,10	0.25	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CSD	G	215	2	-	2/2/6/8	-
1	OCS	B	22	1	-	0/4/7/9	-
2	CSD	A	215	2	-	2/2/6/8	-
1	CSO	B	215	1	-	1/1/5/7	-
3	CSD	E	22	3	-	1/2/6/8	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	B	215	CSO	N-CA-CB-SG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	215	CSD	CA-CB-SG-OD1
3	E	22	CSD	CA-CB-SG-OD1
2	G	215	CSD	N-CA-CB-SG
2	G	215	CSD	CA-CB-SG-OD1
2	A	215	CSD	N-CA-CB-SG

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	PEO	B	401	-	1,1,1	0.11	0	-		
4	PEO	E	402	-	1,1,1	0.11	0	-		
6	EDO	A	401	-	3,3,3	0.43	0	2,2,2	0.38	0
5	SO4	B	402	-	4,4,4	0.24	0	6,6,6	0.09	0
4	PEO	E	401	-	1,1,1	0.11	0	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	EDO	A	401	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	B	320/327 (97%)	0.48	10 (3%)	51 46	26, 60, 107, 201	4 (1%)
2	A	309/327 (94%)	0.81	25 (8%)	18 14	27, 76, 138, 169	6 (1%)
2	G	292/327 (89%)	1.01	51 (17%)	4 3	23, 78, 180, 283	5 (1%)
3	E	309/327 (94%)	0.49	11 (3%)	46 41	23, 64, 115, 222	2 (0%)
All	All	1230/1308 (94%)	0.69	97 (7%)	18 15	23, 67, 139, 283	17 (1%)

All (97) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	G	68	GLY	6.0
2	G	46	LEU	4.8
2	G	213	ASP	4.4
2	G	204	GLY	4.0
2	A	45	ALA	3.8
2	G	212	VAL	3.8
2	G	205	HIS	3.3
2	G	207	LEU	3.3
2	G	28	ALA	3.3
2	G	20	ALA	3.3
2	G	39	LEU	3.2
2	G	67	ALA	3.2
2	A	223	ILE	3.1
2	G	18	THR	3.1
2	G	96	LEU	3.0
2	G	214	LEU	3.0
3	E	320	LYS	3.0
2	A	260	CYS	2.9
1	B	225	SER	2.9
2	G	206	SER	2.9
2	A	75	ALA	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	G	8	PRO	2.9
2	G	210	GLN	2.9
2	A	220	ILE	2.9
2	G	203	ASP	2.9
2	G	36	GLN	2.8
1	B	228	ALA	2.8
2	G	211	ILE	2.8
2	G	198	LEU	2.8
2	G	217	ARG	2.8
2	A	81	ALA	2.7
2	G	7	ARG	2.7
2	A	82	ALA	2.7
2	G	42	ALA	2.7
2	A	20	ALA	2.7
2	G	16	PHE	2.7
2	A	211	ILE	2.7
2	G	19	ILE	2.7
2	G	71	LEU	2.6
2	A	212	VAL	2.6
2	G	75	ALA	2.6
2	G	72	LEU	2.6
3	E	221	ASN	2.5
2	A	89	ALA	2.5
1	B	323	VAL	2.5
2	G	25	PHE	2.5
2	G	63	ILE	2.5
3	E	26	GLY	2.5
2	G	182	PHE	2.5
3	E	29	ALA	2.5
2	G	55	ILE	2.4
1	B	26	GLY	2.4
2	A	262	ARG	2.4
2	G	10	LEU	2.4
2	G	201	LEU	2.4
2	A	296	GLN	2.4
1	B	96	LEU	2.4
1	B	201	LEU	2.4
2	A	13	LEU	2.4
3	E	96	LEU	2.4
3	E	220	ILE	2.4
2	A	159	ALA	2.4
2	G	82	ALA	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	A	6	TYR	2.3
2	G	118	SER	2.3
1	B	226	THR	2.3
2	A	25	PHE	2.3
2	G	69	GLU	2.2
2	A	122	GLU	2.2
2	G	123	GLU	2.2
3	E	228	ALA	2.2
1	B	227	THR	2.2
2	A	79	LEU	2.2
2	A	233	SER	2.2
2	G	43	LEU	2.2
2	G	200	LEU	2.2
1	B	325	VAL	2.2
2	G	74	PHE	2.2
2	G	6	TYR	2.2
3	E	199	LEU	2.2
2	A	219	ASP	2.2
2	A	63	ILE	2.2
2	G	70	LYS	2.1
2	A	195	ASP	2.1
2	G	24	HIS	2.1
2	A	60	ARG	2.1
2	G	160	PRO	2.1
1	B	25	PHE	2.1
2	A	62	VAL	2.1
2	G	22	CYS	2.1
2	G	208	HIS	2.1
3	E	146	GLY	2.1
2	G	195	ASP	2.0
3	E	60	ARG	2.0
2	G	121	ASP	2.0
2	G	157	SER	2.0
3	E	200	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	CSD	G	215	8/9	0.46	0.13	151,154,155,158	0
2	CSD	A	215	8/9	0.57	0.11	124,137,148,155	0
1	CSO	B	215	7/8	0.85	0.12	57,64,68,73	0
3	CSD	E	22	8/9	0.93	0.09	67,76,78,83	0
1	OCS	B	22	9/10	0.94	0.08	64,76,80,83	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	PEO	E	401	2/2	0.77	0.13	50,50,50,52	0
4	PEO	E	402	2/2	0.81	0.19	86,86,86,86	0
4	PEO	B	401	2/2	0.87	0.09	31,31,31,33	0
6	EDO	A	401	4/4	0.87	0.11	56,57,60,62	0
5	SO4	B	402	5/5	0.97	0.08	41,45,51,64	0

6.5 Other polymers [i](#)

There are no such residues in this entry.