



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 09:21 AM UTC

PDB ID : 6G4Z / pdb_00006g4z
Title : Crystal structure of murine NF-kappaB inducing kinase (NIK) in complex with compound 2f
Authors : Leonardo-Silvestre, H.; McEwan, P.A.; Hymowitz, S.G.
Deposited on : 2018-03-28
Resolution : 2.84 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.84 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1520 (2.86-2.82)
Clashscore	190562	1559 (2.86-2.82)
Ramachandran outliers	187476	1517 (2.86-2.82)
Sidechain outliers	187428	1518 (2.86-2.82)
RSRZ outliers	180081	1521 (2.86-2.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	349	
1	B	349	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 5103 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Mitogen-activated protein kinase kinase kinase 14.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	322	Total	C	N	O	S	0	1	0
			2514	1592	446	455	21			
1	B	325	Total	C	N	O	S	0	0	0
			2513	1588	445	459	21			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	327	GLY	-	expression tag	UNP Q9WUL6
A	328	SER	-	expression tag	UNP Q9WUL6
B	327	GLY	-	expression tag	UNP Q9WUL6
B	328	SER	-	expression tag	UNP Q9WUL6

- Molecule 2 is 5-fluoranyl-1-[4-[2-[(3 {R})-1-methyl-3-oxidanyl-2-oxidanylidene-pyrrol-3-yl]ethynyl]pyridin-2-yl]indazole-3-carboxamide (CCD ID: ELW) (formula: C₂₀H₁₄FN₅O₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	F	N	O	0	0
			29	20	1	5	3		
2	B	1	Total	C	F	N	O	0	0
			29	20	1	5	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	9	Total	O	0	0
			9	9		
3	B	9	Total	O	0	0
			9	9		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Mitogen-activated protein kinase kinase kinase 14



- Molecule 1: Mitogen-activated protein kinase kinase kinase 14



4 Data and refinement statistics

Property	Value	Source
Space group	P 43	Depositor
Cell constants a, b, c, α , β , γ	144.83Å 144.83Å 46.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.21 – 2.84 36.21 – 2.84	Depositor EDS
% Data completeness (in resolution range)	98.0 (36.21-2.84) 98.0 (36.21-2.84)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.82 (at 2.85Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
R, R_{free}	0.234 , 0.283 0.236 , 0.281	Depositor DCC
R_{free} test set	1171 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å ²)	74.1	Xtriage
Anisotropy	0.032	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 39.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.036 for h,-k,-l	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	5103	wwPDB-VP
Average B, all atoms (Å ²)	77.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.80% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ELW

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.48	0/2576	0.82	3/3482 (0.1%)
1	B	0.46	0/2570	0.73	0/3474
All	All	0.47	0/5146	0.78	3/6956 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	604	GLY	CA-C-N	11.46	134.16	119.84
1	A	604	GLY	C-N-CA	11.46	134.16	119.84
1	A	408	VAL	N-CA-C	6.65	123.17	109.34

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2514	0	2516	17	0
1	B	2513	0	2519	32	0
2	A	29	0	0	1	0
2	B	29	0	0	0	0
3	A	9	0	0	0	0
3	B	9	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	5103	0	5035	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:405:GLN:HB2	1:B:406:PRO:CD	1.59	1.24
1:B:405:GLN:CD	1:B:406:PRO:HD2	1.77	1.09
1:B:405:GLN:CB	1:B:406:PRO:HD3	1.81	1.07
1:B:405:GLN:CB	1:B:406:PRO:CD	2.34	0.93
1:B:405:GLN:CD	1:B:406:PRO:CD	2.45	0.88
1:B:405:GLN:HB2	1:B:406:PRO:HD3	0.88	0.86
1:A:603:ARG:HH11	1:A:603:ARG:HG3	1.42	0.84
1:A:398:GLU:OE2	1:A:403:THR:HG22	1.81	0.79
1:A:606:LEU:HD12	1:A:606:LEU:N	2.02	0.75
1:B:405:GLN:OE1	1:B:406:PRO:HD2	1.89	0.72
1:A:603:ARG:HG3	1:A:603:ARG:NH1	1.99	0.70
1:A:606:LEU:HD12	1:A:606:LEU:H	1.59	0.67
1:A:603:ARG:HH11	1:A:603:ARG:CG	2.13	0.59
1:A:355:LEU:HD11	1:A:460:ALA:HB3	1.86	0.57
1:B:405:GLN:NE2	1:B:406:PRO:CD	2.69	0.56
1:A:403:THR:HG23	1:A:403:THR:O	2.06	0.55
1:B:405:GLN:CG	1:B:406:PRO:HD2	2.37	0.55
1:B:405:GLN:CG	1:B:406:PRO:CD	2.85	0.55
1:B:493:ARG:HE	1:B:663:LEU:HD11	1.73	0.54
1:B:571:MET:HE3	1:B:611:ALA:HB1	1.89	0.54
1:B:408:VAL:HG21	1:B:418:ARG:HH21	1.73	0.53
1:B:544:GLN:O	1:B:546:ASP:N	2.41	0.53
1:B:408:VAL:HG12	3:B:805:HOH:O	2.09	0.52
1:A:514:LEU:HD11	1:A:579:VAL:CG2	2.42	0.49
1:B:405:GLN:CB	1:B:406:PRO:HD2	2.38	0.49
1:A:639:GLU:HG3	1:A:641:VAL:HG22	1.95	0.49
1:B:552:LEU:O	1:B:553:LEU:HB2	2.13	0.48
1:A:443:LEU:O	1:A:447:ALA:N	2.45	0.48
1:B:625:ALA:HB1	1:B:626:PRO:HD2	1.95	0.48
1:B:527:SER:O	1:B:667:TRP:CH2	2.68	0.46
1:B:355:LEU:HD11	1:B:460:ALA:HB3	1.97	0.46
1:A:606:LEU:N	1:A:606:LEU:CD1	2.73	0.46
1:B:405:GLN:NE2	1:B:406:PRO:HD3	2.30	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:552:LEU:O	1:B:553:LEU:CB	2.64	0.46
1:A:362:TRP:O	1:A:363:SER:C	2.60	0.44
1:B:406:PRO:O	1:B:406:PRO:HG2	2.18	0.44
1:A:606:LEU:H	1:A:606:LEU:CD1	2.28	0.43
1:B:404:HIS:HB2	1:B:418:ARG:HB3	1.99	0.43
1:B:573:LYS:HB3	1:B:574:PRO:HD2	2.01	0.43
1:B:474:LEU:HD22	1:B:524:LEU:HB2	2.00	0.43
1:B:637:ARG:HG2	1:B:642:HIS:HB2	2.00	0.42
1:A:416:VAL:HG21	2:A:701:ELW:C13	2.50	0.42
1:B:391:VAL:O	1:B:392:ASP:C	2.63	0.42
1:B:387:LYS:HB3	1:B:426:PHE:CD2	2.53	0.42
1:B:431:LYS:HE2	1:B:433:VAL:HG22	2.01	0.42
1:B:493:ARG:HD2	1:B:497:TYR:CE2	2.55	0.42
1:A:439:ARG:NH2	1:A:541:LEU:HD23	2.35	0.41
1:A:347:VAL:HA	1:A:381:GLY:O	2.20	0.41
1:B:621:PRO:HA	1:B:622:PRO:HD3	1.99	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	317/349 (91%)	296 (93%)	19 (6%)	2 (1%)	21	39
1	B	319/349 (91%)	296 (93%)	18 (6%)	5 (2%)	7	16
All	All	636/698 (91%)	592 (93%)	37 (6%)	7 (1%)	11	23

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	605	PRO
1	B	405	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	553	LEU
1	B	546	ASP
1	A	407	ARG
1	B	545	PRO
1	B	406	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	271/290 (93%)	261 (96%)	10 (4%)	30	55
1	B	271/290 (93%)	263 (97%)	8 (3%)	36	60
All	All	542/580 (93%)	524 (97%)	18 (3%)	33	58

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	404	HIS
1	A	407	ARG
1	A	454	ILE
1	A	519	LYS
1	A	527	SER
1	A	552	LEU
1	A	603	ARG
1	A	634	GLU
1	A	664	LYS
1	A	667	TRP
1	B	389	LYS
1	B	402	MET
1	B	404	HIS
1	B	405	GLN
1	B	408	VAL
1	B	419	MET
1	B	433	VAL
1	B	542	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	404	HIS
1	A	468	ASN
1	A	593	ASN
1	B	405	GLN
1	B	509	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ELW	B	701	-	29,32,32	1.57	5 (17%)	37,48,48	2.04	9 (24%)
2	ELW	A	701	-	29,32,32	1.59	5 (17%)	37,48,48	1.99	9 (24%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ELW	B	701	-	-	0/10/30/30	0/4/4/4
2	ELW	A	701	-	-	1/10/30/30	0/4/4/4

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	701	ELW	C3-C4	4.29	1.51	1.34
2	A	701	ELW	C3-C4	4.28	1.51	1.34
2	A	701	ELW	C17-C18	-3.54	1.44	1.50
2	A	701	ELW	N15-N16	-3.45	1.34	1.38
2	B	701	ELW	C17-C18	-3.38	1.44	1.50
2	B	701	ELW	N15-N16	-3.10	1.35	1.38
2	B	701	ELW	C3-N2	2.87	1.43	1.37
2	A	701	ELW	C3-N2	2.79	1.43	1.37
2	B	701	ELW	C21-C22	-2.08	1.38	1.41
2	A	701	ELW	C21-C17	-2.05	1.41	1.45

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	701	ELW	C3-N2-C28	6.14	114.47	107.34
2	A	701	ELW	C3-N2-C28	6.09	114.41	107.34
2	A	701	ELW	C14-C13-N15	-5.49	116.04	119.62
2	A	701	ELW	C22-N15-N16	5.04	113.21	111.02
2	B	701	ELW	C22-N15-N16	4.93	113.17	111.02
2	B	701	ELW	C22-N15-C13	-4.34	126.49	130.21
2	B	701	ELW	C14-C13-N15	-4.28	116.83	119.62
2	A	701	ELW	C23-C24-C25	2.77	121.23	118.38
2	B	701	ELW	C23-C24-C25	2.63	121.08	118.38
2	A	701	ELW	C21-C22-N15	-2.43	105.32	106.42
2	A	701	ELW	C22-N15-C13	-2.20	128.32	130.21
2	B	701	ELW	C21-C22-N15	-2.18	105.43	106.42
2	B	701	ELW	C17-C18-N19	-2.18	112.90	116.73
2	A	701	ELW	C1-N2-C3	-2.15	121.53	125.11
2	A	701	ELW	O29-C28-N2	2.11	127.89	125.96
2	B	701	ELW	N12-C13-N15	2.06	119.70	115.91
2	A	701	ELW	N12-C13-N15	2.04	119.67	115.91
2	B	701	ELW	C1-N2-C3	-2.01	121.76	125.11

There are no chirality outliers.

All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	701	ELW	C5-C7-C8-C9


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	701	ELW	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	322/349 (92%)	0.71	25 (7%)	19 14	37, 70, 110, 139	2 (0%)
1	B	325/349 (93%)	0.74	24 (7%)	20 15	41, 74, 116, 141	1 (0%)
All	All	647/698 (92%)	0.73	49 (7%)	20 15	37, 72, 113, 141	3 (0%)

All (49) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	410	ARG	4.9
1	A	551	SER	4.5
1	A	408	VAL	4.3
1	B	604	GLY	4.2
1	A	410	ARG	4.1
1	A	334	PRO	3.8
1	A	604	GLY	3.4
1	B	407	ARG	3.3
1	A	606	LEU	3.2
1	B	548	LEU	3.1
1	B	334	PRO	2.9
1	B	552	LEU	2.8
1	A	607	CYS	2.8
1	B	607	CYS	2.8
1	B	606	LEU	2.8
1	B	608	LEU	2.8
1	B	409	GLY	2.8
1	B	408	VAL	2.7
1	B	667	TRP	2.7
1	B	562	GLU	2.7
1	B	404	HIS	2.7
1	A	571	MET	2.6
1	A	667	TRP	2.5
1	A	554	THR	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	542	CYS	2.5
1	A	402	MET	2.4
1	B	405	GLN	2.4
1	A	608	LEU	2.4
1	B	598	TRP	2.4
1	A	558	ILE	2.4
1	A	407	ARG	2.4
1	B	445	ALA	2.3
1	A	446	CYS	2.3
1	A	544	GLN	2.3
1	A	603	ARG	2.3
1	B	617	ILE	2.3
1	A	600	GLN	2.2
1	A	671	TYR	2.2
1	A	487	GLY	2.2
1	A	552	LEU	2.2
1	A	484	LYS	2.2
1	A	605	PRO	2.1
1	B	600	GLN	2.1
1	B	558	ILE	2.1
1	B	626	PRO	2.0
1	B	553	LEU	2.0
1	B	398	GLU	2.0
1	B	547	GLY	2.0
1	A	445	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ


There are no oligosaccharides in this entry.


6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	ELW	A	701	29/29	0.91	0.12	66,70,72,76	0
2	ELW	B	701	29/29	0.91	0.11	55,61,65,68	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.