



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 4, 2026 – 09:28 PM UTC

PDB ID : 7G48 / pdb_00007g48

Title : Crystal Structure of rat Autotaxin in complex with [(3aS,6aS)-5-(4-methoxyphenyl)-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-(1H-benzotriazol-5-yl)methanone, i.e. SMILES N1(C[C@@H]2[C@@H](C1)CN(C2)C(=O)c1cc(c2c(c1)cccc2)OC)C(=O)c1ccc2c(c1)N=NN2 with IC₅₀=0.11329 microM

Authors : Stihle, M.; Benz, J.; Hunziker, D.; Mattei, P.; Rudolph, M.G.

Deposited on : 2023-06-05

Resolution : 1.77 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.77 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1365 (1.78-1.78)
Clashscore	190562	1395 (1.78-1.78)
Ramachandran outliers	187476	1382 (1.78-1.78)
Sidechain outliers	187428	1382 (1.78-1.78)
RSRZ outliers	180081	1365 (1.78-1.78)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	846	6% 90% 5% 5%
2	B	8	100%

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 7237 atoms, of which 22 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Isoform 2 of Ectonucleotide pyrophosphatase/phosphodiesterase family member 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	806	Total	C	N	O	S	0	5	0
			6529	4142	1122	1213	52			

There are 14 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	53	ALA	ASN	engineered mutation	UNP Q64610
A	410	ALA	ASN	engineered mutation	UNP Q64610
A	591	THR	ARG	engineered mutation	UNP Q64610
A	863	GLY	-	expression tag	UNP Q64610
A	864	GLY	-	expression tag	UNP Q64610
A	865	ARG	-	expression tag	UNP Q64610
A	866	HIS	-	expression tag	UNP Q64610
A	867	HIS	-	expression tag	UNP Q64610
A	868	HIS	-	expression tag	UNP Q64610
A	869	HIS	-	expression tag	UNP Q64610
A	870	HIS	-	expression tag	UNP Q64610
A	871	HIS	-	expression tag	UNP Q64610
A	872	HIS	-	expression tag	UNP Q64610
A	873	HIS	-	expression tag	UNP Q64610

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	8	Total	C	N	O	0	0	0
			94	52	2	40			

- Molecule 3 is [(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl](4-methoxynaphthalen-2-yl)methanone (CCD ID: YP9) (formula: C₂₅H₂₃N₅O₃) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	H	N	O	0	0
			55	25	22	5	3		

- Molecule 4 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula: C₆H₁₃NO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 5 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	Ca	0	0
			2	2		

- Molecule 6 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Zn	0	0
			1	1		

- Molecule 7 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	2	Total	Na	0	0
			2	2		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	542	Total	O	0	0
			542	542		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Isoform 2 of Ectonucleotide pyrophosphatase/phosphodiesterase family member 2



- Molecule 2: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	84.31Å 92.20Å 119.90Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.10 – 1.77 46.10 – 1.77	Depositor EDS
% Data completeness (in resolution range)	96.8 (46.10-1.77) 96.8 (46.10-1.77)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.47 (at 1.77Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.186 , 0.221 0.185 , 0.219	Depositor DCC
R_{free} test set	4655 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	23.9	Xtriage
Anisotropy	0.020	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 26.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7237	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MES, CA, NA, YP9, ZN, MAN, BMA, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.77	1/6726 (0.0%)	0.88	2/9121 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	163	PRO	CA-C	5.67	1.57	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	424	MET	N-CA-C	-5.48	102.17	110.28
1	A	209	THR	N-CA-C	5.01	116.43	110.97

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6529	0	6285	24	0
2	B	94	0	79	0	0
3	A	33	22	0	0	0
4	A	12	0	13	1	0
5	A	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	A	1	0	0	0	0
7	A	2	0	0	0	0
8	A	542	0	0	5	0
All	All	7215	22	6377	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:373:SER:HA	1:A:379:VAL:HG21	1.63	0.81
1:A:269:ARG:HB3	1:A:300:PRO:HA	1.75	0.68
1:A:539:PHE:O	1:A:541:PRO:HD3	1.98	0.64
4:A:902:MES:H52	8:A:1464:HOH:O	2.01	0.60
1:A:163:PRO:HB3	1:A:350:CYS:O	2.03	0.58
1:A:594:ARG:NH2	8:A:1003:HOH:O	2.36	0.58
1:A:546:GLU:CD	1:A:600:ARG:HH22	2.19	0.51
1:A:449:ARG:O	1:A:450:ARG:HB2	2.11	0.50
1:A:210:PHE:HE2	1:A:243:LEU:HD23	1.78	0.48
1:A:371:PHE:CE1	1:A:457:PRO:HA	2.49	0.47
1:A:550:PRO:HB2	1:A:611:ILE:HG12	1.96	0.47
1:A:179:LYS:HE3	1:A:179:LYS:HB2	1.71	0.47
1:A:222:PRO:HA	1:A:225:HIS:CE1	2.49	0.47
1:A:269:ARG:CZ	1:A:269:ARG:HB2	2.44	0.47
1:A:106:ARG:O	1:A:109:GLU:HB3	2.14	0.47
1:A:856:HIS:HD2	1:A:858:TYR:CZ	2.33	0.46
1:A:821:MET:HA	1:A:821:MET:HE2	1.98	0.46
1:A:162:ARG:HB2	8:A:1154:HOH:O	2.18	0.43
1:A:252:ARG:NH1	8:A:1008:HOH:O	2.45	0.43
1:A:450:ARG:NH1	8:A:1001:HOH:O	2.24	0.43
1:A:98:ARG:HG3	1:A:115:ASN:OD1	2.19	0.41
1:A:148[A]:CYS:SG	1:A:496:TYR:HE1	2.44	0.41
1:A:260:TRP:CZ2	1:A:274:PHE:HZ	2.38	0.41
1:A:508:LEU:HD23	1:A:508:LEU:HA	1.91	0.41
1:A:737:ILE:HB	1:A:766:HIS:HB2	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	805/846 (95%)	785 (98%)	20 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	734/765 (96%)	729 (99%)	5 (1%)	76	66

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	389	LEU
1	A	397	ILE
1	A	571	LYS
1	A	572	VAL
1	A	689	PRO

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	436	ASN
1	A	471	GLN
1	A	484	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	582	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

8 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	B	1	1,2	14,14,15	0.70	0	17,19,21	1.44	2 (11%)
2	NAG	B	2	2	14,14,15	0.66	0	17,19,21	1.75	3 (17%)
2	BMA	B	3	2	11,11,12	0.78	0	15,15,17	1.18	3 (20%)
2	MAN	B	4	2	11,11,12	0.77	0	15,15,17	1.47	2 (13%)
2	MAN	B	5	2	11,11,12	0.64	0	15,15,17	2.01	4 (26%)
2	MAN	B	6	2	11,11,12	0.73	0	15,15,17	0.92	1 (6%)
2	MAN	B	7	2	11,11,12	0.59	0	15,15,17	1.24	1 (6%)
2	MAN	B	8	2	11,11,12	0.77	0	15,15,17	0.85	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	1/6/23/26	0/1/1/1
2	BMA	B	3	2	-	0/2/19/22	0/1/1/1
2	MAN	B	4	2	-	0/2/19/22	0/1/1/1
2	MAN	B	5	2	-	0/2/19/22	0/1/1/1
2	MAN	B	6	2	-	0/2/19/22	0/1/1/1
2	MAN	B	7	2	-	1/2/19/22	0/1/1/1
2	MAN	B	8	2	-	2/2/19/22	0/1/1/1

There are no bond length outliers.

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2	NAG	C1-O5-C5	5.33	119.32	112.19
2	B	5	MAN	C1-O5-C5	5.19	119.14	112.19
2	B	4	MAN	C1-O5-C5	3.69	117.13	112.19
2	B	5	MAN	C3-C4-C5	3.38	116.36	110.23
2	B	1	NAG	O5-C5-C6	2.99	113.49	107.66
2	B	4	MAN	O3-C3-C2	2.91	115.99	110.05
2	B	1	NAG	O5-C1-C2	-2.80	106.96	111.29
2	B	5	MAN	C6-C5-C4	-2.76	106.25	113.02
2	B	3	BMA	C1-O5-C5	2.70	115.81	112.19
2	B	2	NAG	C2-N2-C7	2.47	126.21	122.90
2	B	5	MAN	O5-C5-C4	2.40	116.66	110.83
2	B	3	BMA	C1-C2-C3	2.36	113.08	109.64
2	B	6	MAN	C1-O5-C5	2.31	115.28	112.19
2	B	2	NAG	C3-C4-C5	2.26	114.33	110.23
2	B	7	MAN	C3-C4-C5	2.26	114.33	110.23
2	B	8	MAN	O3-C3-C2	2.10	114.35	110.05
2	B	3	BMA	O6-C6-C5	-2.10	104.20	111.33

There are no chirality outliers.


All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	7	MAN	O5-C5-C6-O6
2	B	8	MAN	C4-C5-C6-O6
2	B	2	NAG	C4-C5-C6-O6
2	B	8	MAN	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 5 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	YP9	A	901	6	38,38,38	2.68	19 (50%)	53,56,56	1.93	12 (22%)
4	MES	A	902	-	12,12,12	1.96	1 (8%)	15,16,16	1.12	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	YP9	A	901	6	-	5/18/38/38	0/6/6/6
4	MES	A	902	-	-	2/6/14/14	0/1/1/1

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	902	MES	C8-S	-5.91	1.69	1.77
3	A	901	YP9	C16-C17	5.35	1.48	1.40
3	A	901	YP9	C14-C15	4.84	1.46	1.38
3	A	901	YP9	C16-C12	4.61	1.46	1.39
3	A	901	YP9	C11-N01	4.56	1.44	1.34
3	A	901	YP9	C02-C05	-4.42	1.46	1.53
3	A	901	YP9	C22-N07	4.27	1.43	1.34
3	A	901	YP9	C03-C04	-4.01	1.47	1.53
3	A	901	YP9	O13-C11	3.88	1.30	1.22
3	A	901	YP9	C27-C23	3.79	1.44	1.37
3	A	901	YP9	C15-C18	3.35	1.45	1.39
3	A	901	YP9	O34-C26	3.14	1.42	1.36
3	A	901	YP9	C25-C23	3.13	1.44	1.39
3	A	901	YP9	C26-C29	3.00	1.50	1.42
3	A	901	YP9	C08-C04	-2.60	1.49	1.53
3	A	901	YP9	C29-C28	-2.51	1.38	1.43
3	A	901	YP9	N21-N20	2.51	1.37	1.31
3	A	901	YP9	C32-C31	2.36	1.43	1.38
3	A	901	YP9	C27-C28	2.31	1.47	1.42
3	A	901	YP9	C05-C04	-2.08	1.48	1.54

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	901	YP9	O34-C26-C29	7.45	123.35	114.56
3	A	901	YP9	C02-C05-C04	-4.45	97.19	103.56
3	A	901	YP9	C35-O34-C26	3.92	124.19	117.67
3	A	901	YP9	O13-C11-N01	-3.48	116.87	122.35
3	A	901	YP9	C12-C11-N01	3.30	122.78	118.66
3	A	901	YP9	C03-C04-C05	-2.99	99.28	103.56
3	A	901	YP9	O34-C26-C25	-2.81	118.27	123.27
3	A	901	YP9	C15-C18-C17	-2.62	119.22	122.10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	902	MES	C5-N4-C3	2.56	114.36	108.84
3	A	901	YP9	C33-C28-C27	-2.52	117.66	122.00
3	A	901	YP9	C06-C05-C04	-2.48	100.01	103.56
3	A	901	YP9	C04-C03-N01	-2.12	99.82	103.02
3	A	901	YP9	C05-C02-N01	-2.06	99.92	103.02

There are no chirality outliers.

All (7) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	901	YP9	C23-C22-N07-C06
3	A	901	YP9	O24-C22-N07-C06
3	A	901	YP9	C25-C26-O34-C35
3	A	901	YP9	C29-C26-O34-C35
4	A	902	MES	C8-C7-N4-C3
4	A	902	MES	C8-C7-N4-C5
3	A	901	YP9	O24-C22-N07-C08

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	902	MES	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	806/846 (95%)	0.12	51 (6%) 26 31	11, 24, 54, 80	5 (0%)

All (51) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	468	CYS	7.8
1	A	469	PHE	6.7
1	A	578	LEU	6.5
1	A	572	VAL	5.9
1	A	862	ILE	5.9
1	A	68	VAL	5.7
1	A	51	TRP	5.1
1	A	581	LEU	4.3
1	A	70	PRO	4.2
1	A	71	PRO	4.1
1	A	53	ALA	3.7
1	A	274	PHE	3.3
1	A	379	VAL	3.3
1	A	243	LEU	3.2
1	A	573	GLU	3.1
1	A	54	THR	3.0
1	A	582	ASN	3.0
1	A	55	SER	3.0
1	A	462	LYS	2.9
1	A	461	TYR	2.9
1	A	69	GLY	2.9
1	A	463	LYS	2.8
1	A	161	VAL	2.8
1	A	585	LEU	2.8
1	A	579	GLU	2.7
1	A	237	PHE	2.6
1	A	366	CYS	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	249	PHE	2.6
1	A	82	TYR	2.6
1	A	56	GLY	2.5
1	A	577	LYS	2.5
1	A	580	GLU	2.4
1	A	246	ARG	2.4
1	A	536	THR	2.4
1	A	124	LEU	2.4
1	A	576	ASN	2.3
1	A	397	ILE	2.3
1	A	378	ASN	2.3
1	A	66	GLN	2.3
1	A	584	ARG	2.3
1	A	242	HIS	2.2
1	A	244	ARG	2.2
1	A	566	CYS	2.2
1	A	72	ASP	2.2
1	A	137	CYS	2.2
1	A	470	PHE	2.2
1	A	251	HIS	2.1
1	A	107	CYS	2.1
1	A	245	GLY	2.0
1	A	861	GLU	2.0
1	A	538	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MAN	B	7	11/12	0.37	0.18	78,80,80,80	0
2	BMA	B	3	11/12	0.61	0.15	56,61,67,74	0
2	MAN	B	8	11/12	0.63	0.18	57,73,77,79	0
2	MAN	B	4	11/12	0.76	0.13	47,50,53,53	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	MAN	B	5	11/12	0.78	0.15	44,52,60,69	0
2	MAN	B	6	11/12	0.85	0.15	35,46,50,50	0
2	NAG	B	2	14/15	0.89	0.10	27,32,41,52	0
2	NAG	B	1	14/15	0.97	0.07	17,19,21,24	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	MES	A	902	12/12	0.88	0.16	31,43,50,50	0
3	YP9	A	901	33/33	0.90	0.11	26,37,44,44	0
5	CA	A	903	1/1	0.94	0.21	45,45,45,45	0
7	NA	A	907	1/1	0.94	0.07	30,30,30,30	0
7	NA	A	906	1/1	0.95	0.07	30,30,30,30	0
5	CA	A	905	1/1	1.00	0.01	18,18,18,18	0
6	ZN	A	904	1/1	1.00	0.01	20,20,20,20	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.