



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 05:40 PM UTC

PDB ID : 3G5I / pdb_00003g5i
Title : Crystal Structure of the E.coli RihA pyrimidine nucleosidase bound to a iminoribitol-based inhibitor
Authors : Garau, G.; Muzzolini, L.; Tornaghi, P.; Degano, M.
Deposited on : 2009-02-05
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	312	
1	B	312	
1	C	312	
1	D	312	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	BME	A	1102	-	X	X	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 9716 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Pyrimidine-specific ribonucleoside hydrolase rihA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	305	Total	C	N	O	S	0	1	0
			2325	1488	387	439	11			
1	B	295	Total	C	N	O	S	0	0	0
			2246	1440	374	421	11			
1	C	293	Total	C	N	O	S	0	0	0
			2230	1427	372	420	11			
1	D	304	Total	C	N	O	S	0	2	0
			2334	1495	388	440	11			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	expression tag	UNP P41409
A	1	SER	-	expression tag	UNP P41409
B	0	GLY	-	expression tag	UNP P41409
B	1	SER	-	expression tag	UNP P41409
C	0	GLY	-	expression tag	UNP P41409
C	1	SER	-	expression tag	UNP P41409
D	0	GLY	-	expression tag	UNP P41409
D	1	SER	-	expression tag	UNP P41409

- Molecule 2 is BETA-MERCAPTOETHANOL (CCD ID: BME) (formula: C₂H₆OS).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	O	S	0	0
			4	2	1	1		

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Ca	0	0
			1	1		
3	B	1	Total	Ca	0	0
			1	1		
3	C	1	Total	Ca	0	0
			1	1		
3	D	1	Total	Ca	0	0
			1	1		

- Molecule 4 is (2S,3S,4R,5R)-2-(3,4-diaminophenyl)-5-(hydroxymethyl)pyrrolidine-3,4-diol (CCD ID: DNB) (formula: C₁₁H₁₇N₃O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			16	11	2	3		
4	B	1	Total	C	N	O	0	0
			16	11	2	3		
4	C	1	Total	C	N	O	0	0
			15	11	1	3		
4	D	1	Total	C	N	O	0	0
			16	11	2	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	141	Total	O	0	0
			141	141		
5	B	115	Total	O	0	0
			115	115		
5	C	132	Total	O	0	0
			132	132		
5	D	122	Total	O	0	0
			122	122		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Pyrimidine-specific ribonucleoside hydrolase rihA



- Molecule 1: Pyrimidine-specific ribonucleoside hydrolase rihA




- Molecule 1: Pyrimidine-specific ribonucleoside hydrolase rihA



- Molecule 1: Pyrimidine-specific ribonucleoside hydrolase rihA





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	84.35Å 82.91Å 93.87Å 90.00° 112.15° 90.00°	Depositor
Resolution (Å)	37.50 – 2.10 37.50 – 2.10	Depositor EDS
% Data completeness (in resolution range)	98.0 (37.50-2.10) 98.0 (37.50-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.24 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.203 , 0.241 (Not available) , 0.278	Depositor DCC
R_{free} test set	3453 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	18.7	Xtriage
Anisotropy	1.052	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 47.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	9716	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 41.70 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.2714e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DNB, BME, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.12	0/2378	1.05	0/3249
1	B	1.03	2/2294 (0.1%)	1.03	2/3133 (0.1%)
1	C	1.06	1/2277 (0.0%)	1.01	1/3110 (0.0%)
1	D	1.01	1/2392 (0.0%)	0.99	2/3266 (0.1%)
All	All	1.05	4/9341 (0.0%)	1.02	5/12758 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	135	ASN	CG-ND2	-7.22	1.18	1.33
1	B	135	ASN	CG-OD1	-7.02	1.10	1.23
1	C	215	ILE	CA-CB	5.36	1.60	1.54
1	D	215	ILE	CA-CB	5.01	1.60	1.54

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	165	PHE	N-CA-C	6.01	118.35	111.02
1	B	181	GLY	N-CA-C	-5.76	107.14	115.27
1	B	177	VAL	N-CA-C	5.39	116.02	110.36
1	D	165	PHE	N-CA-C	5.12	117.27	111.02
1	D	303	LEU	CA-CB-CG	5.07	134.05	116.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	283	GLY	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2325	0	2340	26	0
1	B	2246	0	2270	14	0
1	C	2230	0	2248	14	0
1	D	2334	0	2348	14	0
2	A	4	0	0	6	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	16	0	14	1	0
4	B	16	0	13	2	0
4	C	15	0	11	0	0
4	D	16	0	14	1	0
5	A	141	0	0	2	0
5	B	115	0	0	2	0
5	C	132	0	0	3	0
5	D	122	0	0	1	0
All	All	9716	0	9258	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:13:HIS:HB2	1:A:223:PHE:CZ	2.05	0.91
1:A:13:HIS:HB2	1:A:223:PHE:CE2	2.07	0.89
1:B:206:ARG:NH2	1:B:221:ASP:OD2	2.11	0.82

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:206:ARG:NH2	1:A:221:ASP:OD2	2.26	0.68
1:C:110:LYS:O	1:C:114:GLU:HG2	1.94	0.67
1:D:206:ARG:NH2	1:D:221:ASP:OD2	2.27	0.66
1:A:110:LYS:NZ	2:A:1102:BME:C1	2.60	0.65
1:A:110:LYS:O	1:A:114:GLU:HG2	1.98	0.63
1:A:38:GLY:O	1:A:77:ILE:HA	1.98	0.62
1:D:110:LYS:O	1:D:114:GLU:HG2	2.01	0.61
1:A:100:GLN:HB3	2:A:1102:BME:C2	2.31	0.61
1:C:110:LYS:HE2	5:C:376:HOH:O	1.99	0.61
1:B:110:LYS:O	1:B:114:GLU:HG2	2.01	0.60
1:D:116:ALA:HB3	5:D:437:HOH:O	2.03	0.59
1:A:13:HIS:CB	1:A:223:PHE:CE2	2.83	0.59
1:C:16:ALA:HB2	1:C:87:LEU:HD21	1.87	0.55
1:A:235:VAL:HG21	5:A:381:HOH:O	2.06	0.55
1:C:110:LYS:CE	5:C:376:HOH:O	2.54	0.54
1:D:16:ALA:HB2	1:D:87:LEU:HD21	1.89	0.53
1:B:206:ARG:HH22	1:B:221:ASP:CG	2.14	0.53
1:A:110:LYS:HZ2	2:A:1102:BME:C1	2.21	0.53
1:C:220:LEU:HD21	1:C:239:LEU:HD13	1.90	0.53
1:C:206:ARG:HH22	1:C:221:ASP:CG	2.17	0.52
1:D:130:VAL:O	1:D:134:LEU:HG	2.10	0.52
1:D:206:ARG:HH22	1:D:221:ASP:CG	2.17	0.52
1:A:16:ALA:HB2	1:A:87:LEU:HD21	1.91	0.51
1:C:90:PRO:HB2	1:C:215:ILE:HD12	1.93	0.51
1:C:56:LEU:O	5:C:362:HOH:O	2.19	0.50
1:D:218:GLU:O	1:D:221:ASP:HB2	2.12	0.50
4:D:502:DNB:C6	4:D:502:DNB:H5'	2.42	0.50
1:A:110:LYS:HZ1	2:A:1102:BME:C2	2.25	0.49
1:B:261:VAL:HG22	1:B:276:VAL:HG22	1.92	0.49
1:B:16:ALA:HB2	1:B:87:LEU:HD21	1.93	0.49
1:B:90:PRO:HB2	1:B:215:ILE:HD12	1.94	0.49
1:D:211:PRO:O	1:D:215:ILE:HG12	2.13	0.48
1:D:239:LEU:HG	1:D:242:PRO:HG2	1.96	0.48
1:A:110:LYS:NZ	2:A:1102:BME:S2	2.70	0.48
1:A:223:PHE:O	1:A:224:LEU:HG	2.14	0.47
1:A:222:PHE:C	1:A:224:LEU:H	2.23	0.47
1:D:222:PHE:O	1:D:223:PHE:C	2.58	0.47
1:C:211:PRO:O	1:C:215:ILE:HG12	2.15	0.46
1:B:115:SER:OG	1:B:117:GLU:O	2.32	0.45
1:A:206:ARG:HH22	1:A:221:ASP:CG	2.21	0.45
1:A:40:GLN:NE2	1:A:84:GLU:O	2.49	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:39:ASN:OD1	4:B:502:DNB:H1'	2.18	0.44
1:C:160:THR:OG1	1:C:163:ALA:HB3	2.16	0.44
1:B:255:THR:HA	5:B:487:HOH:O	2.17	0.44
1:D:90:PRO:HB2	1:D:215:ILE:HD12	2.00	0.44
1:B:192:THR:HB	1:B:240:HIS:HA	1.99	0.43
1:A:110:LYS:HZ1	2:A:1102:BME:C1	2.26	0.43
1:B:72:MET:HE2	1:B:72:MET:HB3	1.86	0.43
1:C:231:LYS:H	1:C:231:LYS:HG2	1.68	0.43
4:B:502:DNB:H6	4:B:502:DNB:H2'	1.88	0.42
1:A:90:PRO:HB2	1:A:215:ILE:HD12	2.01	0.42
1:C:41:THR:HB	1:C:42:PRO:HD2	2.02	0.42
1:C:72:MET:HE2	1:C:72:MET:HB3	1.95	0.42
1:B:8:ASP:O	1:B:123:SER:HA	2.20	0.42
1:A:27:GLU:CD	1:A:306:ARG:HH12	2.28	0.42
1:C:202:THR:HG21	1:C:221:ASP:OD1	2.20	0.42
1:A:50:LEU:HD11	1:A:99:PRO:HD3	2.02	0.41
1:A:237:ALA:HA	1:A:238:PRO:HD2	1.94	0.41
1:A:150:MET:C	1:A:150:MET:SD	3.03	0.41
1:D:261:VAL:HG22	1:D:276:VAL:HG22	2.03	0.41
1:D:72:MET:HE2	1:D:72:MET:HB3	1.98	0.41
4:A:502:DNB:H5'	4:A:502:DNB:C2	2.51	0.41
1:A:218:GLU:HG2	5:A:410:HOH:O	2.21	0.41
1:D:31:LYS:HA	1:D:31:LYS:HD2	1.90	0.41
1:B:50:LEU:HD11	1:B:99:PRO:HD3	2.03	0.40
1:A:74:GLU:OE1	1:A:74:GLU:HA	2.21	0.40
1:A:5:ILE:HD13	1:A:5:ILE:HG21	1.90	0.40
1:B:271:GLN:NE2	5:B:497:HOH:O	2.55	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	302/312 (97%)	293 (97%)	6 (2%)	3 (1%)	12	9
1	B	289/312 (93%)	283 (98%)	5 (2%)	1 (0%)	36	36
1	C	287/312 (92%)	280 (98%)	6 (2%)	1 (0%)	36	36
1	D	302/312 (97%)	292 (97%)	8 (3%)	2 (1%)	18	15
All	All	1180/1248 (95%)	1148 (97%)	25 (2%)	7 (1%)	21	18

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	223	PHE
1	D	223	PHE
1	A	79	ASP
1	C	38	GLY
1	D	38	GLY
1	A	38	GLY
1	B	38	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	253/258 (98%)	252 (100%)	1 (0%)	84	89
1	B	244/258 (95%)	243 (100%)	1 (0%)	84	89
1	C	242/258 (94%)	239 (99%)	3 (1%)	63	72
1	D	254/258 (98%)	252 (99%)	2 (1%)	73	81
All	All	993/1032 (96%)	986 (99%)	7 (1%)	76	83

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	303	LEU
1	B	5	ILE
1	C	150	MET
1	C	190	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	303	LEU
1	D	76	ILE
1	D	303	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	57	ASN
1	A	179	GLN
1	B	135	ASN
1	B	179	GLN
1	D	227	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 4 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	DNB	C	502	3	16,16,18	1.14	1 (6%)	18,22,26	2.56	5 (27%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BME	A	1102	1	3,3,3	4.62	3 (100%)	2,2,2	2.61	2 (100%)
4	DNB	A	502	3	17,17,18	0.98	0	20,24,26	1.77	5 (25%)
4	DNB	D	502	3	17,17,18	1.19	3 (17%)	20,24,26	2.55	6 (30%)
4	DNB	B	502	3	17,17,18	1.16	2 (11%)	20,24,26	1.87	3 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	DNB	C	502	3	-	5/6/22/22	0/2/2/2
2	BME	A	1102	1	-	1/1/1/1	-
4	DNB	A	502	3	-	4/6/22/22	0/2/2/2
4	DNB	D	502	3	-	5/6/22/22	0/2/2/2
4	DNB	B	502	3	-	2/6/22/22	0/2/2/2

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1102	BME	C2-C1	-5.75	1.12	1.50
2	A	1102	BME	C2-S2	-5.15	1.59	1.80
4	D	502	DNB	C4'-N4'	-2.68	1.44	1.48
4	C	502	DNB	C1-C1'	-2.33	1.48	1.52
4	B	502	DNB	C1-C1'	-2.27	1.49	1.52
2	A	1102	BME	O1-C1	-2.12	1.31	1.42
4	D	502	DNB	C1-C1'	-2.05	1.49	1.52
4	D	502	DNB	C1'-N4'	-2.03	1.45	1.47
4	B	502	DNB	C5'-C4'	2.01	1.56	1.52

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	502	DNB	C1-C1'-N4'	-8.38	99.69	112.98
4	C	502	DNB	C1-C1'-N4'	-8.29	99.83	112.98
4	B	502	DNB	C1-C1'-N4'	-5.99	103.49	112.98
4	D	502	DNB	C2-C1-C1'	-4.19	113.61	120.48
4	B	502	DNB	C2'-C3'-C4'	3.83	108.35	102.55
4	A	502	DNB	C1-C1'-N4'	-3.80	106.95	112.98
4	C	502	DNB	C2'-C3'-C4'	3.49	107.84	102.55

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	502	DNB	C2-C1-C1'	3.46	126.16	120.48
4	D	502	DNB	C6-C1-C1'	3.20	125.86	120.78
4	D	502	DNB	C2'-C3'-C4'	2.88	106.91	102.55
4	C	502	DNB	O2'-C2'-C1'	2.86	116.04	110.47
2	A	1102	BME	O1-C1-C2	2.84	121.90	110.82
4	D	502	DNB	O3'-C3'-C4'	-2.67	105.79	112.92
4	A	502	DNB	C6-C1-C1'	-2.66	116.55	120.78
4	D	502	DNB	C5'-C4'-C3'	2.54	118.06	113.68
4	A	502	DNB	C5-C6-C1	2.46	123.60	120.65
4	C	502	DNB	C5'-C4'-N4'	2.43	115.75	111.47
2	A	1102	BME	C1-C2-S2	2.36	128.39	111.57
4	C	502	DNB	O5'-C5'-C4'	2.26	116.58	111.10
4	B	502	DNB	C2-C1-C1'	-2.08	117.06	120.48
4	A	502	DNB	C4-C3-N3	-2.05	117.13	120.90

There are no chirality outliers.

All (17) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	502	DNB	C6-C1-C1'-C2'
4	A	502	DNB	C2-C1-C1'-C2'
4	B	502	DNB	C6-C1-C1'-C2'
4	B	502	DNB	C2-C1-C1'-C2'
4	C	502	DNB	C6-C1-C1'-C2'
4	C	502	DNB	C2-C1-C1'-C2'
4	C	502	DNB	C3'-C4'-C5'-O5'
4	C	502	DNB	N4'-C4'-C5'-O5'
4	D	502	DNB	C6-C1-C1'-C2'
4	D	502	DNB	C2-C1-C1'-C2'
2	A	1102	BME	O1-C1-C2-S2
4	D	502	DNB	C6-C1-C1'-N4'
4	A	502	DNB	C2-C1-C1'-N4'
4	D	502	DNB	C2-C1-C1'-N4'
4	A	502	DNB	C6-C1-C1'-N4'
4	D	502	DNB	N4'-C4'-C5'-O5'
4	C	502	DNB	C2-C1-C1'-N4'

There are no ring outliers.

4 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1102	BME	6	0
4	A	502	DNB	1	0
4	D	502	DNB	1	0
4	B	502	DNB	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	305/312 (97%)	0.57	23 (7%)	20	21	15, 28, 52, 67	1 (0%)
1	B	295/312 (94%)	0.73	22 (7%)	20	21	21, 27, 41, 58	0
1	C	293/312 (93%)	1.04	38 (12%)	7	7	21, 27, 40, 62	0
1	D	304/312 (97%)	0.74	23 (7%)	20	21	16, 28, 48, 79	2 (0%)
All	All	1197/1248 (95%)	0.77	106 (8%)	15	16	15, 27, 47, 79	3 (0%)

All (106) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	86	GLY	8.2
1	A	78	ALA	5.3
1	D	116	ALA	5.0
1	C	39	ASN	4.9
1	A	82	HIS	4.6
1	C	89	GLY	4.5
1	C	88	ASP	4.4
1	A	224	LEU	4.3
1	C	232	TRP	4.3
1	A	236	GLY	4.3
1	B	222	PHE	4.2
1	D	226	TYR	4.1
1	B	39	ASN	4.0
1	D	86	GLY	3.9
1	A	77	ILE	3.8
1	B	224	LEU	3.8
1	A	79	ASP	3.8
1	D	224	LEU	3.7
1	D	76	ILE	3.7
1	C	101	ASN	3.6
1	C	209	GLY	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	223	PHE	3.5
1	A	81	VAL	3.5
1	C	90	PRO	3.5
1	C	91	ALA	3.5
1	D	227	HIS	3.5
1	D	232	TRP	3.4
1	A	83	GLY	3.4
1	C	87	LEU	3.4
1	A	234	PHE	3.4
1	A	86	GLY	3.3
1	D	0	GLY	3.3
1	B	86	GLY	3.3
1	D	229	ASP	3.3
1	A	232	TRP	3.2
1	D	223	PHE	3.2
1	C	218	GLU	3.2
1	A	223	PHE	3.1
1	C	199	VAL	3.1
1	C	40	GLN	3.1
1	C	222	PHE	3.0
1	C	234	PHE	3.0
1	C	212	VAL	3.0
1	B	200	GLU	3.0
1	D	1	SER	2.9
1	A	76	ILE	2.9
1	B	101	ASN	2.9
1	C	283	GLY	2.9
1	D	282	THR	2.8
1	B	104	ALA	2.8
1	A	87	LEU	2.8
1	B	87	LEU	2.8
1	C	202	THR	2.8
1	D	117	GLU	2.7
1	B	232	TRP	2.7
1	C	235	VAL	2.7
1	C	236	GLY	2.7
1	D	233	GLY	2.7
1	B	90	PRO	2.6
1	A	84	GLU	2.6
1	B	89	GLY	2.6
1	D	285	LYS	2.5
1	C	113	ARG	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	311	ALA	2.5
1	D	230	GLU	2.5
1	C	38	GLY	2.5
1	D	228	LYS	2.5
1	C	2	ALA	2.5
1	C	200	GLU	2.5
1	C	13	HIS	2.5
1	D	38	GLY	2.5
1	C	230	GLU	2.5
1	B	234	PHE	2.5
1	D	283	GLY	2.5
1	D	234	PHE	2.4
1	A	91	ALA	2.4
1	D	101	ASN	2.4
1	A	235	VAL	2.4
1	B	1	SER	2.4
1	C	116	ALA	2.3
1	A	101	ASN	2.3
1	B	13	HIS	2.3
1	B	88	ASP	2.3
1	B	40	GLN	2.3
1	B	199	VAL	2.3
1	A	285	LYS	2.3
1	B	283	GLY	2.3
1	C	281	LEU	2.2
1	C	117	GLU	2.2
1	C	237	ALA	2.2
1	A	92	LEU	2.2
1	B	235	VAL	2.2
1	C	179	GLN	2.2
1	C	280	TYR	2.1
1	B	305	ASP	2.1
1	A	90	PRO	2.1
1	C	282	THR	2.1
1	D	85	SER	2.1
1	A	88	ASP	2.1
1	C	197	ILE	2.1
1	C	14	ASP	2.1
1	C	198	HIS	2.1
1	A	311	ALA	2.1
1	D	237	ALA	2.0
1	B	103	THR	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	26	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	BME	A	1102	4/4	0.78	0.24	59,60,60,61	0
4	DNB	C	502	15/17	0.82	0.19	23,34,41,41	15
4	DNB	B	502	16/17	0.85	0.16	16,27,33,33	16
4	DNB	A	502	16/17	0.90	0.14	16,29,38,39	0
4	DNB	D	502	16/17	0.91	0.12	30,40,45,46	0
3	CA	D	501	1/1	0.98	0.10	20,20,20,20	0
3	CA	C	501	1/1	0.99	0.13	19,19,19,19	0
3	CA	A	501	1/1	0.99	0.13	16,16,16,16	0
3	CA	B	501	1/1	0.99	0.11	19,19,19,19	0

6.5 Other polymers [i](#)

There are no such residues in this entry.