



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 05:34 PM UTC

PDB ID : 3G5K / pdb_00003g5k
Title : Structure and activity of human mitochondrial peptide deformylase, a novel cancer target
Authors : Escobar-Alvarez, S.; Goldgur, Y.; Yang, G.; Ouerfelli, O.; Li, Y.; Scheinberg, D.A.
Deposited on : 2009-02-05
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5551 (1.70-1.70)
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)
RSRZ outliers	180081	5554 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	183	2% 86% 13% .
1	B	183	3% 84% 13% .
1	C	183	4% 85% 14% .
1	D	183	3% 86% 14% .

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6445 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Peptide deformylase, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	183	Total	C	N	O	S	0	0	0
			1445	909	272	251	13			
1	B	183	Total	C	N	O	S	0	0	0
			1445	909	272	251	13			
1	C	183	Total	C	N	O	S	0	0	0
			1445	909	272	251	13			
1	D	183	Total	C	N	O	S	0	0	0
			1445	909	272	251	13			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	3	HIS	-	expression tag	UNP Q9HBH1
A	4	MET	-	expression tag	UNP Q9HBH1
A	5	SER	-	expression tag	UNP Q9HBH1
B	3	HIS	-	expression tag	UNP Q9HBH1
B	4	MET	-	expression tag	UNP Q9HBH1
B	5	SER	-	expression tag	UNP Q9HBH1
C	3	HIS	-	expression tag	UNP Q9HBH1
C	4	MET	-	expression tag	UNP Q9HBH1
C	5	SER	-	expression tag	UNP Q9HBH1
D	3	HIS	-	expression tag	UNP Q9HBH1
D	4	MET	-	expression tag	UNP Q9HBH1
D	5	SER	-	expression tag	UNP Q9HBH1

- Molecule 2 is ACTINONIN (CCD ID: BB2) (formula: C₁₉H₃₅N₃O₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			27	19	3	5		
2	B	1	Total	C	N	O	0	0
			27	19	3	5		
2	C	1	Total	C	N	O	0	0
			27	19	3	5		
2	D	1	Total	C	N	O	0	0
			27	19	3	5		

- Molecule 3 is COBALT (II) ION (CCD ID: CO) (formula: Co).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Co	0	0
			1	1		
3	B	1	Total	Co	0	0
			1	1		
3	C	1	Total	Co	0	0
			1	1		
3	D	1	Total	Co	0	0
			1	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	153	Total	O	0	0
			153	153		

Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	129	Total 129	O 129	0	0
4	C	138	Total 138	O 138	0	0
4	D	133	Total 133	O 133	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Peptide deformylase, mitochondrial




- Molecule 1: Peptide deformylase, mitochondrial



- Molecule 1: Peptide deformylase, mitochondrial



- Molecule 1: Peptide deformylase, mitochondrial



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	116.14Å 77.83Å 110.53Å 90.00° 107.46° 90.00°	Depositor
Resolution (Å)	35.00 – 1.70 35.00 – 1.70	Depositor EDS
% Data completeness (in resolution range)	99.5 (35.00-1.70) 99.8 (35.00-1.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.78 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5.5.0070	Depositor
R, R_{free}	0.164 , 0.193 0.165 , 0.195	Depositor DCC
R_{free} test set	5160 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	16.6	Xtriage
Anisotropy	0.306	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 38.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6445	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.86% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BB2, CO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.73	9/1477 (0.6%)	1.48	6/2002 (0.3%)
1	B	1.80	16/1477 (1.1%)	1.48	10/2002 (0.5%)
1	C	1.77	12/1477 (0.8%)	1.46	2/2002 (0.1%)
1	D	1.77	12/1477 (0.8%)	1.44	4/2002 (0.2%)
All	All	1.77	49/5908 (0.8%)	1.47	22/8008 (0.3%)

All (49) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	112	GLU	CB-CG	-9.22	1.24	1.52
1	B	93	ARG	CZ-NH2	8.57	1.44	1.33
1	B	55	ALA	CA-CB	7.64	1.65	1.54
1	D	97	ASN	C-O	6.87	1.27	1.23
1	C	126	ARG	CZ-NH1	6.64	1.42	1.32
1	A	94	VAL	C-O	6.56	1.31	1.24
1	C	171	ASP	CG-OD1	6.51	1.37	1.25
1	A	55	ALA	CA-CB	6.48	1.62	1.53
1	D	180	TRP	N-CA	6.43	1.53	1.46
1	D	107	LEU	N-CA	6.34	1.53	1.46
1	B	114	CYS	N-CA	6.29	1.53	1.46
1	C	114	CYS	N-CA	5.87	1.53	1.46
1	D	120	PHE	N-CA	5.79	1.53	1.45
1	B	162	GLN	CG-CD	-5.75	1.37	1.52
1	C	113	GLY	N-CA	5.66	1.50	1.44
1	B	109	THR	C-O	5.64	1.30	1.24
1	D	157	GLU	CD-OE2	-5.64	1.14	1.25
1	B	141	GLN	N-CA	5.64	1.53	1.46
1	B	25	GLU	N-CA	5.58	1.53	1.45
1	A	138	ASN	N-CA	5.55	1.53	1.46
1	B	24	VAL	CA-CB	-5.55	1.47	1.54
1	D	84	LEU	N-CA	5.55	1.53	1.46

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	94	VAL	C-O	5.51	1.30	1.24
1	D	150	ALA	N-CA	5.45	1.52	1.46
1	A	147	SER	N-CA	5.45	1.53	1.45
1	B	93	ARG	CD-NE	5.43	1.53	1.46
1	C	153	ILE	CG1-CD1	5.43	1.73	1.51
1	B	97	ASN	CA-CB	5.40	1.57	1.52
1	D	178	VAL	CA-CB	5.38	1.62	1.54
1	C	143	VAL	N-CA	-5.36	1.40	1.46
1	C	165	LEU	CA-CB	5.34	1.62	1.53
1	A	33	GLU	C-O	5.32	1.30	1.24
1	B	8	HIS	C-O	5.31	1.30	1.23
1	A	141	GLN	N-CA	5.29	1.52	1.46
1	D	132	ILE	CA-CB	5.24	1.60	1.54
1	C	167	ILE	N-CA	5.23	1.53	1.46
1	D	111	PRO	C-O	5.21	1.29	1.23
1	B	93	ARG	CZ-NH1	5.21	1.40	1.32
1	A	78	PRO	CA-C	5.18	1.54	1.51
1	D	92	LEU	C-O	5.18	1.30	1.23
1	B	67	LEU	CA-CB	5.16	1.62	1.53
1	C	92	LEU	CA-C	5.15	1.59	1.52
1	B	4	MET	N-CA	5.12	1.52	1.46
1	C	74	CYS	CB-SG	-5.12	1.64	1.81
1	C	147	SER	CA-C	5.08	1.59	1.52
1	A	111	PRO	C-O	5.07	1.29	1.23
1	A	129	ALA	N-CA	5.02	1.52	1.46
1	C	108	VAL	CA-C	5.01	1.58	1.52
1	B	37	LEU	N-CA	5.00	1.52	1.46

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	93	ARG	NE-CZ-NH1	-10.56	110.94	121.50
1	A	142	VAL	N-CA-C	-7.90	96.69	108.46
1	B	19	GLY	CA-C-N	-6.78	114.31	123.12
1	B	19	GLY	C-N-CA	-6.78	114.31	123.12
1	B	93	ARG	NH1-CZ-NH2	6.75	128.08	119.30
1	D	157	GLU	CG-CD-OE2	-6.47	103.53	118.40
1	C	138	ASN	N-CA-C	6.20	119.01	111.82
1	C	136	ASP	N-CA-CB	6.06	117.43	109.55
1	B	13	GLY	N-CA-C	-5.83	107.34	113.58
1	A	78	PRO	O-C-N	5.81	123.98	121.31
1	A	26	ARG	CG-CD-NE	-5.80	99.23	112.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	108	VAL	CA-C-N	-5.64	115.04	122.99
1	B	108	VAL	C-N-CA	-5.64	115.04	122.99
1	B	112	GLU	N-CA-CB	-5.46	101.36	111.53
1	D	78	PRO	CB-CA-C	-5.36	106.35	111.39
1	B	74	CYS	N-CA-CB	-5.28	102.34	110.16
1	B	47	ARG	NE-CZ-NH1	-5.20	116.31	121.50
1	D	108	VAL	O-C-N	5.18	128.80	123.20
1	A	85	ARG	CD-NE-CZ	5.15	131.60	124.40
1	A	88	GLU	CA-C-N	-5.14	114.94	120.03
1	A	88	GLU	C-N-CA	-5.14	114.94	120.03
1	D	157	GLU	CG-CD-OE1	5.11	130.15	118.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1445	0	1456	7	1
1	B	1445	0	1456	14	0
1	C	1445	0	1456	9	1
1	D	1445	0	1456	9	0
2	A	27	0	34	0	0
2	B	27	0	35	0	0
2	C	27	0	34	0	0
2	D	27	0	34	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	153	0	0	0	0
4	B	129	0	0	5	0
4	C	138	0	0	3	0
4	D	133	0	0	2	0
All	All	6445	0	5961	39	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:40:ARG:NH2	4:B:1047:HOH:O	2.01	0.93
1:D:81:GLN:HG3	1:D:85:ARG:NH1	1.93	0.83
1:A:35:GLN:OE1	1:A:39:GLN:NE2	2.23	0.71
1:C:80:ARG:O	1:C:80:ARG:HD2	1.90	0.71
1:B:100:LEU:HD23	1:B:101:ARG:N	2.06	0.71
1:D:7:SER:OG	1:D:40:ARG:HD2	1.92	0.69
1:A:86:GLN:HE22	1:A:128:GLN:HE22	1.41	0.69
1:C:48:ARG:NH1	4:C:1061:HOH:O	2.28	0.66
1:B:85:ARG:CZ	4:B:1069:HOH:O	2.44	0.65
1:B:100:LEU:HD23	1:B:100:LEU:C	2.23	0.64
1:D:131:GLN:HE21	1:D:145:GLN:HE21	1.44	0.64
1:D:131:GLN:NE2	1:D:145:GLN:HE21	1.96	0.62
1:C:4:MET:HA	4:C:1102:HOH:O	2.00	0.62
1:B:85:ARG:NH1	4:B:1069:HOH:O	2.35	0.59
1:B:43:GLN:HE21	1:B:47:ARG:HE	1.52	0.58
1:B:85:ARG:NH2	4:B:1069:HOH:O	2.36	0.58
1:C:131:GLN:HB3	1:C:145:GLN:HG2	1.87	0.57
1:A:63:GLN:HE21	1:A:97:ASN:HA	1.72	0.55
1:C:43:GLN:HE21	1:C:47:ARG:HE	1.55	0.55
1:B:90:PHE:HB2	1:B:93:ARG:HH12	1.73	0.53
1:A:81:GLN:HG3	1:A:85:ARG:NH1	2.24	0.53
1:D:47:ARG:HD3	4:D:1088:HOH:O	2.08	0.52
1:C:65:LEU:C	1:C:65:LEU:HD12	2.38	0.48
1:A:86:GLN:NE2	1:A:128:GLN:HE22	2.09	0.47
1:D:140:GLU:HB3	4:D:1039:HOH:O	2.15	0.47
1:A:136:ASP:HB2	1:A:137:PRO:CD	2.46	0.46
1:D:43:GLN:HE21	1:D:47:ARG:HH21	1.64	0.44
1:C:80:ARG:HD2	1:C:80:ARG:C	2.43	0.43
1:B:65:LEU:HD12	1:B:65:LEU:C	2.43	0.43
1:B:136:ASP:OD1	1:B:136:ASP:C	2.61	0.43
1:C:171:ASP:HB2	4:C:1117:HOH:O	2.18	0.43
1:D:63:GLN:HE21	1:D:97:ASN:HA	1.83	0.42
1:B:25:GLU:HB2	1:B:28:GLN:OE1	2.18	0.42
1:A:136:ASP:HB2	1:A:137:PRO:HD2	2.02	0.42
1:B:111:PRO:HD2	4:B:1069:HOH:O	2.21	0.41
1:C:7:SER:OG	1:C:40:ARG:HD2	2.21	0.41
1:B:100:LEU:C	1:B:100:LEU:CD2	2.93	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:65:LEU:C	1:D:65:LEU:HD12	2.46	0.40
1:B:97:ASN:N	1:B:98:PRO:CD	2.85	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:185:ASP:OD2	1:C:3:HIS:O[4_555]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	181/183 (99%)	180 (99%)	1 (1%)	0	100	100
1	B	181/183 (99%)	180 (99%)	1 (1%)	0	100	100
1	C	181/183 (99%)	180 (99%)	1 (1%)	0	100	100
1	D	181/183 (99%)	180 (99%)	1 (1%)	0	100	100
All	All	724/732 (99%)	720 (99%)	4 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	158/158 (100%)	156 (99%)	2 (1%)	61	48
1	B	158/158 (100%)	156 (99%)	2 (1%)	61	48
1	C	158/158 (100%)	155 (98%)	3 (2%)	50	34
1	D	158/158 (100%)	157 (99%)	1 (1%)	78	72
All	All	632/632 (100%)	624 (99%)	8 (1%)	61	48

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	47	ARG
1	A	56	PRO
1	B	4	MET
1	B	100	LEU
1	C	73	LEU
1	C	136	ASP
1	C	157	GLU
1	D	56	PRO

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	39	GLN
1	A	63	GLN
1	A	86	GLN
1	A	138	ASN
1	A	177	ASN
1	B	43	GLN
1	B	63	GLN
1	C	43	GLN
1	C	81	GLN
1	D	43	GLN
1	D	63	GLN
1	D	81	GLN
1	D	131	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	BB2	D	1004	3	27,27,27	1.89	5 (18%)	34,35,35	1.98	11 (32%)
2	BB2	A	1001	3	27,27,27	1.96	7 (25%)	34,35,35	3.12	11 (32%)
2	BB2	C	1003	3	27,27,27	1.96	6 (22%)	34,35,35	2.00	8 (23%)
2	BB2	B	1002	3	27,27,27	2.31	10 (37%)	34,35,35	2.91	12 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BB2	D	1004	3	-	0/33/43/43	0/1/1/1
2	BB2	A	1001	3	-	2/33/43/43	0/1/1/1
2	BB2	C	1003	3	-	1/33/43/43	0/1/1/1
2	BB2	B	1002	3	-	1/33/43/43	0/1/1/1

All (28) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	1003	BB2	C3-N1	-7.57	1.24	1.32

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	1004	BB2	C3-N1	-7.15	1.24	1.32
2	B	1002	BB2	C3-N1	-6.45	1.25	1.32
2	A	1001	BB2	C6-C12	-5.02	1.42	1.51
2	B	1002	BB2	C6-C12	-5.01	1.43	1.51
2	A	1001	BB2	C3-N1	-4.21	1.27	1.32
2	B	1002	BB2	O20-C19	3.48	1.28	1.22
2	A	1001	BB2	C12-N14	3.25	1.41	1.34
2	D	1004	BB2	O20-C19	3.00	1.27	1.22
2	B	1002	BB2	C25-C22	2.96	1.60	1.53
2	A	1001	BB2	O20-C19	2.95	1.27	1.22
2	B	1002	BB2	C5-C6	2.85	1.59	1.53
2	B	1002	BB2	C24-C23	2.77	1.61	1.51
2	B	1002	BB2	O2-N1	2.70	1.46	1.40
2	C	1003	BB2	C23-N21	2.67	1.52	1.47
2	B	1002	BB2	C23-N21	2.52	1.52	1.47
2	D	1004	BB2	C23-N21	2.43	1.52	1.47
2	C	1003	BB2	O2-N1	2.40	1.45	1.40
2	D	1004	BB2	C5-C3	-2.37	1.46	1.51
2	C	1003	BB2	C8-C7	2.35	1.62	1.52
2	A	1001	BB2	O4-C3	2.33	1.27	1.23
2	C	1003	BB2	C25-C22	2.31	1.59	1.53
2	C	1003	BB2	C15-N14	2.25	1.50	1.45
2	B	1002	BB2	C15-N14	2.25	1.50	1.45
2	A	1001	BB2	C23-N21	2.23	1.51	1.47
2	D	1004	BB2	O4-C3	2.23	1.27	1.23
2	A	1001	BB2	C25-C22	2.22	1.58	1.53
2	B	1002	BB2	C8-C7	2.04	1.60	1.52

All (42) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1001	BB2	O4-C3-N1	-13.31	106.93	123.27
2	B	1002	BB2	O4-C3-N1	-12.45	107.99	123.27
2	C	1003	BB2	O2-N1-C3	-5.94	111.02	119.79
2	C	1003	BB2	O4-C3-N1	-5.61	116.39	123.27
2	A	1001	BB2	C5-C3-N1	5.33	123.22	115.14
2	A	1001	BB2	O4-C3-C5	5.26	129.22	121.54
2	D	1004	BB2	O4-C3-N1	-5.16	116.93	123.27
2	D	1004	BB2	O2-N1-C3	-4.75	112.77	119.79
2	B	1002	BB2	C5-C3-N1	4.69	122.24	115.14
2	A	1001	BB2	C17-C16-C18	4.60	123.23	110.58
2	B	1002	BB2	O27-C26-C22	-4.13	95.52	111.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1002	BB2	C24-C23-N21	-3.94	96.46	103.24
2	A	1001	BB2	O13-C12-N14	-3.55	116.61	122.96
2	C	1003	BB2	O27-C26-C22	-3.49	97.98	111.51
2	A	1001	BB2	C6-C12-N14	3.49	122.20	116.19
2	C	1003	BB2	C23-N21-C22	3.46	117.53	111.44
2	B	1002	BB2	O4-C3-C5	3.33	126.41	121.54
2	D	1004	BB2	C18-C16-C15	-3.29	102.14	111.16
2	B	1002	BB2	O13-C12-N14	-3.26	117.12	122.96
2	C	1003	BB2	C5-C3-N1	3.14	119.90	115.14
2	B	1002	BB2	C7-C6-C12	-3.01	104.74	109.63
2	D	1004	BB2	O20-C19-N21	-2.99	116.01	121.38
2	D	1004	BB2	C5-C3-N1	2.86	119.47	115.14
2	B	1002	BB2	O13-C12-C6	2.78	126.49	122.19
2	D	1004	BB2	C25-C22-C26	-2.57	105.85	113.07
2	A	1001	BB2	O2-N1-C3	-2.53	116.05	119.79
2	B	1002	BB2	O20-C19-C15	2.53	124.71	119.96
2	D	1004	BB2	C6-C5-C3	-2.52	107.38	112.33
2	A	1001	BB2	O27-C26-C22	-2.52	101.75	111.51
2	D	1004	BB2	O27-C26-C22	-2.45	102.03	111.51
2	C	1003	BB2	C5-C6-C12	-2.44	106.24	109.71
2	A	1001	BB2	C7-C6-C5	2.41	118.00	110.30
2	B	1002	BB2	O2-N1-C3	-2.39	116.25	119.79
2	D	1004	BB2	O13-C12-N14	-2.36	118.72	122.96
2	C	1003	BB2	O20-C19-N21	-2.34	117.17	121.38
2	B	1002	BB2	C25-C22-C26	-2.31	106.58	113.07
2	D	1004	BB2	C16-C15-C19	-2.30	106.21	110.71
2	C	1003	BB2	O20-C19-C15	2.27	124.22	119.96
2	D	1004	BB2	O20-C19-C15	2.25	124.18	119.96
2	B	1002	BB2	C26-C22-N21	-2.17	103.18	112.51
2	A	1001	BB2	C24-C23-N21	2.08	106.82	103.24
2	A	1001	BB2	C17-C16-C15	-2.04	105.56	111.16

There are no chirality outliers.


All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	1003	BB2	N1-C3-C5-C6
2	A	1001	BB2	N1-C3-C5-C6
2	A	1001	BB2	O4-C3-C5-C6
2	B	1002	BB2	N1-C3-C5-C6


There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	183/183 (100%)	-0.36	3 (1%) 70 75	6, 12, 33, 59	0
1	B	183/183 (100%)	-0.18	6 (3%) 49 53	5, 12, 34, 68	0
1	C	183/183 (100%)	-0.12	8 (4%) 39 42	6, 13, 37, 68	0
1	D	183/183 (100%)	-0.15	6 (3%) 49 53	6, 14, 34, 58	0
All	All	732/732 (100%)	-0.20	23 (3%) 51 55	5, 13, 35, 68	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	4	MET	6.7
1	B	3	HIS	6.6
1	A	3	HIS	4.9
1	C	3	HIS	4.3
1	D	3	HIS	4.0
1	C	136	ASP	3.7
1	D	185	ASP	3.3
1	C	4	MET	3.1
1	B	5	SER	3.0
1	D	4	MET	3.0
1	A	4	MET	3.0
1	C	138	ASN	3.0
1	C	137	PRO	2.9
1	A	185	ASP	2.7
1	C	80	ARG	2.6
1	C	139	GLY	2.6
1	D	27	ALA	2.5
1	B	185	ASP	2.3
1	C	5	SER	2.2
1	D	140	GLU	2.1
1	B	75	ARG	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	36	ARG	2.1
1	B	80	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	BB2	B	1002	27/27	0.87	0.13	6,22,33,40	0
2	BB2	A	1001	27/27	0.90	0.12	7,21,28,31	0
2	BB2	C	1003	27/27	0.94	0.09	10,17,25,36	0
2	BB2	D	1004	27/27	0.95	0.08	9,17,27,29	0
3	CO	B	1003	1/1	0.99	0.05	9,9,9,9	0
3	CO	D	2	1/1	0.99	0.04	10,10,10,10	0
3	CO	C	1	1/1	1.00	0.05	8,8,8,8	0
3	CO	A	1002	1/1	1.00	0.04	9,9,9,9	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around BB2 B 1002:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around BB2 A 1001:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around BB2 C 1003:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around BB2 D 1004:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.