



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 06:21 AM UTC

PDB ID : 1G67 / pdb_00001g67
Title : THIAMIN PHOSPHATE SYNTHASE
Authors : Peapus, D.H.; Chiu, H.-J.; Campobasso, N.; Reddick, J.J.; Begley, T.P.; Ealick, S.E.
Deposited on : 2000-11-03
Resolution : 1.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2563 (1.40-1.40)
Clashscore	190562	2660 (1.40-1.40)
Ramachandran outliers	187476	2611 (1.40-1.40)
Sidechain outliers	187428	2610 (1.40-1.40)
RSRZ outliers	180081	2561 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	225	12% 88% 11%
1	B	225	11% 83% 17%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 3723 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called THIAMIN PHOSPHATE SYNTHASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	225	Total	C	N	O	S	0	3	0
			1688	1047	298	335	8			
1	B	225	Total	C	N	O	S	0	6	0
			1700	1057	303	332	8			


There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	11	GLY	-	cloning artifact	UNP P39594
A	12	ILE	-	cloning artifact	UNP P39594
A	13	ARG	-	cloning artifact	UNP P39594
A	130	ALA	SER	engineered mutation	UNP P39594
B	1011	GLY	-	cloning artifact	UNP P39594
B	1012	ILE	-	cloning artifact	UNP P39594
B	1013	ARG	-	cloning artifact	UNP P39594
B	1130	ALA	SER	engineered mutation	UNP P39594

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		
2	B	1	Total	Mg	0	0
			1	1		

- Molecule 3 is 2-METHYL-5-METHYLENE-5H-PYRIMIDIN-4-YLIDENEAMINE (CCD ID: ICP) (formula: C₆H₇N₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	N	0	0
			9	6	3		
3	B	1	Total	C	N	0	0
			9	6	3		

- Molecule 4 is PYROPHOSPHATE 2- (CCD ID: POP) (formula: $\text{H}_2\text{O}_7\text{P}_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	P	0	1
			13	10	3		
4	B	1	Total	O	P	0	1
			13	10	3		

- Molecule 5 is 4-METHYL-5-HYDROXYETHYLTHIAZOLE PHOSPHATE (CCD ID: TZP) (formula: $C_6H_{10}NO_4PS$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	S	0	0
			13	6	1	4	1	1		
5	B	1	Total	C	N	O	P	S	0	0
			13	6	1	4	1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	132	Total	O	0	0
			132	132		
6	B	131	Total	O	0	0
			131	131		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: THIAMIN PHOSPHATE SYNTHASE



• Molecule 1: THIAMIN PHOSPHATE SYNTHASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	76.98Å 76.98Å 140.15Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.99 – 1.40 42.99 – 1.40	Depositor EDS
% Data completeness (in resolution range)	77.0 (42.99-1.40) 77.1 (42.99-1.40)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.90 (at 1.40Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.216 , 0.240 0.221 , 0.242	Depositor DCC
R_{free} test set	3240 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	13.4	Xtriage
Anisotropy	0.035	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 39.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3723	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 37.36 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.3396e-04.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: POP, MG, ICP, TZP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.35	0/1721	0.79	2/2317 (0.1%)
1	B	0.35	0/1750	0.80	2/2356 (0.1%)
All	All	0.35	0/3471	0.80	4/4673 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	1209	SER	N-CA-C	7.12	119.04	111.28
1	A	209	SER	N-CA-C	6.72	119.44	111.71
1	A	122	ILE	N-CA-C	5.69	118.36	112.90
1	B	1122	ILE	N-CA-C	5.47	117.44	112.29

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1688	0	1711	25	0
1	B	1700	0	1733	32	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	9	0	6	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	9	0	6	0	0
4	A	13	0	0	0	0
4	B	13	0	0	1	0
5	A	13	0	8	0	0
5	B	13	0	8	0	0
6	A	132	0	0	1	0
6	B	131	0	0	3	0
All	All	3723	0	3472	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:114:ASN:ND2	1:A:116:LYS:HG2	1.64	1.11
1:B:1189:ILE:HG21	1:B:1205[B]:VAL:HG11	1.40	1.02
1:A:114:ASN:HD22	1:A:116:LYS:HG2	1.35	0.85
1:A:12:ILE:H	1:A:12:ILE:HD13	1.45	0.80
1:B:1189:ILE:HG21	1:B:1205[B]:VAL:CG1	2.13	0.79
1:A:35:ASN:HD22	1:A:35:ASN:H	1.33	0.76
1:A:105:GLY:HA2	1:A:125:MET:HG2	1.70	0.74
1:B:1060:GLU:HB3	1:B:1071[B]:ARG:HH12	1.54	0.71
1:A:114:ASN:HD21	1:A:116:LYS:HG2	1.52	0.70
1:B:1225:ARG:HG3	1:B:1229:GLN:HE21	1.56	0.69
1:B:1060:GLU:HB3	1:B:1071[B]:ARG:NH1	2.08	0.68
1:A:35:ASN:H	1:A:35:ASN:ND2	1.93	0.67
1:A:12:ILE:HD13	1:A:12:ILE:N	2.13	0.64
1:A:12:ILE:H	1:A:12:ILE:CD1	2.12	0.63
1:B:1195:ALA:HB3	1:B:1196:PRO:HD3	1.81	0.63
1:A:35:ASN:HD22	1:A:35:ASN:N	1.94	0.62
1:B:1091:VAL:O	1:B:1106:ILE:HD12	1.99	0.62
1:B:1189:ILE:CG2	1:B:1205[B]:VAL:HG11	2.23	0.61
1:B:1189:ILE:CD1	1:B:1205[B]:VAL:HG13	2.32	0.60
1:A:195:ALA:HB3	1:A:196:PRO:HD3	1.85	0.58
1:B:1050:LYS:HG3	6:B:3110:HOH:O	2.02	0.58
1:B:1189:ILE:HD13	1:B:1205[B]:VAL:HG13	1.84	0.58
1:B:1232:LYS:HA	1:B:1235:ARG:HD2	1.86	0.57
1:A:134:MET:HE3	1:A:177:GLN:NE2	2.21	0.56
1:A:91:VAL:O	1:A:106:ILE:HD12	2.05	0.56
1:A:225:ARG:HG3	1:A:229:GLN:HE21	1.71	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1096:GLU:H	1:B:1096:GLU:CD	2.13	0.55
1:B:1225:ARG:O	1:B:1229:GLN:HG3	2.06	0.55
1:A:105:GLY:HA2	1:A:125:MET:CG	2.39	0.52
1:A:128:GLY:HA3	1:A:147:TYR:CZ	2.44	0.52
1:A:83:ARG:HH12	1:B:1079:GLN:NE2	2.08	0.51
1:A:76[B]:GLU:HG2	6:B:3070:HOH:O	2.11	0.49
1:B:1055:LEU:HD21	1:B:1090[B]:ILE:HD11	1.94	0.48
1:B:1177:GLN:HG3	6:B:3094:HOH:O	2.13	0.48
1:B:1224:PHE:O	1:B:1228:ILE:HG12	2.14	0.46
1:A:72:ILE:O	1:A:76[A]:GLU:HG3	2.15	0.46
1:B:1128:GLY:HA3	1:B:1147:TYR:CZ	2.50	0.46
1:B:1043:THR:CG2	1:B:1047:LYS:HE3	2.45	0.46
1:B:1082:CYS:HB3	1:B:1087:VAL:O	2.17	0.45
1:B:1055:LEU:CD1	1:B:1088:PRO:HB2	2.45	0.45
1:B:1043:THR:HG22	1:B:1047:LYS:HE3	1.99	0.45
1:A:21:MET:HG2	6:A:3099:HOH:O	2.15	0.45
1:B:1216:ASP:HB3	1:B:1219:SER:OG	2.17	0.45
1:A:226:GLU:O	1:A:230[B]:THR:HG23	2.17	0.44
1:A:35:ASN:ND2	1:A:35:ASN:N	2.58	0.43
1:B:1109:GLY:HA3	4:B:2004[B]:POP:O3	2.18	0.43
1:B:1060:GLU:CB	1:B:1071[B]:ARG:HH12	2.27	0.43
1:B:1020:GLU:HG2	1:B:1021:MET:HE2	2.00	0.42
1:A:31:ILE:HG12	1:A:57:GLN:HB3	2.00	0.42
1:A:225:ARG:O	1:A:229:GLN:HG3	2.21	0.41
1:B:1080:ALA:O	1:B:1084:GLU:HG3	2.21	0.41
1:B:1055:LEU:HD21	1:B:1090[B]:ILE:CD1	2.51	0.41
1:B:1019:ARG:HH11	1:B:1019:ARG:HG2	1.84	0.40
1:B:1225:ARG:HG3	1:B:1229:GLN:NE2	2.29	0.40
1:A:134:MET:HE3	1:A:177:GLN:HE21	1.85	0.40
1:B:1096:GLU:CD	1:B:1096:GLU:N	2.78	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	226/225 (100%)	218 (96%)	7 (3%)	1 (0%)	30	10
1	B	229/225 (102%)	221 (96%)	8 (4%)	0	100	100
All	All	455/450 (101%)	439 (96%)	15 (3%)	1 (0%)	43	19

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	64	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	175/172 (102%)	173 (99%)	2 (1%)	65	37
1	B	178/172 (104%)	177 (99%)	1 (1%)	78	56
All	All	353/344 (103%)	350 (99%)	3 (1%)	70	48

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	12	ILE
1	A	35	ASN
1	B	1192	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	35	ASN
1	A	100	ASN
1	A	110	GLN
1	A	114	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	166	GLN
1	A	229	GLN
1	B	1079	GLN
1	B	1139	GLN
1	B	1213	GLN
1	B	1229	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 10 ligands modelled in this entry, 2 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	POP	A	2003[B]	2	6,8,8	1.86	1 (16%)	12,13,13	1.67	3 (25%)
4	POP	B	2004[A]	2	6,8,8	1.84	1 (16%)	12,13,13	1.63	3 (25%)
4	POP	A	2003[A]	2	6,8,8	1.83	1 (16%)	12,13,13	1.66	3 (25%)
4	POP	B	2004[B]	2	6,8,8	1.89	2 (33%)	12,13,13	1.66	3 (25%)
5	TZP	A	2005	-	12,13,13	1.49	2 (16%)	15,18,18	1.65	3 (20%)
3	ICP	A	2001	-	4,9,9	4.08	2 (50%)	4,12,12	3.86	2 (50%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ICP	B	2002	-	4,9,9	4.08	2 (50%)	4,12,12	3.86	2 (50%)
5	TZP	B	2006	-	12,13,13	1.50	2 (16%)	15,18,18	1.65	4 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	POP	A	2003[B]	2	-	0/6/6/6	-
4	POP	B	2004[A]	2	-	1/6/6/6	-
4	POP	A	2003[A]	2	-	1/6/6/6	-
4	POP	B	2004[B]	2	-	0/6/6/6	-
5	TZP	A	2005	-	-	0/7/7/7	0/1/1/1
3	ICP	A	2001	-	-	-	0/1/1/1
3	ICP	B	2002	-	-	-	0/1/1/1
5	TZP	B	2006	-	-	0/7/7/7	0/1/1/1

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	2001	ICP	C7A-C5A	7.47	1.50	1.33
3	B	2002	ICP	C7A-C5A	7.46	1.50	1.33
5	B	2006	TZP	C4-N3	-3.16	1.33	1.38
5	A	2005	TZP	C4-N3	-3.13	1.33	1.38
3	A	2001	ICP	C6A-N1A	2.76	1.37	1.32
3	B	2002	ICP	C6A-N1A	2.73	1.37	1.32
5	B	2006	TZP	C2-S1	-2.62	1.67	1.72
4	A	2003[B]	POP	P1-O1	2.57	1.58	1.50
4	B	2004[B]	POP	P1-O1	2.53	1.58	1.50
5	A	2005	TZP	C2-S1	-2.52	1.67	1.72
4	B	2004[A]	POP	P1-O1	2.50	1.58	1.50
4	A	2003[A]	POP	P1-O1	2.46	1.58	1.50
4	B	2004[B]	POP	P1-O2	-2.00	1.47	1.54

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	2001	ICP	C5A-C6A-N1A	-6.08	121.83	126.20
3	B	2002	ICP	C5A-C6A-N1A	-6.05	121.85	126.20
3	B	2002	ICP	N1A-C2A-N3A	-4.31	123.04	126.92
3	A	2001	ICP	N1A-C2A-N3A	-4.25	123.09	126.92

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	2005	TZP	S1-C2-N3	-3.52	113.06	116.08
5	B	2006	TZP	S1-C2-N3	-3.42	113.14	116.08
5	A	2005	TZP	C2-N3-C4	3.33	114.46	110.71
5	B	2006	TZP	C2-N3-C4	3.28	114.40	110.71
4	B	2004[B]	POP	O3-P1-O2	2.99	119.00	107.80
4	A	2003[B]	POP	O3-P1-O2	2.97	118.95	107.80
4	A	2003[A]	POP	O3-P1-O2	2.93	118.78	107.80
4	B	2004[A]	POP	O3-P1-O2	2.87	118.56	107.80
4	A	2003[A]	POP	O3-P1-O	2.84	114.16	104.64
4	A	2003[B]	POP	O3-P1-O	2.80	114.04	104.64
4	B	2004[B]	POP	O3-P1-O	2.78	113.95	104.64
4	B	2004[A]	POP	O3-P1-O	2.69	113.67	104.64
4	A	2003[B]	POP	O3-P1-O1	-2.30	101.87	110.83
4	B	2004[B]	POP	O3-P1-O1	-2.30	101.88	110.83
4	A	2003[A]	POP	O3-P1-O1	-2.26	102.02	110.83
4	B	2004[A]	POP	O3-P1-O1	-2.25	102.07	110.83
5	B	2006	TZP	C7-C6-C5	2.20	119.64	112.73
5	A	2005	TZP	C7-C6-C5	2.17	119.54	112.73
5	B	2006	TZP	O1-P1-O2	2.00	118.64	110.83

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	2004[A]	POP	P1-O-P2-O4
4	A	2003[A]	POP	P1-O-P2-O4

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	2004[B]	POP	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	225/225 (100%)	0.76	28 (12%)	8 8	7, 14, 25, 35	4 (1%)
1	B	225/225 (100%)	0.77	25 (11%)	10 10	7, 14, 25, 54	6 (2%)
All	All	450/450 (100%)	0.76	53 (11%)	9 9	7, 14, 25, 54	10 (2%)

All (53) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	63	GLY	6.3
1	B	1235	ARG	5.6
1	B	1234	GLY	4.7
1	A	160	LYS	4.3
1	A	64	ASP	4.0
1	B	1163	ARG	3.8
1	B	1192	ASP	3.7
1	A	12	ILE	3.6
1	B	1084	GLU	3.6
1	A	15	THR	3.5
1	B	1063	GLY	3.5
1	A	161	ASP	3.5
1	B	1161	ASP	3.4
1	A	117	GLU	3.4
1	B	1216	ASP	3.3
1	B	1062	GLY	3.2
1	A	11	GLY	3.2
1	B	1064	ASP	3.1
1	A	192	ASP	3.1
1	B	1231	TYR	3.1
1	A	235	ARG	2.9
1	A	234	GLY	2.9
1	B	1157	GLU	2.9
1	B	1164	ALA	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	116	LYS	2.8
1	A	163	ARG	2.7
1	A	191	ILE	2.7
1	B	1011	GLY	2.6
1	B	1233	THR	2.6
1	A	132	HIS	2.6
1	B	1019	ARG	2.6
1	B	1114	ASN	2.4
1	B	1215	GLU	2.4
1	B	1229	GLN	2.4
1	A	84	GLU	2.4
1	A	229	GLN	2.4
1	A	176	ARG	2.3
1	A	157	GLU	2.3
1	B	1142	GLU	2.2
1	A	21	MET	2.2
1	A	179	ILE	2.2
1	B	1096	GLU	2.2
1	A	69	GLU	2.2
1	B	1050	LYS	2.2
1	B	1179	ILE	2.1
1	A	162	THR	2.1
1	A	226	GLU	2.1
1	A	214	ALA	2.1
1	B	1162	THR	2.1
1	B	1111	GLU	2.0
1	A	178	GLY	2.0
1	A	111	GLU	2.0
1	A	65	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	ICP	B	2002	9/9	0.88	0.10	13,14,17,17	0
3	ICP	A	2001	9/9	0.92	0.10	10,14,17,20	0
4	POP	B	2004[A]	9/9	0.94	0.09	5,12,16,18	4
4	POP	B	2004[B]	9/9	0.94	0.09	5,12,16,17	4
4	POP	A	2003[A]	9/9	0.95	0.09	9,12,14,16	4
4	POP	A	2003[B]	9/9	0.95	0.09	9,12,14,16	4
5	TZP	A	2005	13/13	0.97	0.07	8,10,18,18	0
5	TZP	B	2006	13/13	0.97	0.08	10,11,16,19	0
2	MG	A	2007	1/1	0.99	0.04	11,11,11,11	0
2	MG	B	2008	1/1	0.99	0.04	12,12,12,12	0

6.5 Other polymers

There are no such residues in this entry.