



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 10:55 PM UTC

PDB ID : 3G6G / pdb_00003g6g
Title : Equally potent inhibition of c-Src and Abl by compounds that recognize inactive kinase conformations
Authors : Seeliger, M.A.; Ranjitkar, P.; Kasap, C.; Shan, Y.; Shaw, D.E.; Shah, N.P.; Kuriyan, J.; Maly, D.J.
Deposited on : 2009-02-06
Resolution : 2.31 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.31 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7754 (2.34-2.30)
Clashscore	190562	8383 (2.34-2.30)
Ramachandran outliers	187476	8303 (2.34-2.30)
Sidechain outliers	187428	8303 (2.34-2.30)
RSRZ outliers	180081	7760 (2.34-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	286	12% 78% 13% 8%
1	B	286	15% 73% 15% 8%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4394 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Proto-oncogene tyrosine-protein kinase Src.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	264	Total	C	N	O	S	0	0	0
			2118	1359	354	388	17			
1	B	263	Total	C	N	O	S	0	0	0
			2114	1357	353	387	17			

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	248	GLY	-	expression tag	UNP P00523
A	249	HIS	-	expression tag	UNP P00523
A	250	MET	-	expression tag	UNP P00523
B	248	GLY	-	expression tag	UNP P00523
B	249	HIS	-	expression tag	UNP P00523
B	250	MET	-	expression tag	UNP P00523

- Molecule 2 is N-{3-[(3-{4-[(4-methoxyphenyl)amino]-1,3,5-triazin-2-yl}pyridin-2-yl)amino]-4-methylphenyl}-4-[(4-methylpiperazin-1-yl)methyl]benzamide (CCD ID: G6G) (formula: C₃₅H₃₇N₉O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			46	35	9	2		
2	B	1	Total	C	N	O	0	0
			46	35	9	2		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	40	Total 40	O 40	0	0
4	B	18	Total 18	O 18	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Proto-oncogene tyrosine-protein kinase Src



- Molecule 1: Proto-oncogene tyrosine-protein kinase Src



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	41.88Å 63.85Å 76.62Å 102.50° 89.40° 90.29°	Depositor
Resolution (Å)	43.48 – 2.31 43.48 – 2.31	Depositor EDS
% Data completeness (in resolution range)	95.1 (43.48-2.31) 95.2 (43.48-2.31)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.19 (at 2.32Å)	Xtriage
Refinement program	REFMAC 5.5.0035	Depositor
R, R_{free}	0.229 , 0.277 0.228 , 0.272	Depositor DCC
R_{free} test set	1652 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	42.0	Xtriage
Anisotropy	0.095	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 27.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.023 for h,-k,-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4394	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.25% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: G6G, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.88	1/2170 (0.0%)	0.98	0/2936
1	B	0.86	0/2166	1.00	4/2931 (0.1%)
All	All	0.87	1/4336 (0.0%)	0.99	4/5867 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	422	ALA	CA-C	5.32	1.55	1.52

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	356	LYS	N-CA-C	-8.28	102.97	113.23
1	B	424	PHE	N-CA-C	8.26	122.62	109.40
1	B	468	ASN	N-CA-C	6.43	118.83	111.11
1	B	519	TYR	N-CA-C	6.39	118.78	111.11

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2118	0	2104	26	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	2114	0	2101	35	0
2	A	46	0	37	1	0
2	B	46	0	37	5	0
3	A	6	0	8	0	0
3	B	6	0	8	0	0
4	A	40	0	0	0	0
4	B	18	0	0	0	0
All	All	4394	0	4295	65	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:426:ILE:HD11	1:B:472:LEU:HG	1.42	0.98
1:A:269:LEU:HD22	1:A:294:ILE:HD13	1.62	0.82
1:B:526:GLN:OE1	1:B:526:GLN:N	2.12	0.79
1:A:426:ILE:HD11	1:A:472:LEU:HD11	1.63	0.79
1:B:476:GLU:HA	1:B:476:GLU:OE1	1.89	0.73
1:A:502:ASP:HB3	1:A:505:GLU:HG3	1.71	0.73
1:B:354:MET:O	1:B:356:LYS:N	2.21	0.72
1:B:426:ILE:CD1	1:B:472:LEU:HG	2.18	0.72
1:B:502:ASP:HB3	1:B:505:GLU:HG3	1.72	0.71
1:B:272:LYS:HZ1	1:B:275:GLN:HG3	1.60	0.66
1:A:384:HIS:O	1:A:385:ARG:HB2	1.99	0.62
1:B:334:ILE:O	1:B:335:TYR:HD1	1.83	0.60
1:B:272:LYS:NZ	1:B:275:GLN:HG3	2.15	0.60
1:B:354:MET:CA	1:B:354:MET:HE3	2.32	0.60
1:A:424:PHE:HD1	1:A:425:PRO:HD2	1.67	0.60
1:B:258:ASP:O	1:B:261:GLU:HB2	2.01	0.60
1:B:383:VAL:HG12	1:B:385:ARG:HG3	1.83	0.59
2:B:2:G6G:H5	2:B:2:G6G:N11	2.17	0.59
1:A:354:MET:HE3	1:A:354:MET:CA	2.34	0.57
1:A:426:ILE:HG21	1:A:468:ASN:CG	2.31	0.55
2:A:1:G6G:N11	2:A:1:G6G:H5	2.22	0.55
2:B:2:G6G:H24	2:B:2:G6G:O33	2.06	0.55
1:A:347:LEU:O	1:A:351:LYS:HG2	2.07	0.54
1:B:256:ALA:O	1:B:257:LYS:HG3	2.08	0.54
1:A:270:GLU:OE1	1:A:285:THR:OG1	2.26	0.54
1:A:426:ILE:HD11	1:A:472:LEU:CD1	2.35	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:354:MET:C	1:B:356:LYS:H	2.19	0.50
1:B:283:MET:HG3	1:B:340:TYR:CE1	2.47	0.50
1:A:354:MET:O	1:A:355:GLY:C	2.55	0.48
1:B:349:PHE:CZ	1:B:354:MET:HG3	2.49	0.47
1:A:424:PHE:CD1	1:A:425:PRO:HD2	2.49	0.47
1:A:264:ARG:NH2	1:A:331:GLU:O	2.47	0.47
1:B:354:MET:HE3	1:B:354:MET:C	2.40	0.47
1:B:453:THR:HG22	1:B:495:MET:SD	2.54	0.47
1:A:258:ASP:HB3	1:A:260:TRP:H	1.79	0.46
1:B:269:LEU:HD22	1:B:294:ILE:HD13	1.98	0.45
2:B:2:G6G:N11	2:B:2:G6G:C5	2.79	0.45
1:A:354:MET:HE3	1:A:354:MET:HA	1.98	0.45
1:B:483:CYS:SG	1:B:487:CYS:O	2.73	0.45
1:A:263:PRO:O	1:A:266:SER:OG	2.34	0.45
1:A:287:ASN:C	1:A:289:THR:H	2.25	0.45
1:B:297:LEU:HD11	1:B:302:MET:HB3	1.99	0.44
1:A:384:HIS:O	1:A:385:ARG:CB	2.66	0.44
1:B:258:ASP:OD1	1:B:258:ASP:C	2.60	0.44
1:B:396:GLU:O	1:B:397:ASN:HB3	2.17	0.44
1:A:361:PRO:HA	1:A:520:PHE:CE1	2.52	0.43
1:B:524:GLU:OE2	1:B:526:GLN:NE2	2.51	0.43
1:A:458:LYS:HD3	1:A:458:LYS:HA	1.75	0.43
1:B:361:PRO:HA	1:B:520:PHE:CE1	2.54	0.43
1:B:384:HIS:O	1:B:385:ARG:HB2	2.19	0.43
1:B:503:PRO:HA	1:B:506:ARG:CZ	2.49	0.43
1:B:354:MET:HE3	1:B:354:MET:HA	1.99	0.42
1:A:354:MET:HE3	1:A:354:MET:O	2.18	0.42
1:B:458:LYS:HA	1:B:458:LYS:HD3	1.79	0.42
1:A:476:GLU:OE1	1:A:476:GLU:HA	2.20	0.42
1:A:278:PHE:CD2	1:A:302:MET:HB2	2.55	0.42
1:B:424:PHE:HA	1:B:425:PRO:HD3	1.87	0.41
1:A:260:TRP:NE1	1:A:315:LYS:HG2	2.35	0.41
1:B:385:ARG:HD3	2:B:2:G6G:H46A	2.02	0.41
1:B:281:VAL:HG21	2:B:2:G6G:C19	2.51	0.41
1:A:428:TRP:HE1	1:A:454:GLU:CD	2.29	0.41
1:A:283:MET:HG2	1:A:284:GLY:N	2.35	0.40
1:B:258:ASP:OD1	1:B:259:ALA:N	2.54	0.40
1:B:347:LEU:O	1:B:351:LYS:HG3	2.21	0.40
1:B:354:MET:HA	1:B:354:MET:CE	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	260/286 (91%)	254 (98%)	6 (2%)	0	100	100
1	B	259/286 (91%)	242 (93%)	14 (5%)	3 (1%)	10	11
All	All	519/572 (91%)	496 (96%)	20 (4%)	3 (1%)	21	25

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	354	MET
1	B	301	THR
1	B	355	GLY

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	227/245 (93%)	218 (96%)	9 (4%)	28	41
1	B	227/245 (93%)	216 (95%)	11 (5%)	23	33
All	All	454/490 (93%)	434 (96%)	20 (4%)	25	37

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	265	GLU
1	A	269	LEU
1	A	285	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	296	THR
1	A	351	LYS
1	A	360	LEU
1	A	398	LEU
1	A	424	PHE
1	A	426	ILE
1	B	258	ASP
1	B	269	LEU
1	B	275	GLN
1	B	285	THR
1	B	328	VAL
1	B	351	LYS
1	B	360	LEU
1	B	426	ILE
1	B	438	ARG
1	B	451	LEU
1	B	476	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	381	ASN
1	B	275	GLN
1	B	397	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	G6G	A	1	-	51,51,51	1.21	5 (9%)	69,70,70	2.97	13 (18%)
3	GOL	A	1003	-	5,5,5	0.33	0	5,5,5	0.54	0
2	G6G	B	2	-	51,51,51	1.23	5 (9%)	69,70,70	2.92	13 (18%)
3	GOL	B	1002	-	5,5,5	0.36	0	5,5,5	0.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	G6G	A	1	-	-	0/26/36/36	0/6/6/6
3	GOL	A	1003	-	-	1/4/4/4	-
2	G6G	B	2	-	-	2/26/36/36	0/6/6/6
3	GOL	B	1002	-	-	3/4/4/4	-

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1	G6G	C16-C12	-3.76	1.40	1.48
2	B	2	G6G	C16-C12	-3.74	1.40	1.48
2	A	1	G6G	C25-N30	-3.46	1.34	1.41
2	B	2	G6G	C25-N30	-3.15	1.35	1.41
2	B	2	G6G	C6-N9	-2.67	1.35	1.40
2	B	2	G6G	O7-C8	2.49	1.49	1.42
2	A	1	G6G	C29-C28	2.40	1.55	1.51
2	A	1	G6G	C6-N9	-2.20	1.36	1.40
2	B	2	G6G	C32-C31	2.12	1.54	1.50
2	A	1	G6G	O7-C8	2.10	1.48	1.42

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1	G6G	C14-N13-C12	11.57	120.28	114.14
2	B	2	G6G	C14-N15-C10	11.37	120.54	113.27
2	A	1	G6G	C10-N11-C12	11.32	121.72	114.50
2	B	2	G6G	C10-N11-C12	10.35	121.10	114.50
2	B	2	G6G	C14-N13-C12	9.73	119.30	114.14
2	A	1	G6G	C14-N15-C10	9.63	119.42	113.27
2	B	2	G6G	N15-C10-N11	-8.63	118.08	126.42
2	A	1	G6G	N15-C10-N11	-7.63	119.04	126.42
2	A	1	G6G	N13-C12-N11	-7.28	118.09	125.23
2	B	2	G6G	N13-C12-N11	-6.13	119.22	125.23
2	A	1	G6G	C16-C12-N13	5.31	123.45	117.22
2	A	1	G6G	N15-C14-N13	-4.72	121.44	128.58
2	B	2	G6G	N15-C14-N13	-4.52	121.75	128.58
2	B	2	G6G	C16-C12-N13	4.06	121.98	117.22
2	B	2	G6G	C16-C21-N22	3.61	123.75	119.97
2	B	2	G6G	C23-N22-C21	-3.17	119.98	129.28
2	B	2	G6G	C39-N40-C41	3.15	117.85	111.07
2	B	2	G6G	C19-N20-C21	3.14	122.86	116.74
2	A	1	G6G	C19-N20-C21	3.07	122.71	116.74
2	A	1	G6G	C44-N43-C42	2.79	113.99	109.54
2	A	1	G6G	C41-C42-N43	-2.54	106.77	110.86
2	B	2	G6G	C27-C28-C23	2.30	119.65	117.50
2	B	2	G6G	C44-N43-C42	2.25	113.12	109.54
2	A	1	G6G	C18-C19-N20	-2.08	120.12	123.42
2	A	1	G6G	C8-O7-C3	-2.02	113.17	117.50
2	A	1	G6G	C27-C28-C23	2.01	119.38	117.50

There are no chirality outliers.

All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	B	1002	GOL	O1-C1-C2-C3
3	A	1003	GOL	O2-C2-C3-O3
3	B	1002	GOL	O1-C1-C2-O2
3	B	1002	GOL	O2-C2-C3-O3
2	B	2	G6G	C4-C3-O7-C8
2	B	2	G6G	C2-C3-O7-C8


There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1	G6G	1	0
2	B	2	G6G	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	264/286 (92%)	0.82	35 (13%) 7 8	25, 42, 74, 83	0
1	B	263/286 (91%)	0.93	42 (15%) 5 5	27, 42, 83, 91	0
All	All	527/572 (92%)	0.87	77 (14%) 6 6	25, 42, 78, 91	0

All (77) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	422	ALA	7.4
1	B	424	PHE	6.6
1	A	422	ALA	6.1
1	B	423	LYS	6.0
1	B	421	GLY	5.1
1	B	256	ALA	4.9
1	B	278	PHE	4.7
1	A	423	LYS	4.6
1	B	277	CYS	4.0
1	B	301	THR	3.9
1	A	424	PHE	3.8
1	A	259	ALA	3.8
1	A	355	GLY	3.7
1	A	278	PHE	3.6
1	B	269	LEU	3.6
1	A	256	ALA	3.4
1	B	271	VAL	3.3
1	B	425	PRO	3.2
1	B	282	TRP	3.2
1	B	297	LEU	3.2
1	A	421	GLY	3.1
1	A	277	CYS	3.1
1	B	279	GLY	3.1
1	B	273	LEU	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	262	ILE	3.1
1	A	257	LYS	3.0
1	A	469	ARG	3.0
1	B	354	MET	2.9
1	B	284	GLY	2.9
1	B	267	LEU	2.9
1	B	307	PHE	2.9
1	A	258	ASP	2.8
1	B	355	GLY	2.8
1	A	353	GLU	2.8
1	A	299	PRO	2.7
1	A	405	PHE	2.7
1	B	299	PRO	2.7
1	A	269	LEU	2.7
1	B	304	PRO	2.7
1	B	264	ARG	2.7
1	A	273	LEU	2.7
1	B	333	PRO	2.6
1	A	301	THR	2.5
1	A	526	GLN	2.5
1	A	276	GLY	2.5
1	A	475	VAL	2.4
1	A	302	MET	2.4
1	B	274	GLY	2.4
1	B	328	VAL	2.4
1	B	286	TRP	2.4
1	B	276	GLY	2.4
1	B	306	ALA	2.4
1	A	304	PRO	2.4
1	A	426	ILE	2.4
1	B	281	VAL	2.3
1	A	265	GLU	2.3
1	B	288	GLY	2.3
1	B	526	GLN	2.3
1	A	332	GLU	2.3
1	B	296	THR	2.3
1	A	275	GLN	2.2
1	A	267	LEU	2.2
1	A	288	GLY	2.2
1	A	300	GLY	2.2
1	B	335	TYR	2.2
1	A	268	ARG	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	334	ILE	2.2
1	B	291	ARG	2.1
1	B	436	TYR	2.1
1	B	268	ARG	2.1
1	B	353	GLU	2.1
1	B	295	LYS	2.1
1	A	271	VAL	2.1
1	A	260	TRP	2.1
1	B	305	GLU	2.0
1	B	311	ALA	2.0
1	A	425	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	GOL	B	1002	6/6	0.87	0.17	54,55,55,56	0
3	GOL	A	1003	6/6	0.88	0.19	55,56,57,58	0
2	G6G	B	2	46/46	0.90	0.10	34,40,45,46	0
2	G6G	A	1	46/46	0.93	0.09	29,37,41,45	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around G6G B 2:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around G6G A 1:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.