



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 03:14 PM UTC

PDB ID : 4G65 / pdb_00004g65
Title : Potassium transporter peripheral membrane component (trkA) from *Vibrio vulnificus*
Authors : Filippova, E.V.; Wawrzak, Z.; Skarina, T.; Onopriyenko, O.; Edwards, A.; Savchenko, A.; Anderson, W.F.; Center for Structural Genomics of Infectious Diseases (CSGID)
Deposited on : 2012-07-18
Resolution : 2.09 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	461	 3% 89% 9% .
1	B	461	 3% 89% 10% ..

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 7583 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Trk system potassium uptake protein trkA.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	459	Total	C	N	O	S	Se	0	8	0
			3583	2246	633	690	2	12			
1	B	458	Total	C	N	O	S	Se	0	2	0
			3533	2218	622	680	2	11			

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	expression tag	UNP Q8DDE6
A	-1	ASN	-	expression tag	UNP Q8DDE6
A	0	ALA	-	expression tag	UNP Q8DDE6
B	-2	SER	-	expression tag	UNP Q8DDE6
B	-1	ASN	-	expression tag	UNP Q8DDE6
B	0	ALA	-	expression tag	UNP Q8DDE6

- Molecule 2 is TETRAETHYLENE GLYCOL (CCD ID: PG4) (formula: C₈H₁₈O₅).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	7	Total	Cl	0	0
			7	7		
4	B	7	Total	Cl	0	0
			7	7		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	234	Total	O	0	6
			240	240		
5	B	192	Total	O	0	4
			196	196		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Trk system potassium uptake protein trkA



- Molecule 1: Trk system potassium uptake protein trkA



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	68.11Å 225.26Å 139.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.67 – 2.09 29.67 – 2.09	Depositor EDS
% Data completeness (in resolution range)	97.5 (29.67-2.09) 97.5 (29.67-2.09)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.75 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.179 , 0.228 0.186 , 0.231	Depositor DCC
R_{free} test set	3155 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	23.6	Xtriage
Anisotropy	0.029	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 40.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7583	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, PG4, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.04	2/3621 (0.1%)	1.03	7/4890 (0.1%)
1	B	0.96	1/3571 (0.0%)	0.96	3/4823 (0.1%)
All	All	1.00	3/7192 (0.0%)	1.00	10/9713 (0.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	217	HIS	CG-ND1	-6.81	1.30	1.38
1	A	135	ARG	C-O	-5.41	1.17	1.24
1	B	178	MSE	SE-CE	-5.14	1.80	1.95

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	238	VAL	CB-CA-C	6.28	119.69	110.77
1	A	238	VAL	N-CA-CB	-6.22	103.55	111.64
1	B	332	ARG	N-CA-C	5.54	119.08	110.17
1	A	333	GLY	N-CA-C	5.47	119.51	112.83
1	B	212	VAL	CB-CA-C	-5.41	103.66	111.34
1	A	178	MSE	CB-CA-C	-5.40	104.14	111.21
1	B	444	VAL	CB-CA-C	-5.28	108.75	114.35
1	A	196	ARG	CA-C-N	-5.20	113.83	120.23
1	A	196	ARG	C-N-CA	-5.20	113.83	120.23
1	A	108	LYS	CB-CA-C	-5.10	102.32	110.79

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3583	0	3619	37	0
1	B	3533	0	3573	29	0
2	A	7	0	9	4	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
4	A	7	0	0	0	0
4	B	7	0	0	1	0
5	A	240	0	0	3	0
5	B	196	0	0	0	0
All	All	7583	0	7201	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:437:PHE:C	1:A:438:LEU:HD23	1.82	1.02
1:B:80:MSE:CE	1:B:117:ILE:HD13	2.06	0.86
1:A:438:LEU:HD23	1:A:438:LEU:N	1.96	0.80
1:A:313[A]:MSE:HE3	1:A:313[A]:MSE:HA	1.67	0.76
1:A:156[B]:VAL:HG11	1:A:222:MSE:SE	2.36	0.76
1:B:80:MSE:HE2	1:B:117:ILE:HD13	1.68	0.75
1:A:332:ARG:NH1	5:A:817:HOH:O	2.22	0.72
1:A:438:LEU:HD11	1:A:447:VAL:HG21	1.73	0.68
1:A:46:ARG:HH21	1:A:46:ARG:HG3	1.60	0.66
1:B:45:LEU:HD12	1:B:45:LEU:C	2.21	0.66
1:B:408:THR:HG23	1:B:438:LEU:HD22	1.80	0.63
1:B:333:GLY:HA3	1:B:352:GLN:HG3	1.81	0.62
1:A:413:ILE:HD12	1:A:426:THR:HG21	1.83	0.61
1:B:384:VAL:HG22	1:B:433:HIS:CE1	2.36	0.60
1:B:333:GLY:CA	1:B:352:GLN:HG3	2.32	0.59
1:A:332:ARG:H	1:A:332:ARG:HD2	1.68	0.58
1:A:438:LEU:N	1:A:438:LEU:CD2	2.68	0.56
1:B:220:SER:O	1:B:224:GLU:HG2	2.05	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:103:GLN:NE2	5:A:780:HOH:O	2.40	0.55
1:A:438:LEU:HD11	1:A:447:VAL:CG2	2.37	0.54
1:B:212:VAL:HG11	1:B:412:ALA:HB2	1.89	0.54
1:B:152:LYS:HE3	1:B:416:GLY:O	2.08	0.54
1:A:332:ARG:HG2	1:A:335:TYR:CD1	2.44	0.53
1:A:335:TYR:O	1:A:339:VAL:HG23	2.08	0.53
1:A:266:ARG:HG3	1:A:266:ARG:HH11	1.74	0.53
1:A:332:ARG:CG	1:A:335:TYR:CD1	2.95	0.50
1:A:332:ARG:HD2	1:A:332:ARG:N	2.27	0.49
1:A:126:GLU:OE2	5:A:786:HOH:O	2.20	0.49
1:B:291:THR:HG23	1:B:321:MSE:HE2	1.93	0.49
1:B:45:LEU:HD12	1:B:45:LEU:O	2.13	0.49
1:B:291:THR:HG23	1:B:321:MSE:CE	2.43	0.49
1:B:384:VAL:HG22	1:B:433:HIS:ND1	2.28	0.49
1:B:309:GLU:OE1	1:B:313:MSE:HE1	2.12	0.49
1:A:205:ALA:O	1:A:206:ASP:HB2	2.13	0.49
1:A:309:GLU:H	2:A:501:PG4:H41	1.77	0.49
1:A:332:ARG:HG2	1:A:335:TYR:CG	2.48	0.49
1:A:331:GLN:NE2	1:A:331:GLN:HA	2.29	0.48
1:B:333:GLY:HA3	1:B:352:GLN:CG	2.43	0.48
1:A:309:GLU:N	2:A:501:PG4:H41	2.28	0.48
2:A:501:PG4:H32	2:A:501:PG4:O1	2.13	0.48
1:B:352:GLN:HG2	4:B:505:CL:CL	2.50	0.47
1:B:453:PRO:HB2	1:B:455:PRO:HD2	1.97	0.47
1:A:397:ARG:HB2	1:A:402:ILE:HD11	1.95	0.47
1:B:363:ARG:NH2	1:B:384:VAL:HG21	2.30	0.46
1:B:452:GLN:HG3	1:B:456:PHE:CD2	2.50	0.46
1:B:238:VAL:HG11	1:B:284:ALA:HB1	1.97	0.46
1:A:103:GLN:HG2	1:B:56:ASP:OD1	2.17	0.44
1:B:174:LEU:HD22	1:B:178:MSE:HE2	1.99	0.44
1:B:178:MSE:HE1	1:B:225:LEU:HD23	2.00	0.44
1:A:437:PHE:C	1:A:438:LEU:CD2	2.72	0.44
1:B:45:LEU:C	1:B:45:LEU:CD1	2.90	0.44
1:A:45:LEU:C	1:A:45:LEU:HD12	2.43	0.43
1:A:17:GLU:O	1:A:20:VAL:HG22	2.18	0.43
2:A:501:PG4:O1	2:A:501:PG4:C3	2.65	0.43
1:B:178:MSE:HE1	1:B:225:LEU:CD2	2.49	0.43
1:A:235[A]:ILE:HD13	1:A:251:LEU:HD13	2.01	0.42
1:A:261:GLU:O	1:A:281:CYS:HA	2.19	0.42
1:A:408:THR:HG23	1:A:438:LEU:HD22	2.02	0.42
1:A:313[A]:MSE:HE3	1:A:313[A]:MSE:CA	2.42	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:329:LEU:HD11	1:B:350:PRO:HD3	2.01	0.42
1:B:101[B]:SER:OG	1:B:103:GLN:OE1	2.35	0.41
1:A:196:ARG:HG3	1:A:407:GLY:HA2	2.03	0.41
1:A:121:HIS:CE1	1:A:364:ARG:HH22	2.39	0.41
1:A:331:GLN:O	1:A:351:GLN:OE1	2.39	0.41
1:A:330:ILE:O	1:A:349:SER:HB2	2.21	0.41
1:A:332:ARG:H	1:A:332:ARG:CD	2.32	0.41
1:B:349:SER:HB3	1:B:352:GLN:HB2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	465/461 (101%)	454 (98%)	11 (2%)	0	100	100
1	B	456/461 (99%)	449 (98%)	7 (2%)	0	100	100
All	All	921/922 (100%)	903 (98%)	18 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	386/369 (105%)	372 (96%)	14 (4%)	31	34

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	379/369 (103%)	364 (96%)	15 (4%)	28	29
All	All	765/738 (104%)	736 (96%)	29 (4%)	30	32

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	19	LEU
1	A	38	GLU
1	A	41	ASP
1	A	46	ARG
1	A	138[A]	GLN
1	A	138[B]	GLN
1	A	177	HIS
1	A	193	ARG
1	A	196	ARG
1	A	238	VAL
1	A	331	GLN
1	A	332	ARG
1	A	346	VAL
1	A	438	LEU
1	B	19	LEU
1	B	41	ASP
1	B	181	ILE
1	B	195	ILE
1	B	254	THR
1	B	262	ARG
1	B	269	LYS
1	B	272	GLU
1	B	291	THR
1	B	297	GLN
1	B	307	GLU
1	B	393	LYS
1	B	413	ILE
1	B	438	LEU
1	B	452	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	180	HIS
1	A	216	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	276	ASN
1	A	352	GLN
1	A	423	HIS
1	A	430	GLN
1	B	64	GLN
1	B	276	ASN
1	B	352	GLN
1	B	423	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 17 ligands modelled in this entry, 14 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PG4	A	501	-	6,6,12	0.43	0	5,5,11	0.31	0
3	SO4	B	501	-	4,4,4	0.40	0	6,6,6	0.55	0
3	SO4	A	502	-	4,4,4	0.38	0	6,6,6	0.76	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PG4	A	501	-	-	3/4/4/10	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	PG4	C1-C2-O2-C3
2	A	501	PG4	O1-C1-C2-O2
2	A	501	PG4	O2-C3-C4-O3

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	PG4	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	448/461 (97%)	-0.06	15 (3%)	49	51	6, 23, 47, 70	7 (1%)
1	B	447/461 (96%)	0.00	13 (2%)	53	56	10, 29, 50, 79	2 (0%)
All	All	895/922 (97%)	-0.03	28 (3%)	51	54	6, 26, 49, 79	9 (1%)

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	162[A]	TYR	5.6
1	A	457	PHE	4.9
1	B	455	PRO	4.0
1	B	179	PRO	4.0
1	A	335	TYR	3.5
1	B	456	PHE	3.5
1	A	162	TYR	3.5
1	A	338	LEU	3.1
1	A	177	HIS	2.9
1	A	332	ARG	2.8
1	A	331	GLN	2.6
1	A	423	HIS	2.6
1	A	425	ARG	2.5
1	B	177	HIS	2.5
1	B	192	GLY	2.5
1	A	333	GLY	2.4
1	A	334	ALA	2.3
1	B	334	ALA	2.3
1	B	193	ARG	2.3
1	A	438	LEU	2.3
1	B	337	ASP	2.3
1	B	283	ASP	2.2
1	B	341	GLY	2.2
1	B	-1	ASN	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	174	LEU	2.1
1	A	193	ARG	2.1
1	B	174	LEU	2.1
1	A	336	VAL	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PG4	A	501	7/13	0.68	0.17	35,46,49,50	0
4	CL	A	505	1/1	0.82	0.16	48,48,48,48	0
4	CL	B	508	1/1	0.89	0.14	59,59,59,59	0
4	CL	A	506	1/1	0.90	0.11	51,51,51,51	0
4	CL	B	505	1/1	0.91	0.10	49,49,49,49	0
4	CL	B	506	1/1	0.93	0.08	47,47,47,47	0
4	CL	A	509	1/1	0.93	0.08	46,46,46,46	0
4	CL	B	504	1/1	0.94	0.08	29,29,29,29	1
4	CL	A	504	1/1	0.94	0.09	29,29,29,29	0
4	CL	A	507	1/1	0.96	0.08	30,30,30,30	0
3	SO4	A	502	5/5	0.97	0.10	27,28,32,35	0
4	CL	B	502	1/1	0.97	0.06	33,33,33,33	0
3	SO4	B	501	5/5	0.97	0.07	27,30,32,36	0
4	CL	A	503	1/1	0.98	0.04	36,36,36,36	0
4	CL	B	507	1/1	0.98	0.04	34,34,34,34	0
4	CL	A	508	1/1	0.98	0.08	28,28,28,28	0
4	CL	B	503	1/1	0.99	0.04	15,15,15,15	0

6.5 Other polymers [i](#)

There are no such residues in this entry.