



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 1, 2026 – 04:19 PM UTC

PDB ID : 4G7H / pdb_00004g7h
Title : Crystal structure of Thermus thermophilus transcription initiation complex
Authors : Zhang, Y.; Ebright, R.H.
Deposited on : 2012-07-20
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2481 (2.90-2.90)
Clashscore	190562	2690 (2.90-2.90)
Ramachandran outliers	187476	2623 (2.90-2.90)
Sidechain outliers	187428	2625 (2.90-2.90)
RSRZ outliers	180081	2481 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	315	% 53% 14% • 28%
1	B	315	3% 54% 13% • 30%
1	K	315	53% 16% • 28%
1	L	315	% 56% 14% • 29%
2	C	1119	% 76% 21% ••

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	M	1119	% 75% 21% • •
3	D	1524	% 77% 19% • •
3	N	1524	% 77% 19% • •
4	E	99	% 79% 16% 5%
4	O	99	2% 75% 19% • 5%
5	F	443	% 62% 15% • 22%
5	P	443	3% 57% 16% 5% • 22%
6	G	19	5% 79% 5% 16%
6	Q	19	11% 79% 5% 16%
7	H	27	4% 41% 48% 11%
7	R	27	7% 52% 37% 11%

2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 57420 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-directed RNA polymerase subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	226	Total	C	N	O	S	0	0	0
			1782	1138	310	332	2			
1	B	222	Total	C	N	O	S	0	0	0
			1750	1118	304	326	2			
1	K	226	Total	C	N	O	S	0	0	0
			1782	1138	310	332	2			
1	L	225	Total	C	N	O	S	0	0	0
			1773	1133	308	330	2			

- Molecule 2 is a protein called DNA-directed RNA polymerase subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	1111	Total	C	N	O	S	0	0	0
			8770	5548	1564	1634	24			
2	M	1111	Total	C	N	O	S	0	0	0
			8770	5548	1564	1634	24			

- Molecule 3 is a protein called DNA-directed RNA polymerase subunit beta'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	1486	Total	C	N	O	S	0	0	0
			11738	7441	2067	2195	35			
3	N	1486	Total	C	N	O	S	0	0	0
			11738	7441	2067	2195	35			

- Molecule 4 is a protein called DNA-directed RNA polymerase subunit omega.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	E	94	Total	C	N	O	S	0	0	0
			761	486	132	139	4			
4	O	94	Total	C	N	O	S	0	0	0
			761	486	132	139	4			

- Molecule 5 is a protein called RNA polymerase sigma factor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	F	346	Total	C	N	O	S	0	0	0
			2807	1770	509	524	4			
5	P	347	Total	C	N	O	S	0	0	0
			2814	1774	510	526	4			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
F	-19	MET	-	expression tag	UNP Q5SKW1
F	-18	GLY	-	expression tag	UNP Q5SKW1
F	-17	SER	-	expression tag	UNP Q5SKW1
F	-16	SER	-	expression tag	UNP Q5SKW1
F	-15	HIS	-	expression tag	UNP Q5SKW1
F	-14	HIS	-	expression tag	UNP Q5SKW1
F	-13	HIS	-	expression tag	UNP Q5SKW1
F	-12	HIS	-	expression tag	UNP Q5SKW1
F	-11	HIS	-	expression tag	UNP Q5SKW1
F	-10	HIS	-	expression tag	UNP Q5SKW1
F	-9	SER	-	expression tag	UNP Q5SKW1
F	-8	SER	-	expression tag	UNP Q5SKW1
F	-7	GLY	-	expression tag	UNP Q5SKW1
F	-6	LEU	-	expression tag	UNP Q5SKW1
F	-5	VAL	-	expression tag	UNP Q5SKW1
F	-4	PRO	-	expression tag	UNP Q5SKW1
F	-3	ARG	-	expression tag	UNP Q5SKW1
F	-2	GLY	-	expression tag	UNP Q5SKW1
F	-1	SER	-	expression tag	UNP Q5SKW1
F	0	HIS	-	expression tag	UNP Q5SKW1
P	-19	MET	-	expression tag	UNP Q5SKW1
P	-18	GLY	-	expression tag	UNP Q5SKW1
P	-17	SER	-	expression tag	UNP Q5SKW1
P	-16	SER	-	expression tag	UNP Q5SKW1
P	-15	HIS	-	expression tag	UNP Q5SKW1
P	-14	HIS	-	expression tag	UNP Q5SKW1
P	-13	HIS	-	expression tag	UNP Q5SKW1
P	-12	HIS	-	expression tag	UNP Q5SKW1
P	-11	HIS	-	expression tag	UNP Q5SKW1
P	-10	HIS	-	expression tag	UNP Q5SKW1
P	-9	SER	-	expression tag	UNP Q5SKW1
P	-8	SER	-	expression tag	UNP Q5SKW1
P	-7	GLY	-	expression tag	UNP Q5SKW1
P	-6	LEU	-	expression tag	UNP Q5SKW1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
P	-5	VAL	-	expression tag	UNP Q5SKW1
P	-4	PRO	-	expression tag	UNP Q5SKW1
P	-3	ARG	-	expression tag	UNP Q5SKW1
P	-2	GLY	-	expression tag	UNP Q5SKW1
P	-1	SER	-	expression tag	UNP Q5SKW1
P	0	HIS	-	expression tag	UNP Q5SKW1

- Molecule 6 is a DNA chain called 5'-D(*CP*CP*T*GP*CP*AP*TP*CP*CP*GP*TP*GP*AP*GP*TP*CP*GP*AP*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	G	16	Total	C	N	O	P	0	0	0
			328	156	63	94	15			
6	Q	16	Total	C	N	O	P	0	0	0
			328	156	63	94	15			

- Molecule 7 is a DNA chain called 5'-D(*TP*AP*TP*AP*AP*TP*GP*GP*GP*AP*GP*C P*TP*GP*TP*CP*AP*CP*GP*GP*AP*TP*GP*CP*AP*GP*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	H	24	Total	C	N	O	P	0	0	0
			495	236	94	142	23			
7	R	24	Total	C	N	O	P	0	0	0
			495	236	94	142	23			

- Molecule 8 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	B	1	Total	Mg	0	0
			1	1		
8	D	3	Total	Mg	0	0
			3	3		
8	F	1	Total	Mg	0	0
			1	1		
8	K	1	Total	Mg	0	0
			1	1		
8	N	3	Total	Mg	0	0
			3	3		
8	P	1	Total	Mg	0	0
			1	1		

- Molecule 9 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	D	2	Total 2	Zn 2	0	0
9	N	2	Total 2	Zn 2	0	0

- Molecule 10 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	16	Total 16	O 16	0	0
10	B	4	Total 4	O 4	0	0
10	C	104	Total 104	O 104	0	0
10	D	138	Total 138	O 138	0	0
10	E	5	Total 5	O 5	0	0
10	F	32	Total 32	O 32	0	0
10	G	7	Total 7	O 7	0	0
10	H	6	Total 6	O 6	0	0
10	K	12	Total 12	O 12	0	0
10	L	8	Total 8	O 8	0	0
10	M	58	Total 58	O 58	0	0
10	N	89	Total 89	O 89	0	0
10	O	6	Total 6	O 6	0	0
10	P	21	Total 21	O 21	0	0
10	Q	3	Total 3	O 3	0	0
10	R	5	Total 5	O 5	0	0




• Molecule 1: DNA-directed RNA polymerase subunit alpha




• Molecule 2: DNA-directed RNA polymerase subunit beta







• Molecule 2: DNA-directed RNA polymerase subunit beta





• Molecule 3: DNA-directed RNA polymerase subunit beta'






● Molecule 3: DNA-directed RNA polymerase subunit beta'







- Molecule 4: DNA-directed RNA polymerase subunit omega




- Molecule 4: DNA-directed RNA polymerase subunit omega




- Molecule 5: RNA polymerase sigma factor







• Molecule 5: RNA polymerase sigma factor



• Molecule 6: 5'-D(*CP*CP*T*GP*CP*AP*TP*CP*CP*GP*TP*GP*AP*GP*TP*CP*GP*AP*G)-3'



• Molecule 6: 5'-D(*CP*CP*T*GP*CP*AP*TP*CP*CP*GP*TP*GP*AP*GP*TP*CP*GP*AP*G)-3'



• Molecule 7: 5'-D(*TP*AP*TP*AP*AP*TP*GP*GP*GP*AP*GP*CP*TP*GP*TP*CP*AP*C*P*GP*GP*AP*TP*GP*CP*AP*GP*G)-3'





● Molecule 7: 5'-D(*TP*AP*TP*AP*AP*TP*GP*GP*GP*AP*GP*CP*TP*GP*TP*CP*AP*CP*GP*GP*AP*TP*GP*CP*AP*GP*G)-3'



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	185.56Å 104.57Å 297.55Å 90.00° 98.32° 90.00°	Depositor
Resolution (Å)	49.27 – 2.90 49.27 – 2.90	Depositor EDS
% Data completeness (in resolution range)	99.5 (49.27-2.90) 99.6 (49.27-2.90)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.11 (at 2.91Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.3_928)	Depositor
R, R_{free}	0.188 , 0.226 0.195 , 0.227	Depositor DCC
R_{free} test set	2205 reflections (0.88%)	wwPDB-VP
Wilson B-factor (Å ²)	61.5	Xtriage
Anisotropy	0.653	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 42.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	57420	wwPDB-VP
Average B, all atoms (Å ²)	71.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 25.73 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.0020e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.44	0/1814	0.94	12/2466 (0.5%)
1	B	0.39	0/1782	0.90	4/2424 (0.2%)
1	K	0.39	0/1814	0.90	8/2466 (0.3%)
1	L	0.38	0/1805	0.91	9/2454 (0.4%)
2	C	0.43	0/8937	0.85	24/12087 (0.2%)
2	M	0.41	0/8937	0.84	14/12087 (0.1%)
3	D	0.41	0/11944	0.82	13/16149 (0.1%)
3	N	0.41	0/11944	0.82	9/16149 (0.1%)
4	E	0.34	0/775	0.76	0/1045
4	O	0.36	0/775	0.74	0/1045
5	F	0.39	0/2852	0.82	5/3837 (0.1%)
5	P	0.40	0/2859	0.89	6/3847 (0.2%)
6	G	0.24	0/368	0.72	0/567
6	Q	0.19	0/368	0.74	0/567
7	H	0.25	0/556	0.84	0/858
7	R	0.22	0/556	0.82	0/858
All	All	0.40	0/58086	0.84	104/78906 (0.1%)

There are no bond length outliers.

The worst 5 of 104 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	P	417	LYS	N-CA-C	8.85	123.58	111.54
5	P	416	ARG	N-CA-C	8.15	121.68	108.32
1	L	112	ARG	NE-CZ-NH2	7.76	126.19	119.20
2	C	422	ARG	NE-CZ-NH2	7.60	126.04	119.20
1	A	187	GLY	N-CA-C	-7.50	105.47	115.21

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1782	0	1834	40	0
1	B	1750	0	1797	33	0
1	K	1782	0	1834	35	0
1	L	1773	0	1826	27	0
2	C	8770	0	8874	148	0
2	M	8770	0	8874	165	0
3	D	11738	0	11971	163	0
3	N	11738	0	11971	165	0
4	E	761	0	778	11	0
4	O	761	0	778	13	0
5	F	2807	0	2882	41	0
5	P	2814	0	2889	85	0
6	G	328	0	181	1	0
6	Q	328	0	181	1	0
7	H	495	0	272	12	0
7	R	495	0	272	10	0
8	B	1	0	0	0	0
8	D	3	0	0	0	0
8	F	1	0	0	0	0
8	K	1	0	0	0	0
8	N	3	0	0	0	0
8	P	1	0	0	0	0
9	D	2	0	0	0	0
9	N	2	0	0	0	0
10	A	16	0	0	2	0
10	B	4	0	0	0	0
10	C	104	0	0	8	0
10	D	138	0	0	5	0
10	E	5	0	0	0	0
10	F	32	0	0	3	0
10	G	7	0	0	0	0
10	H	6	0	0	1	0
10	K	12	0	0	0	0
10	L	8	0	0	0	0
10	M	58	0	0	2	0
10	N	89	0	0	3	0
10	O	6	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
10	P	21	0	0	0	0
10	Q	3	0	0	0	0
10	R	5	0	0	0	0
All	All	57420	0	57214	864	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 864 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:P:415:THR:HG22	5:P:416:ARG:CG	1.50	1.41
5:P:415:THR:CG2	5:P:416:ARG:HG3	1.51	1.40
2:M:172:ILE:HG13	2:M:186:VAL:HG22	1.19	1.14
5:P:415:THR:HG22	5:P:416:ARG:CD	1.89	1.01
2:C:627:ARG:NH1	2:C:638:ASP:OD2	1.92	1.01

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	224/315 (71%)	222 (99%)	2 (1%)	0	100	100
1	B	220/315 (70%)	215 (98%)	5 (2%)	0	100	100
1	K	10/315 (3%)	10 (100%)	0	0	100	100
1	L	223/315 (71%)	218 (98%)	5 (2%)	0	100	100
2	C	1107/1119 (99%)	1086 (98%)	21 (2%)	0	100	100
2	M	1107/1119 (99%)	1080 (98%)	27 (2%)	0	100	100
3	D	1482/1524 (97%)	1455 (98%)	26 (2%)	1 (0%)	48	77

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	N	1482/1524 (97%)	1454 (98%)	27 (2%)	1 (0%)	48	77
4	E	92/99 (93%)	89 (97%)	2 (2%)	1 (1%)	11	36
4	O	92/99 (93%)	90 (98%)	2 (2%)	0	100	100
5	F	344/443 (78%)	339 (98%)	5 (2%)	0	100	100
5	P	345/443 (78%)	341 (99%)	4 (1%)	0	100	100
All	All	6728/7630 (88%)	6599 (98%)	126 (2%)	3 (0%)	100	100

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	E	94	PRO
3	D	530	VAL
3	N	530	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	199/273 (73%)	183 (92%)	16 (8%)	11	34
1	B	195/273 (71%)	181 (93%)	14 (7%)	13	39
1	K	199/273 (73%)	186 (94%)	13 (6%)	15	44
1	L	198/273 (72%)	186 (94%)	12 (6%)	17	46
2	C	936/941 (100%)	858 (92%)	78 (8%)	10	32
2	M	936/941 (100%)	864 (92%)	72 (8%)	12	35
3	D	1253/1279 (98%)	1152 (92%)	101 (8%)	11	33
3	N	1253/1279 (98%)	1153 (92%)	100 (8%)	11	34
4	E	83/88 (94%)	82 (99%)	1 (1%)	63	86
4	O	83/88 (94%)	81 (98%)	2 (2%)	43	75
5	F	301/388 (78%)	285 (95%)	16 (5%)	20	52
5	P	302/388 (78%)	264 (87%)	38 (13%)	4	14

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	5938/6484 (92%)	5475 (92%)	463 (8%)	11	35

5 of 463 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	K	142	VAL
5	P	361	LEU
2	M	480	THR
5	P	310	ILE
3	N	1062	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 28 such sidechains are listed below:

Mol	Chain	Res	Type
5	F	381	HIS
5	P	90	GLN
2	M	565	GLN
3	N	1172	HIS
2	M	187	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 14 ligands modelled in this entry, 14 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	226/315 (71%)	-0.07	2 (0%) 81 75	45, 63, 96, 118	0
1	B	222/315 (70%)	0.21	8 (3%) 46 38	47, 77, 114, 137	0
1	K	226/315 (71%)	-0.04	1 (0%) 88 85	49, 66, 97, 116	0
1	L	225/315 (71%)	0.25	3 (1%) 75 67	51, 81, 118, 139	0
2	C	1111/1119 (99%)	-0.13	6 (0%) 87 83	32, 59, 111, 149	0
2	M	1111/1119 (99%)	0.01	14 (1%) 75 67	34, 65, 129, 152	0
3	D	1486/1524 (97%)	-0.12	11 (0%) 84 79	29, 61, 120, 177	1 (0%)
3	N	1486/1524 (97%)	-0.11	10 (0%) 84 79	32, 64, 118, 178	1 (0%)
4	E	94/99 (94%)	0.10	1 (1%) 78 71	41, 64, 110, 121	0
4	O	94/99 (94%)	-0.10	2 (2%) 63 54	45, 67, 112, 122	0
5	F	346/443 (78%)	-0.08	3 (0%) 81 75	37, 63, 108, 132	0
5	P	347/443 (78%)	0.09	13 (3%) 45 37	42, 72, 139, 172	0
6	G	16/19 (84%)	0.10	1 (6%) 26 20	56, 84, 176, 182	0
6	Q	16/19 (84%)	0.31	2 (12%) 8 7	68, 93, 180, 182	0
7	H	24/27 (88%)	0.25	1 (4%) 40 32	54, 98, 145, 192	0
7	R	24/27 (88%)	0.22	2 (8%) 17 14	55, 112, 147, 199	0
All	All	7054/7722 (91%)	-0.05	80 (1%) 78 71	29, 65, 121, 199	2 (0%)

The worst 5 of 80 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	D	1252	ILE	5.1
3	N	1252	ILE	4.6
2	C	363	SER	4.1
5	P	77	THR	3.6
1	B	190	THR	3.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
8	MG	N	2004	1/1	0.81	0.32	66,66,66,66	0
8	MG	D	2004	1/1	0.83	0.34	66,66,66,66	0
8	MG	P	2001	1/1	0.84	0.18	71,71,71,71	0
8	MG	B	2001	1/1	0.85	0.13	71,71,71,71	0
8	MG	K	1001	1/1	0.85	0.29	67,67,67,67	0
8	MG	N	2005	1/1	0.89	0.11	67,67,67,67	0
8	MG	F	2001	1/1	0.91	0.07	63,63,63,63	0
8	MG	D	2003	1/1	0.93	0.15	32,32,32,32	0
8	MG	N	2003	1/1	0.94	0.15	38,38,38,38	0
8	MG	D	2005	1/1	0.94	0.14	71,71,71,71	0
9	ZN	D	2002	1/1	0.99	0.03	73,73,73,73	0
9	ZN	N	2002	1/1	0.99	0.04	95,95,95,95	0
9	ZN	N	2001	1/1	1.00	0.03	54,54,54,54	0
9	ZN	D	2001	1/1	1.00	0.10	77,77,77,77	0

6.5 Other polymers [i](#)

There are no such residues in this entry.