



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 10:12 PM UTC

PDB ID : 4G7Z / pdb_00004g7z
Title : Crystal structure of Thermus thermophilus transcription initiation complex containing 5-BrU at template-strand position +1
Authors : Zhang, Y.; Ebright, R.H.
Deposited on : 2012-07-20
Resolution : 3.81 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.81 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1043 (3.98-3.66)
Clashscore	190562	1075 (3.98-3.66)
Ramachandran outliers	187476	1029 (3.98-3.66)
Sidechain outliers	187428	1024 (3.98-3.66)
RSRZ outliers	180081	1042 (3.98-3.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	315	40% 27% 5% 28%
1	B	315	41% 26% • 30%
1	K	315	54% 16% • 28%
1	L	315	52% 18% • 29%
2	C	1119	57% 36% 6% •

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	M	1119	
3	D	1524	
3	N	1524	
4	E	99	
4	O	99	
5	F	443	
5	P	443	
6	H	27	
6	R	27	
7	G	21	
7	Q	21	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
7	BRU	Q	15	-	-	X	-

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 56872 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-directed RNA polymerase subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	226	Total	C	N	O	S	0	0	0
			1782	1138	310	332	2			
1	B	222	Total	C	N	O	S	0	0	0
			1750	1118	304	326	2			
1	K	226	Total	C	N	O	S	0	0	0
			1782	1138	310	332	2			
1	L	225	Total	C	N	O	S	0	0	0
			1773	1133	308	330	2			

- Molecule 2 is a protein called DNA-directed RNA polymerase subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	1111	Total	C	N	O	S	0	0	0
			8770	5548	1564	1634	24			
2	M	1111	Total	C	N	O	S	0	0	0
			8770	5548	1564	1634	24			

- Molecule 3 is a protein called DNA-directed RNA polymerase subunit beta'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	1482	Total	C	N	O	S	0	0	0
			11704	7421	2059	2189	35			
3	N	1489	Total	C	N	O	S	0	0	0
			11746	7446	2066	2198	36			

- Molecule 4 is a protein called DNA-directed RNA polymerase subunit omega.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	E	94	Total	C	N	O	S	0	0	0
			761	486	132	139	4			
4	O	94	Total	C	N	O	S	0	0	0
			761	486	132	139	4			

- Molecule 5 is a protein called RNA polymerase sigma factor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	F	346	Total	C	N	O	S	0	0	0
			2807	1770	509	524	4			
5	P	347	Total	C	N	O	S	0	0	0
			2814	1774	510	526	4			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
F	-19	MET	-	expression tag	UNP Q5SKW1
F	-18	GLY	-	expression tag	UNP Q5SKW1
F	-17	SER	-	expression tag	UNP Q5SKW1
F	-16	SER	-	expression tag	UNP Q5SKW1
F	-15	HIS	-	expression tag	UNP Q5SKW1
F	-14	HIS	-	expression tag	UNP Q5SKW1
F	-13	HIS	-	expression tag	UNP Q5SKW1
F	-12	HIS	-	expression tag	UNP Q5SKW1
F	-11	HIS	-	expression tag	UNP Q5SKW1
F	-10	HIS	-	expression tag	UNP Q5SKW1
F	-9	SER	-	expression tag	UNP Q5SKW1
F	-8	SER	-	expression tag	UNP Q5SKW1
F	-7	GLY	-	expression tag	UNP Q5SKW1
F	-6	LEU	-	expression tag	UNP Q5SKW1
F	-5	VAL	-	expression tag	UNP Q5SKW1
F	-4	PRO	-	expression tag	UNP Q5SKW1
F	-3	ARG	-	expression tag	UNP Q5SKW1
F	-2	GLY	-	expression tag	UNP Q5SKW1
F	-1	SER	-	expression tag	UNP Q5SKW1
F	0	HIS	-	expression tag	UNP Q5SKW1
P	-19	MET	-	expression tag	UNP Q5SKW1
P	-18	GLY	-	expression tag	UNP Q5SKW1
P	-17	SER	-	expression tag	UNP Q5SKW1
P	-16	SER	-	expression tag	UNP Q5SKW1
P	-15	HIS	-	expression tag	UNP Q5SKW1
P	-14	HIS	-	expression tag	UNP Q5SKW1
P	-13	HIS	-	expression tag	UNP Q5SKW1
P	-12	HIS	-	expression tag	UNP Q5SKW1
P	-11	HIS	-	expression tag	UNP Q5SKW1
P	-10	HIS	-	expression tag	UNP Q5SKW1
P	-9	SER	-	expression tag	UNP Q5SKW1
P	-8	SER	-	expression tag	UNP Q5SKW1
P	-7	GLY	-	expression tag	UNP Q5SKW1
P	-6	LEU	-	expression tag	UNP Q5SKW1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
P	-5	VAL	-	expression tag	UNP Q5SKW1
P	-4	PRO	-	expression tag	UNP Q5SKW1
P	-3	ARG	-	expression tag	UNP Q5SKW1
P	-2	GLY	-	expression tag	UNP Q5SKW1
P	-1	SER	-	expression tag	UNP Q5SKW1
P	0	HIS	-	expression tag	UNP Q5SKW1

- Molecule 6 is a DNA chain called 5'-D(*TP*AP*TP*AP*AP*TP*GP*GP*GP*AP*GP*C
P*TP*GP*TP*CP*AP*CP*GP*GP*AP*TP*GP*CP*AP*GP*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	H	24	Total	C	N	O	P	0	0	0
			495	236	94	142	23			
6	R	24	Total	C	N	O	P	0	0	0
			495	236	94	142	23			

- Molecule 7 is a DNA chain called 5'-D(*CP*CP*TP*GP*CP*AP*TP*CP*CP*GP*TP*G
P*AP*GP*(BRU)P*CP*GP*AP*GP*GP*G)-3'.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	G	16	Total	Br	C	N	O	P	0	0
			328	1	155	63	94	15		
7	Q	16	Total	Br	C	N	O	P	0	0
			328	1	155	63	94	15		

- Molecule 8 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	D	2	Total	Zn	0	0
			2	2		
8	N	2	Total	Zn	0	0
			2	2		


- Molecule 9 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	D	1	Total	Mg	0	0
			1	1		
9	N	1	Total	Mg	0	0
			1	1		



• Molecule 2: DNA-directed RNA polymerase subunit beta


Chain M: 69% 28% ..






• Molecule 3: DNA-directed RNA polymerase subunit beta'


Chain D: 55% 36% 6% .






• Molecule 3: DNA-directed RNA polymerase subunit beta'


Chain N: 70% 25%






- Molecule 4: DNA-directed RNA polymerase subunit omega

Chain E: 56% 33% 6% 5%




- Molecule 4: DNA-directed RNA polymerase subunit omega


Chain O: 72% 21% 5%




- Molecule 5: RNA polymerase sigma factor

Chain F: 56% 21% 22%







• Molecule 5: RNA polymerase sigma factor




• Molecule 6: 5'-D(*TP*AP*TP*AP*AP*TP*GP*GP*GP*AP*GP*CP*TP*GP*TP*CP*AP*C
P*GP*GP*AP*TP*GP*CP*AP*GP*G)-3'



• Molecule 6: 5'-D(*TP*AP*TP*AP*AP*TP*GP*GP*GP*AP*GP*CP*TP*GP*TP*CP*AP*C
P*GP*GP*AP*TP*GP*CP*AP*GP*G)-3'



• Molecule 7: 5'-D(*CP*CP*TP*GP*CP*AP*TP*CP*CP*GP*TP*GP*AP*GP*(BRU)P*CP*
GP*AP*GP*GP*G)-3'



• Molecule 7: 5'-D(*CP*CP*TP*GP*CP*AP*TP*CP*CP*GP*TP*GP*AP*GP*(BRU)P*CP*
GP*AP*GP*GP*G)-3'



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	185.47Å 103.59Å 296.51Å 90.00° 98.53° 90.00°	Depositor
Resolution (Å)	49.85 – 3.81 49.85 – 3.81	Depositor EDS
% Data completeness (in resolution range)	51.6 (49.85-3.81) 81.7 (49.85-3.81)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	0.17	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.44 (at 3.77Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.3_928)	Depositor
R, R_{free}	0.176 , 0.232 0.193 , 0.251	Depositor DCC
R_{free} test set	1776 reflections (1.63%)	wwPDB-VP
Wilson B-factor (Å ²)	120.0	Xtriage
Anisotropy	0.137	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 107.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	56872	wwPDB-VP
Average B, all atoms (Å ²)	151.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 40.98 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.5249e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, BRU, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.61	0/1814	1.10	14/2466 (0.6%)
1	B	0.60	0/1782	1.11	11/2424 (0.5%)
1	K	0.34	0/1814	0.84	4/2466 (0.2%)
1	L	0.34	0/1805	0.82	5/2454 (0.2%)
2	C	0.62	0/8937	1.08	40/12087 (0.3%)
2	M	0.35	0/8937	0.85	18/12087 (0.1%)
3	D	0.63	0/11910	1.08	59/16105 (0.4%)
3	N	0.34	0/11952	0.83	19/16162 (0.1%)
4	E	0.57	0/775	1.03	5/1045 (0.5%)
4	O	0.31	0/775	0.78	0/1045
5	F	0.35	0/2852	0.80	4/3837 (0.1%)
5	P	0.36	0/2859	0.79	2/3847 (0.1%)
6	H	0.37	0/556	1.06	0/858
6	R	0.41	0/556	1.04	1/858 (0.1%)
7	G	0.36	0/345	0.81	0/529
7	Q	0.34	0/345	0.75	0/529
All	All	0.49	0/58014	0.95	182/78799 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	M	0	1

There are no bond length outliers.

The worst 5 of 182 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	8	ALA	CA-C-N	10.36	130.46	120.31
1	A	8	ALA	C-N-CA	10.36	130.46	120.31
2	C	398	THR	N-CA-C	-9.73	99.97	112.23
1	B	48	ILE	CA-C-N	8.93	128.87	119.85
1	B	48	ILE	C-N-CA	8.93	128.87	119.85

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	M	294	GLU	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1782	0	1834	71	0
1	B	1750	0	1797	67	0
1	K	1782	0	1834	32	0
1	L	1773	0	1826	37	0
2	C	8770	0	8874	320	0
2	M	8770	0	8874	217	0
3	D	11704	0	11934	408	0
3	N	11746	0	11974	284	0
4	E	761	0	778	32	0
4	O	761	0	778	20	0
5	F	2807	0	2882	74	0
5	P	2814	0	2889	123	0
6	H	495	0	272	18	0
6	R	495	0	272	16	0
7	G	328	0	178	7	0
7	Q	328	0	178	13	0
8	D	2	0	0	0	0
8	N	2	0	0	0	0
9	D	1	0	0	0	0
9	N	1	0	0	0	0
All	All	56872	0	57174	1572	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 1572 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:P:408:LEU:O	5:P:412:GLU:CB	1.72	1.37
5:P:408:LEU:O	5:P:412:GLU:HB2	1.07	1.23
5:P:408:LEU:HD23	5:P:412:GLU:CG	1.69	1.21
7:Q:14:DG:H2''	7:Q:15:BRU:O5'	1.43	1.15
3:N:1238:MET:SD	3:N:1359:GLN:OE1	2.07	1.12

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	224/315 (71%)	219 (98%)	3 (1%)	2 (1%)	14	46
1	B	220/315 (70%)	209 (95%)	11 (5%)	0	100	100
1	K	224/315 (71%)	220 (98%)	4 (2%)	0	100	100
1	L	223/315 (71%)	214 (96%)	9 (4%)	0	100	100
2	C	1107/1119 (99%)	1066 (96%)	40 (4%)	1 (0%)	48	79
2	M	1107/1119 (99%)	1065 (96%)	40 (4%)	2 (0%)	43	73
3	D	1478/1524 (97%)	1418 (96%)	59 (4%)	1 (0%)	48	79
3	N	1485/1524 (97%)	1423 (96%)	59 (4%)	3 (0%)	43	73
4	E	92/99 (93%)	88 (96%)	4 (4%)	0	100	100
4	O	92/99 (93%)	88 (96%)	4 (4%)	0	100	100
5	F	344/443 (78%)	337 (98%)	7 (2%)	0	100	100
5	P	345/443 (78%)	331 (96%)	12 (4%)	2 (1%)	21	55
All	All	6941/7630 (91%)	6678 (96%)	252 (4%)	11 (0%)	43	73

5 of 11 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	C	325	ILE
2	M	295	ASP
2	M	325	ILE
5	P	413	SER
3	N	1234	THR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	199/273 (73%)	181 (91%)	18 (9%)	9	32
1	B	195/273 (71%)	179 (92%)	16 (8%)	10	34
1	K	199/273 (73%)	187 (94%)	12 (6%)	17	43
1	L	198/273 (72%)	191 (96%)	7 (4%)	32	54
2	C	936/941 (100%)	846 (90%)	90 (10%)	8	29
2	M	936/941 (100%)	900 (96%)	36 (4%)	29	53
3	D	1250/1279 (98%)	1118 (89%)	132 (11%)	6	26
3	N	1253/1279 (98%)	1187 (95%)	66 (5%)	20	45
4	E	83/88 (94%)	74 (89%)	9 (11%)	6	25
4	O	83/88 (94%)	81 (98%)	2 (2%)	43	62
5	F	301/388 (78%)	293 (97%)	8 (3%)	39	59
5	P	302/388 (78%)	281 (93%)	21 (7%)	14	40
All	All	5935/6484 (92%)	5518 (93%)	417 (7%)	14	40

5 of 417 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	D	1237	THR
1	L	34	VAL
5	P	156	VAL
3	D	1291	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
4	E	68	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 32 such sidechains are listed below:

Mol	Chain	Res	Type
3	N	1116	ASN
3	N	1404	ASN
2	C	899	GLN
2	C	704	HIS
3	N	1442	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	BRU	Q	15	7	18,21,22	2.77	8 (44%)	25,30,33	2.14	5 (20%)
7	BRU	G	15	7	18,21,22	2.78	8 (44%)	25,30,33	2.17	5 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	BRU	Q	15	7	-	0/7/21/22	0/2/2/2
7	BRU	G	15	7	-	2/7/21/22	0/2/2/2

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	G	15	BRU	O4-C4	6.81	1.36	1.23
7	Q	15	BRU	O4-C4	6.79	1.36	1.23
7	G	15	BRU	C6-N1	5.38	1.47	1.38
7	Q	15	BRU	C6-N1	5.36	1.47	1.38
7	Q	15	BRU	C2-N1	3.52	1.44	1.38

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	G	15	BRU	C5-C4-N3	5.83	120.05	113.34
7	Q	15	BRU	C5-C4-N3	5.75	119.95	113.34
7	G	15	BRU	C4-N3-C2	-5.64	119.94	127.34
7	Q	15	BRU	C4-N3-C2	-5.58	120.02	127.34
7	G	15	BRU	O4-C4-C5	-4.59	119.95	125.80

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	G	15	BRU	C3'-C4'-C5'-O5'
7	G	15	BRU	O4'-C4'-C5'-O5'

There are no ring outliers.

2 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	Q	15	BRU	10	0
7	G	15	BRU	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2			OWAB(Å ²)	Q < 0.9
1	A	226/315 (71%)	-0.85	0	100	100	112, 138, 163, 174	0
1	B	222/315 (70%)	-0.78	0	100	100	110, 145, 190, 215	0
1	K	226/315 (71%)	-0.74	0	100	100	114, 145, 171, 193	0
1	L	225/315 (71%)	-0.69	0	100	100	114, 150, 198, 222	0
2	C	1111/1119 (99%)	-0.66	0	100	100	89, 136, 212, 240	0
2	M	1111/1119 (99%)	-0.48	4 (0%)	88	75	98, 154, 239, 269	0
3	D	1482/1524 (97%)	-0.73	1 (0%)	92	87	86, 136, 196, 258	1 (0%)
3	N	1489/1524 (97%)	-0.63	1 (0%)	92	87	84, 139, 199, 263	1 (0%)
4	E	94/99 (94%)	-0.60	0	100	100	107, 138, 175, 185	0
4	O	94/99 (94%)	-0.61	0	100	100	109, 141, 191, 201	0
5	F	346/443 (78%)	-0.53	0	100	100	112, 152, 238, 251	0
5	P	347/443 (78%)	-0.41	3 (0%)	81	61	117, 161, 253, 290	0
6	H	24/27 (88%)	-0.22	0	100	100	144, 191, 241, 247	0
6	R	24/27 (88%)	-0.32	0	100	100	142, 204, 241, 254	0
7	G	15/21 (71%)	-0.32	0	100	100	143, 181, 229, 233	0
7	Q	15/21 (71%)	-0.26	0	100	100	154, 197, 234, 235	0
All	All	7051/7726 (91%)	-0.63	9 (0%)	92	87	84, 144, 219, 290	2 (0%)

The worst 5 of 9 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	M	196	LEU	4.8
3	N	353	VAL	3.1
3	D	302	GLN	2.6
2	M	307	LEU	2.2
5	P	357	ALA	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	BRU	Q	15	20/21	0.88	0.08	177,196,247,303	1
7	BRU	G	15	20/21	0.94	0.07	138,145,192,257	1

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
9	MG	D	2003	1/1	0.96	0.04	111,111,111,111	0
9	MG	N	2003	1/1	0.98	0.04	127,127,127,127	0
8	ZN	D	2002	1/1	0.99	0.02	164,164,164,164	0
8	ZN	N	2001	1/1	0.99	0.03	166,166,166,166	0
8	ZN	D	2001	1/1	1.00	0.02	168,168,168,168	0
8	ZN	N	2002	1/1	1.00	0.03	138,138,138,138	0

6.5 Other polymers [i](#)

There are no such residues in this entry.