



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 12:45 PM UTC

PDB ID : 1G8L / pdb_00001g8l
Title : CRYSTAL STRUCTURE OF ESCHERICHIA COLI MOEA
Authors : Xiang, S.; Nichols, J.; Rajagopalan, K.V.; Schindelin, H.
Deposited on : 2000-11-17
Resolution : 1.95 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	411	
1	B	411	

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 6781 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MOLYBDOPTERIN BIOSYNTHESIS MOEA PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	403	Total	C	N	O	S	0	0	0
			3040	1918	531	578	13			
1	B	403	Total	C	N	O	S	0	0	0
			3040	1918	531	578	13			

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total 6	C 3	O 3	0	0
2	A	1	Total 6	C 3	O 3	0	0
2	A	1	Total 6	C 3	O 3	0	0
2	B	1	Total 6	C 3	O 3	0	0
2	B	1	Total 6	C 3	O 3	0	0
2	B	1	Total 6	C 3	O 3	0	0
2	B	1	Total 6	C 3	O 3	0	0
2	B	1	Total 6	C 3	O 3	0	0
2	B	1	Total 6	C 3	O 3	0	0
2	B	1	Total 6	C 3	O 3	0	0
2	B	1	Total 6	C 3	O 3	0	0
2	B	1	Total 6	C 3	O 3	0	0
2	B	1	Total 6	C 3	O 3	0	0
2	B	1	Total 6	C 3	O 3	0	0
2	B	1	Total 6	C 3	O 3	0	0

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	312	Total 312	O 312	0	0
3	B	281	Total 281	O 281	0	0

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: MOLYBDOPTERIN BIOSYNTHESIS MOEA PROTEIN



• Molecule 1: MOLYBDOPTERIN BIOSYNTHESIS MOEA PROTEIN



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	88.32Å 97.41Å 98.74Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.95	Depositor
% Data completeness (in resolution range)	91.5 (20.00-1.95)	Depositor
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
Refinement program	X-PLOR	Depositor
R, R_{free}	0.228 , 0.284	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	6781	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.90	0/3099	0.92	4/4212 (0.1%)
1	B	0.89	0/3099	0.92	7/4212 (0.2%)
All	All	0.90	0/6198	0.92	11/8424 (0.1%)

There are no bond length outliers.

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	336	ALA	N-CA-C	-7.67	103.49	112.92
1	B	336	ALA	N-CA-C	-7.47	103.90	113.16
1	B	89	GLU	N-CA-C	7.01	118.57	111.07
1	B	21	VAL	N-CA-C	6.93	118.64	109.21
1	B	255	VAL	N-CA-C	-6.30	105.21	111.88

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3040	0	3037	128	0
1	B	3040	0	3037	130	0
2	A	42	0	56	9	0
2	B	66	0	88	9	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	312	0	0	12	0
3	B	281	0	0	15	0
All	All	6781	0	6218	255	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

The worst 5 of 255 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:356:ASN:HD21	1:A:360:GLU:HB2	1.09	1.17
1:B:368:HIS:HD2	1:B:370:GLY:H	1.00	0.95
1:B:374:PHE:H	2:B:703:GOL:H31	1.30	0.94
1:A:177:LYS:HA	2:A:706:GOL:H11	1.52	0.92
1:A:356:ASN:ND2	1:A:360:GLU:HB2	1.83	0.92

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	401/411 (98%)	383 (96%)	13 (3%)	5 (1%)	10	4
1	B	401/411 (98%)	378 (94%)	18 (4%)	5 (1%)	10	4
All	All	802/822 (98%)	761 (95%)	31 (4%)	10 (1%)	10	4

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	198	ASP
1	A	282	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	282	LYS
1	A	82	ALA
1	A	228	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	325/331 (98%)	276 (85%)	49 (15%)	3	0
1	B	325/331 (98%)	277 (85%)	48 (15%)	3	0
All	All	650/662 (98%)	553 (85%)	97 (15%)	3	0

5 of 97 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	79	LYS
1	B	157	THR
1	B	104	VAL
1	B	119	GLN
1	B	196	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 29 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	368	HIS
1	B	354	GLN
1	B	15	ASN
1	B	222	ASN
1	A	392	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

18 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	B	710	-	5,5,5	0.35	0	5,5,5	0.48	0
2	GOL	B	712	-	5,5,5	0.33	0	5,5,5	0.55	0
2	GOL	A	714	-	5,5,5	0.40	0	5,5,5	0.26	0
2	GOL	B	718	-	5,5,5	0.39	0	5,5,5	0.40	0
2	GOL	B	711	-	5,5,5	0.29	0	5,5,5	0.41	0
2	GOL	B	703	-	5,5,5	0.36	0	5,5,5	0.47	0
2	GOL	A	704	-	5,5,5	0.44	0	5,5,5	0.52	0
2	GOL	B	705	-	5,5,5	0.38	0	5,5,5	0.24	0
2	GOL	A	709	-	5,5,5	0.38	0	5,5,5	0.35	0
2	GOL	A	717	-	5,5,5	0.39	0	5,5,5	0.33	0
2	GOL	B	716	-	5,5,5	0.41	0	5,5,5	0.24	0
2	GOL	B	702	-	5,5,5	0.49	0	5,5,5	0.78	0
2	GOL	B	707	-	5,5,5	0.32	0	5,5,5	0.32	0
2	GOL	B	708	-	5,5,5	0.38	0	5,5,5	0.49	0
2	GOL	A	701	-	5,5,5	0.60	0	5,5,5	0.79	0
2	GOL	B	713	-	5,5,5	0.35	0	5,5,5	0.46	0
2	GOL	A	715	-	5,5,5	0.43	0	5,5,5	0.44	0
2	GOL	A	706	-	5,5,5	0.57	0	5,5,5	0.99	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	B	710	-	-	2/4/4/4	-
2	GOL	B	712	-	-	2/4/4/4	-
2	GOL	A	714	-	-	2/4/4/4	-
2	GOL	B	718	-	-	0/4/4/4	-
2	GOL	B	711	-	-	2/4/4/4	-
2	GOL	B	703	-	-	0/4/4/4	-
2	GOL	A	704	-	-	2/4/4/4	-
2	GOL	B	705	-	-	2/4/4/4	-
2	GOL	A	709	-	-	2/4/4/4	-
2	GOL	A	717	-	-	2/4/4/4	-
2	GOL	B	716	-	-	2/4/4/4	-
2	GOL	B	702	-	-	0/4/4/4	-
2	GOL	B	707	-	-	2/4/4/4	-
2	GOL	B	708	-	-	2/4/4/4	-
2	GOL	A	701	-	-	4/4/4/4	-
2	GOL	B	713	-	-	4/4/4/4	-
2	GOL	A	715	-	-	0/4/4/4	-
2	GOL	A	706	-	-	4/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 34 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	706	GOL	O1-C1-C2-C3
2	A	709	GOL	O1-C1-C2-C3
2	A	714	GOL	O1-C1-C2-C3
2	B	707	GOL	O1-C1-C2-C3
2	B	708	GOL	C1-C2-C3-O3

There are no ring outliers.

10 monomers are involved in 18 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	711	GOL	2	0
2	B	703	GOL	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	704	GOL	1	0
2	A	709	GOL	1	0
2	B	702	GOL	2	0
2	B	707	GOL	2	0
2	B	708	GOL	1	0
2	A	701	GOL	3	0
2	A	715	GOL	1	0
2	A	706	GOL	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.