



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 28, 2026 – 09:03 PM UTC

PDB ID : 1G8Q / pdb_00001g8q
Title : CRYSTAL STRUCTURE OF HUMAN CD81 EXTRACELLULAR DOMAIN, A RECEPTOR FOR HEPATITIS C VIRUS
Authors : Kitadokoro, K.; Bolognesi, M.; Bordo, D.; Grandi, G.; Galli, G.; Petracca, R.; Falugi, F.
Deposited on : 2000-11-20
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	4931 (1.60-1.60)
Ramachandran outliers	187476	4831 (1.60-1.60)
Sidechain outliers	187428	4830 (1.60-1.60)
RSRZ outliers	180081	4672 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	90	% 50% 39% 8% .
1	B	90	3% 47% 33% 14% .

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1550 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CD81 ANTIGEN, EXTRACELLULAR DOMAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	90	Total	C	N	O	S	3	0	0
			693	430	118	141	4			
1	B	86	Total	C	N	O	S	17	0	0
			663	414	113	132	4			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	202	HIS	LEU	conflict	UNP P60033
B	302	HIS	LEU	conflict	UNP P60033


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	104	Total	O	0	0
			104	104		
2	B	90	Total	O	0	0
			90	90		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: CD81 ANTIGEN, EXTRACELLULAR DOMAIN



• Molecule 1: CD81 ANTIGEN, EXTRACELLULAR DOMAIN



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	31.48Å 77.17Å 38.46Å 90.00° 107.39° 90.00°	Depositor
Resolution (Å)	20.00 – 1.60 20.00 – 1.60	Depositor EDS
% Data completeness (in resolution range)	93.0 (20.00-1.60) 87.9 (20.00-1.60)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.12 (at 1.60Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.188 , 0.238 0.188 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	22.3	Xtriage
Anisotropy	0.380	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.51 , 101.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	1550	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.14% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.74	7/702 (1.0%)	2.66	56/948 (5.9%)
1	B	1.77	7/671 (1.0%)	3.42	52/902 (5.8%)
All	All	1.76	14/1373 (1.0%)	3.05	108/1850 (5.8%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	277	SER	C-N	24.00	1.68	1.33
1	A	138	ASP	CB-CG	11.64	1.81	1.52
1	B	280	ASN	N-CA	8.29	1.57	1.46
1	A	147	VAL	CA-CB	6.70	1.63	1.54
1	B	222	ASP	C-N	6.47	1.42	1.33
1	A	146	VAL	N-CA	5.50	1.53	1.46
1	A	124	LYS	N-CA	5.50	1.53	1.46
1	A	165	LEU	C-N	5.38	1.41	1.33
1	B	223	VAL	CA-CB	5.31	1.61	1.54
1	B	213	PHE	CA-CB	5.23	1.64	1.53
1	B	290	CYS	N-CA	5.13	1.52	1.46
1	B	255	ASP	C-N	5.05	1.40	1.33
1	A	143	ALA	C-N	5.02	1.40	1.33
1	A	188	GLU	CA-C	5.01	1.59	1.52

All (108) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	277	SER	O-C-N	-66.90	30.09	122.41
1	B	213	PHE	CA-CB-CG	18.29	132.09	113.80
1	A	138	ASP	CA-CB-CG	-14.99	97.61	112.60
1	A	167	THR	CA-C-O	11.94	133.67	120.90
1	B	292	GLN	OE1-CD-NE2	-11.45	111.15	122.60
1	B	284	ASN	CA-CB-CG	-11.22	101.38	112.60
1	B	280	ASN	N-CA-C	-10.72	98.72	112.23
1	A	141	ASN	O-C-N	9.92	134.47	122.27
1	B	273	ASN	CA-CB-CG	8.93	121.53	112.60
1	B	231	LEU	O-C-N	8.80	131.13	122.07
1	A	192	GLN	OE1-CD-NE2	-8.49	114.11	122.60
1	B	268	SER	CB-CA-C	8.12	124.26	110.79
1	B	302	HIS	CA-CB-CG	8.09	121.89	113.80
1	A	147	VAL	CA-CB-CG2	-8.06	96.70	110.40
1	B	269	VAL	CA-C-O	-7.93	112.08	120.57
1	A	141	ASN	CA-C-O	-7.73	111.08	119.97
1	A	138	ASP	CA-C-O	-7.72	111.09	119.97
1	A	143	ALA	O-C-N	-7.70	113.96	122.12
1	A	129	GLN	OE1-CD-NE2	7.39	129.99	122.60
1	B	287	LYS	CA-C-O	-7.34	112.64	120.42
1	A	173	ASN	OD1-CG-ND2	7.33	129.93	122.60
1	B	284	ASN	OD1-CG-ND2	7.20	129.80	122.60
1	A	145	ALA	O-C-N	7.19	131.11	122.27
1	B	263	THR	N-CA-C	7.16	119.08	111.28
1	A	117	ASP	CA-C-O	-7.09	113.03	120.55
1	B	280	ASN	CA-CB-CG	7.04	119.64	112.60
1	B	255	ASP	CB-CA-C	-6.92	102.00	112.11
1	B	229	GLN	O-C-N	-6.91	114.27	122.15
1	B	218	GLN	N-CA-CB	6.86	119.96	110.01
1	B	225	GLN	CA-C-O	6.86	127.70	120.42
1	A	125	GLN	OE1-CD-NE2	-6.82	115.78	122.60
1	A	188	GLU	O-C-N	6.78	130.85	122.86
1	B	255	ASP	CA-C-O	6.77	130.49	122.03
1	B	290	CYS	CA-C-O	-6.73	112.50	120.10
1	B	268	SER	CA-C-O	6.68	127.64	120.55
1	A	134	ALA	CA-C-O	6.68	127.65	120.10
1	A	139	ASP	CA-C-N	-6.66	111.60	122.26
1	A	139	ASP	C-N-CA	-6.66	111.60	122.26
1	B	301	LYS	CB-CA-C	-6.65	99.27	110.44
1	B	229	GLN	OE1-CD-NE2	6.64	129.24	122.60
1	A	147	VAL	N-CA-CB	-6.60	100.67	110.58
1	A	151	HIS	CB-CG-ND1	-6.52	112.92	122.70
1	A	190	CYS	CA-C-O	-6.50	112.49	119.97

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	189	ASP	OD1-CG-OD2	-6.48	107.35	122.90
1	A	169	VAL	CA-C-O	6.47	127.68	120.95
1	A	126	PHE	CA-C-O	6.46	127.81	120.90
1	A	189	ASP	CB-CG-OD2	6.44	133.22	118.40
1	B	291	HIS	CA-CB-CG	-6.39	107.41	113.80
1	A	141	ASN	OD1-CG-ND2	6.36	128.96	122.60
1	B	224	LYS	CA-C-O	-6.33	112.69	119.97
1	B	225	GLN	CA-C-N	6.30	128.63	120.44
1	B	225	GLN	C-N-CA	6.30	128.63	120.44
1	A	166	THR	O-C-N	-6.29	115.46	122.12
1	A	176	PRO	O-C-N	6.26	131.09	122.64
1	A	145	ALA	CA-C-N	-6.18	112.77	120.56
1	A	145	ALA	C-N-CA	-6.18	112.77	120.56
1	B	214	VAL	O-C-N	-6.16	115.89	122.61
1	A	141	ASN	CA-C-N	-6.13	111.59	120.29
1	A	141	ASN	C-N-CA	-6.13	111.59	120.29
1	B	281	ILE	N-CA-CB	6.04	117.61	110.55
1	B	291	HIS	CB-CG-ND1	-6.01	113.69	122.70
1	A	122	ASP	N-CA-CB	6.01	119.05	110.16
1	A	143	ALA	CA-C-O	5.98	126.89	120.55
1	A	134	ALA	N-CA-C	5.88	118.47	111.71
1	A	173	ASN	CA-CB-CG	-5.87	106.73	112.60
1	B	279	SER	N-CA-C	-5.75	105.52	112.54
1	A	114	VAL	CA-C-O	5.73	127.36	121.23
1	B	216	LYS	CA-C-O	5.73	127.03	120.90
1	A	114	VAL	N-CA-CB	-5.71	101.22	110.13
1	B	299	SER	CA-CB-OG	-5.71	99.68	111.10
1	B	225	GLN	O-C-N	-5.70	115.66	122.15
1	B	233	GLN	N-CA-C	5.69	118.21	111.33
1	A	165	LEU	CD1-CG-CD2	5.66	123.25	110.80
1	B	277	SER	CA-C-N	5.60	132.38	121.41
1	B	277	SER	C-N-CA	5.60	132.38	121.41
1	B	222	ASP	CA-C-O	5.56	126.45	120.55
1	A	115	ASN	O-C-N	5.56	131.10	122.87
1	A	169	VAL	O-C-N	-5.53	116.50	121.87
1	B	251	HIS	ND1-CE1-NE2	5.52	113.92	108.40
1	B	293	LYS	CG-CD-CE	-5.46	98.73	111.30
1	A	159	SER	N-CA-CB	5.46	120.11	111.43
1	A	145	ALA	CA-C-O	-5.45	113.70	119.97
1	B	217	ASP	N-CA-CB	-5.38	102.20	110.16
1	B	269	VAL	O-C-N	5.37	127.29	121.87
1	B	291	HIS	CB-CG-CD2	5.34	138.15	131.20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	242	ASN	CA-CB-CG	-5.34	107.26	112.60
1	A	139	ASP	N-CA-CB	5.28	119.83	111.17
1	A	136	VAL	O-C-N	-5.28	114.13	122.04
1	B	232	GLN	N-CA-C	-5.26	105.23	110.97
1	A	157	CYS	O-C-N	-5.25	116.31	123.24
1	A	159	SER	O-C-N	5.24	130.11	122.94
1	A	202	HIS	CA-CB-CG	-5.23	108.57	113.80
1	A	144	LYS	CA-C-O	-5.21	114.89	120.42
1	A	135	VAL	CA-CB-CG2	5.20	119.24	110.40
1	B	231	LEU	CA-C-O	-5.18	115.38	120.82
1	A	192	GLN	CA-CB-CG	-5.18	103.74	114.10
1	B	229	GLN	CA-C-O	5.18	125.91	120.42
1	A	147	VAL	CA-CB-CG1	5.17	119.19	110.40
1	B	275	CYS	CA-C-N	5.12	125.12	119.90
1	B	275	CYS	C-N-CA	5.12	125.12	119.90
1	B	216	LYS	N-CA-C	5.11	117.24	111.11
1	B	301	LYS	N-CA-CB	-5.08	102.83	111.27
1	A	166	THR	N-CA-C	5.08	117.48	111.33
1	B	255	ASP	CA-CB-CG	-5.07	107.53	112.60
1	A	156	CYS	CA-C-N	-5.07	114.11	121.76
1	A	156	CYS	C-N-CA	-5.07	114.11	121.76
1	A	175	CYS	CA-C-N	5.05	126.16	119.84
1	A	175	CYS	C-N-CA	5.05	126.16	119.84

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	259	SER	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	693	0	672	32	1
1	B	663	0	648	18	8
2	A	104	0	0	18	9

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	90	0	0	7	2
All	All	1550	0	1320	48	10

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:229:GLN:HG2	2:B:194:HOH:O	1.24	1.28
1:A:202:HIS:HA	2:A:306:HOH:O	1.71	0.89
1:B:216:LYS:HE3	2:B:138:HOH:O	1.76	0.84
1:A:118:GLN:HG3	1:A:121:LYS:NZ	1.93	0.83
1:A:155:ASP:OD2	2:A:292:HOH:O	2.03	0.75
1:B:269:VAL:HG12	1:B:274:LEU:HD12	1.71	0.72
1:A:202:HIS:O	2:A:216:HOH:O	2.08	0.71
1:A:149:THR:HA	2:A:279:HOH:O	1.91	0.70
1:A:118:GLN:HG3	1:A:121:LYS:HZ2	1.56	0.69
1:B:236:VAL:HG23	1:B:237:ASP:N	2.05	0.69
1:A:113:PHE:CD1	2:A:250:HOH:O	2.45	0.68
1:A:118:GLN:NE2	2:A:205:HOH:O	2.27	0.68
1:A:113:PHE:N	2:A:276:HOH:O	2.28	0.67
1:A:201:LYS:O	1:A:202:HIS:C	2.38	0.65
1:A:113:PHE:CD1	2:A:271:HOH:O	2.51	0.61
1:A:180:ASN:ND2	1:A:183:SER:H	2.00	0.58
1:B:273:ASN:HA	2:B:22:HOH:O	2.03	0.58
1:B:244:LYS:HG3	2:B:12:HOH:O	2.02	0.58
1:A:118:GLN:O	1:A:121:LYS:HD3	2.03	0.58
1:A:113:PHE:CG	2:A:271:HOH:O	2.57	0.57
1:A:188:GLU:HG3	2:A:245:HOH:O	2.05	0.56
1:A:149:THR:HG23	2:A:279:HOH:O	2.04	0.56
1:A:113:PHE:HA	2:A:286:HOH:O	2.04	0.55
1:B:236:VAL:HG23	1:B:237:ASP:HB2	1.88	0.54
1:A:118:GLN:CG	1:A:121:LYS:NZ	2.70	0.53
1:B:215:ASN:HD22	1:B:218:GLN:H	1.55	0.52
1:A:149:THR:HG23	1:B:302:HIS:HB3	1.92	0.51
1:B:216:LYS:NZ	2:B:150:HOH:O	2.45	0.50
1:B:233:GLN:CD	2:B:194:HOH:O	2.55	0.48
1:B:287:LYS:NZ	2:B:131:HOH:O	2.47	0.48
1:A:121:LYS:HE2	2:A:247:HOH:O	2.14	0.47
1:A:121:LYS:CE	1:A:122:ASP:OD1	2.64	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:236:VAL:O	1:B:237:ASP:O	2.33	0.46
1:B:235:VAL:HG21	1:B:265:LEU:CD2	2.46	0.46
1:A:121:LYS:HG2	2:A:247:HOH:O	2.16	0.45
1:A:149:THR:CG2	1:B:302:HIS:HB3	2.46	0.45
1:A:149:THR:CA	2:A:279:HOH:O	2.59	0.44
1:A:118:GLN:CG	1:A:121:LYS:HZ1	2.30	0.43
1:A:202:HIS:CE1	2:A:284:HOH:O	2.72	0.43
1:B:215:ASN:HD21	1:B:217:ASP:HB2	1.83	0.43
1:A:118:GLN:HG3	1:A:121:LYS:HZ1	1.77	0.42
1:B:235:VAL:HG21	1:B:265:LEU:HD22	2.01	0.42
1:A:149:THR:CB	2:A:279:HOH:O	2.67	0.41
1:A:193:LYS:HB3	1:A:202:HIS:NE2	2.35	0.41
1:B:224:LYS:CE	1:B:295:ASP:OD1	2.68	0.41
1:A:176:PRO:HD2	2:A:252:HOH:O	2.21	0.41
1:A:180:ASN:HD21	1:A:183:SER:H	1.66	0.41
1:A:142:ASN:HD22	1:A:142:ASN:HA	1.79	0.40

All (10) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:213:PHE:CZ	2:A:300:HOH:O[1_655]	0.78	1.42
1:A:138:ASP:OD1	2:B:180:HOH:O[2_646]	1.37	0.83
1:B:213:PHE:CE1	2:A:300:HOH:O[1_655]	1.43	0.77
1:B:213:PHE:CE2	2:A:300:HOH:O[1_655]	1.66	0.54
1:B:282:ILE:C	2:A:285:HOH:O[2_556]	1.83	0.37
1:B:282:ILE:CG2	2:A:285:HOH:O[2_556]	1.87	0.33
1:B:213:PHE:CE2	2:A:223:HOH:O[1_655]	2.15	0.05
2:A:245:HOH:O	2:B:143:HOH:O[1_455]	2.15	0.05
1:B:282:ILE:O	2:A:285:HOH:O[2_556]	2.17	0.03
1:B:282:ILE:CB	2:A:285:HOH:O[2_556]	2.17	0.03

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	88/90 (98%)	85 (97%)	3 (3%)	0	100	100
1	B	82/90 (91%)	78 (95%)	4 (5%)	0	100	100
All	All	170/180 (94%)	163 (96%)	7 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	80/80 (100%)	72 (90%)	8 (10%)	7	1
1	B	76/80 (95%)	72 (95%)	4 (5%)	20	4
All	All	156/160 (98%)	144 (92%)	12 (8%)	12	2

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	118	GLN
1	A	121	LYS
1	A	138	ASP
1	A	139	ASP
1	A	147	VAL
1	A	167	THR
1	A	176	PRO
1	A	202	HIS
1	B	216	LYS
1	B	235	VAL
1	B	236	VAL
1	B	277	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	142	ASN
1	A	180	ASN
1	B	215	ASN
1	B	272	ASN
1	B	302	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	B	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	B	277:SER	C	278:GLY	N	1.68

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	90/90 (100%)	0.27	1 (1%) 78 82	21, 31, 46, 68	1 (1%)
1	B	84/90 (93%)	0.35	3 (3%) 46 48	11, 30, 49, 73	2 (2%)
All	All	174/180 (96%)	0.31	4 (2%) 61 64	11, 31, 48, 73	3 (1%)

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	213	PHE	4.2
1	A	163	THR	2.8
1	B	236	VAL	2.7
1	B	257	CYS	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.