



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 17, 2026 – 08:02 PM UTC

PDB ID : 3G8C / pdb_00003g8c
Title : Crystal Structure of Biotin Carboxylase in Complex with Biotin, Bicarbonate, ADP and Mg Ion
Authors : Chou, C.Y.; Yu, L.P.; Tong, L.
Deposited on : 2009-02-12
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	444	14% 89% 10% .
1	B	444	4% 92% 8% .

2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 7794 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Biotin carboxylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	444	Total	C	N	O	S	0	0	0
			3421	2155	611	633	22			
1	B	444	Total	C	N	O	S	0	0	0
			3421	2155	611	633	22			

- Molecule 2 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
2	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Mg 1 1	0	0
3	B	1	Total Mg 1 1	0	0

- Molecule 4 is BIOTIN (CCD ID: BTN) (formula: $C_{10}H_{16}N_2O_3S$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C N O S 16 10 2 3 1	0	0
4	B	1	Total C N O S 16 10 2 3 1	0	0

- Molecule 5 is BICARBONATE ION (CCD ID: BCT) (formula: CHO_3).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	1	3		
5	B	1	Total	C	O	0	0
			4	1	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	396	Total	O	0	0
			396	396		
6	B	460	Total	O	0	0
			460	460		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Biotin carboxylase



• Molecule 1: Biotin carboxylase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	83.32Å 106.17Å 121.49Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.00 30.00 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.2 (30.00-2.00) 99.2 (30.00-2.00)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.29 (at 2.00Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.182 , 0.216 0.182 , 0.216	Depositor DCC
R_{free} test set	3678 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	26.6	Xtriage
Anisotropy	0.382	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 41.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7794	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BTN, BCT, MG, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.56	1/3484 (0.0%)	0.75	0/4702
1	B	0.55	1/3484 (0.0%)	0.75	0/4702
All	All	0.56	2/6968 (0.0%)	0.75	0/9404

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	83	GLY	C-N	-7.10	1.23	1.33
1	A	139	GLY	C-N	6.54	1.42	1.33

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3421	0	3445	28	0
1	B	3421	0	3445	26	0
2	A	27	0	12	0	0
2	B	27	0	12	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	16	0	15	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	16	0	15	0	0
5	A	4	0	0	0	0
5	B	4	0	0	0	0
6	A	396	0	0	3	4
6	B	460	0	0	9	4
All	All	7794	0	6944	54	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:106:LYS:HE2	1:B:270:ARG:HH12	1.34	0.91
1:A:172:VAL:HG11	1:A:178:LEU:HD13	1.55	0.89
1:B:106:LYS:HE2	1:B:270:ARG:NH1	1.86	0.88
1:B:428:GLU:H	1:B:428:GLU:CD	1.90	0.80
1:A:1:MET:HE3	6:A:646:HOH:O	1.83	0.77
1:A:370:HIS:CD2	1:A:370:HIS:H	2.13	0.64
1:B:106:LYS:CE	1:B:270:ARG:HH12	2.10	0.64
1:B:407:GLN:HG2	6:B:881:HOH:O	1.98	0.64
1:A:436:ASN:C	1:A:436:ASN:HD22	2.05	0.64
1:A:178:LEU:HD12	1:A:181:SER:HB2	1.81	0.62
1:B:67:ILE:CD1	1:B:94:GLN:HG2	2.29	0.62
1:B:370:HIS:H	1:B:370:HIS:CD2	2.20	0.59
1:B:278:LEU:HD22	1:B:287:ILE:HG21	1.85	0.58
1:A:155:PRO:HB3	1:A:173:ARG:HG2	1.85	0.57
1:A:311:GLU:OE2	1:A:314:ARG:NH1	2.38	0.56
1:A:172:VAL:HG11	1:A:178:LEU:CD1	2.32	0.55
1:B:370:HIS:HE1	6:B:652:HOH:O	1.90	0.55
1:A:1:MET:HE2	1:A:24:LEU:HD13	1.88	0.54
1:A:206:ASN:N	1:A:207:PRO:HD3	2.24	0.53
1:B:252:LEU:HD12	6:B:764:HOH:O	2.09	0.52
1:B:298:PRO:O	1:B:302:MET:HG2	2.09	0.52
1:B:131:VAL:HG22	1:B:285:TYR:HB3	1.93	0.50
1:B:314:ARG:NH1	6:B:879:HOH:O	2.44	0.50
1:A:131:VAL:HG22	1:A:285:TYR:HB3	1.94	0.50
1:A:34:SER:HB2	1:A:37:ARG:NH2	2.28	0.49
1:B:323:ILE:HG12	1:B:327:GLU:HB2	1.94	0.48
1:B:108:GLU:HG3	6:B:696:HOH:O	2.13	0.47
1:B:314:ARG:HD3	6:B:496:HOH:O	2.14	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:105:PRO:HG2	1:B:291:THR:HB	1.95	0.47
1:A:158:ILE:HB	1:A:170:ARG:HB3	1.97	0.47
1:A:167:ARG:HD2	1:A:192:ALA:HB2	1.97	0.47
1:B:206:ASN:O	1:B:206:ASN:CG	2.57	0.47
1:B:251:GLU:H	1:B:251:GLU:CD	2.22	0.46
1:B:423:ARG:HD3	6:B:767:HOH:O	2.15	0.46
1:A:298:PRO:O	1:A:302:MET:HG2	2.15	0.46
1:B:146:ARG:NH1	6:B:683:HOH:O	2.48	0.46
1:A:370:HIS:HE1	6:A:473:HOH:O	1.99	0.45
1:B:159:LYS:HG2	1:B:169:MET:HG2	1.99	0.45
1:A:436:ASN:ND2	1:A:438:HIS:H	2.15	0.45
1:A:160:ALA:HA	1:A:198:VAL:HG12	1.98	0.44
1:A:278:LEU:HD22	1:A:287:ILE:HG21	1.99	0.44
1:A:133:GLY:HA2	1:A:152:ILE:HD13	1.99	0.44
1:A:211:GLU:C	1:A:212:ILE:HD12	2.43	0.43
1:B:407:GLN:HG3	6:B:801:HOH:O	2.18	0.43
1:B:211:GLU:C	1:B:212:ILE:HD12	2.44	0.42
1:A:442:LYS:NZ	6:A:687:HOH:O	2.49	0.42
1:B:369:SER:OG	1:B:371:ILE:HG12	2.20	0.42
1:A:18:LEU:HD21	1:A:22:LYS:HE2	2.02	0.41
1:A:142:MET:SD	1:A:179:ALA:HA	2.61	0.41
1:A:157:ILE:HB	1:A:203:TYR:HD1	1.86	0.41
1:A:140:ASP:O	1:A:142:MET:HG2	2.21	0.41
1:A:251:GLU:CD	1:A:251:GLU:H	2.28	0.41
1:B:170:ARG:CZ	1:B:184:MET:HG2	2.51	0.40
1:A:301:GLU:HG2	1:A:306:VAL:O	2.22	0.40

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:821:HOH:O	6:B:660:HOH:O[2_655]	2.02	0.18
6:A:861:HOH:O	6:B:707:HOH:O[2_655]	2.03	0.17
6:A:831:HOH:O	6:B:790:HOH:O[2_655]	2.06	0.14
6:A:835:HOH:O	6:B:852:HOH:O[2_655]	2.09	0.11

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	442/444 (100%)	425 (96%)	17 (4%)	0	100	100
1	B	442/444 (100%)	430 (97%)	12 (3%)	0	100	100
All	All	884/888 (100%)	855 (97%)	29 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	357/357 (100%)	351 (98%)	6 (2%)	53	60
1	B	357/357 (100%)	350 (98%)	7 (2%)	48	54
All	All	714/714 (100%)	701 (98%)	13 (2%)	51	58

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	125	LYS
1	A	193	PHE
1	A	251	GLU
1	A	278	LEU
1	A	436	ASN
1	A	444	LEU
1	B	138	LEU
1	B	184	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	205	GLU
1	B	234	ARG
1	B	251	GLU
1	B	278	LEU
1	B	444	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	236	HIS
1	A	340	ASN
1	A	370	HIS
1	A	404	ASN
1	A	407	GLN
1	A	426	ASN
1	A	436	ASN
1	B	145	ASN
1	B	221	ASN
1	B	340	ASN
1	B	370	HIS
1	B	404	ASN
1	B	429	ASN
1	B	438	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 2 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	BCT	A	1006	-	3,3,3	0.49	0	2,3,3	0.54	0
4	BTN	A	1004	-	17,17,17	3.06	2 (11%)	23,23,23	1.32	3 (13%)
5	BCT	B	1007	-	3,3,3	0.54	0	2,3,3	0.15	0
2	ADP	B	1001	3	28,29,29	1.54	5 (17%)	43,45,45	1.85	8 (18%)
4	BTN	B	1005	-	17,17,17	3.08	2 (11%)	23,23,23	1.39	2 (8%)
2	ADP	A	1000	3	28,29,29	1.50	5 (17%)	43,45,45	1.84	8 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	BTN	A	1004	-	-	4/7/28/28	0/2/2/2
2	ADP	B	1001	3	-	3/16/32/32	0/3/3/3
4	BTN	B	1005	-	-	3/7/28/28	0/2/2/2
2	ADP	A	1000	3	-	2/16/32/32	0/3/3/3

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	1005	BTN	C2-S1	-9.07	1.68	1.82
4	A	1004	BTN	C2-S1	-8.92	1.68	1.82
4	B	1005	BTN	O3-C3	8.52	1.41	1.23
4	A	1004	BTN	O3-C3	8.50	1.41	1.23
2	A	1000	ADP	C5-C4	5.08	1.48	1.39
2	B	1001	ADP	C5-C4	5.00	1.48	1.39
2	A	1000	ADP	C5-C6	2.95	1.49	1.41
2	B	1001	ADP	C5-C6	2.86	1.49	1.41
2	B	1001	ADP	PA-O3A	2.69	1.62	1.59
2	B	1001	ADP	C8-N7	2.53	1.36	1.31
2	A	1000	ADP	PA-O3A	2.52	1.62	1.59
2	B	1001	ADP	C5-N7	-2.46	1.34	1.39

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1000	ADP	C8-N7	2.30	1.36	1.31
2	A	1000	ADP	C5-N7	-2.21	1.35	1.39

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1001	ADP	C5-C4-N3	-6.10	118.32	126.72
2	A	1000	ADP	C5-C4-N3	-6.07	118.36	126.72
2	A	1000	ADP	N3-C4-N9	4.84	135.40	127.17
2	B	1001	ADP	N3-C4-N9	4.47	134.77	127.17
2	B	1001	ADP	C2-N3-C4	3.90	121.35	111.83
2	A	1000	ADP	C2-N3-C4	3.89	121.33	111.83
4	B	1005	BTN	C4-C2-S1	3.70	109.23	105.03
2	B	1001	ADP	N3-C2-N1	-3.57	123.18	128.58
2	B	1001	ADP	C4-C5-N7	-3.55	106.52	110.58
4	A	1004	BTN	C4-C2-S1	3.40	108.90	105.03
2	A	1000	ADP	C4-C5-N7	-3.39	106.71	110.58
2	A	1000	ADP	N3-C2-N1	-3.37	123.48	128.58
2	A	1000	ADP	C5-N7-C8	2.51	107.40	103.45
2	B	1001	ADP	C5-N7-C8	2.42	107.25	103.45
4	A	1004	BTN	O12-C11-C10	2.36	121.46	114.00
4	B	1005	BTN	O12-C11-C10	2.36	121.45	114.00
2	A	1000	ADP	C4-N9-C8	2.23	108.08	105.74
2	B	1001	ADP	C6-C5-N7	2.12	136.18	132.09
2	B	1001	ADP	C2-N1-C6	2.10	122.18	118.73
2	A	1000	ADP	C6-C5-N7	2.09	136.11	132.09
4	A	1004	BTN	C6-C5-N1	-2.06	110.53	113.18

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1004	BTN	C4-C2-C7-C8
2	B	1001	ADP	PA-O3A-PB-O2B
2	A	1000	ADP	PB-O3A-PA-O2A
2	B	1001	ADP	PB-O3A-PA-O1A
4	B	1005	BTN	C2-C7-C8-C9
4	A	1004	BTN	C2-C7-C8-C9
2	A	1000	ADP	PB-O3A-PA-O1A
4	B	1005	BTN	C9-C10-C11-O11
4	A	1004	BTN	C9-C10-C11-O11
4	A	1004	BTN	C9-C10-C11-O12

Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Atoms
4	B	1005	BTN	C9-C10-C11-O12
2	B	1001	ADP	PB-O3A-PA-O2A

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	444/444 (100%)	0.30	60 (13%) 7 6	15, 26, 76, 80	0
1	B	444/444 (100%)	-0.03	16 (3%) 46 45	17, 26, 50, 69	0
All	All	888/888 (100%)	0.13	76 (8%) 16 15	15, 26, 70, 80	0

All (76) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	179	ALA	6.5
1	A	176	ALA	6.1
1	A	152	ILE	6.0
1	A	148	ILE	5.9
1	A	178	LEU	5.8
1	A	157	ILE	5.4
1	A	166	GLY	5.0
1	A	193	PHE	4.7
1	A	189	ALA	4.6
1	A	192	ALA	4.4
1	A	154	TYR	4.4
1	A	137	PRO	4.4
1	A	149	ALA	4.3
1	A	171	VAL	4.1
1	A	180	GLN	4.1
1	A	191	ALA	4.0
1	A	174	GLY	4.0
1	A	200	MET	4.0
1	A	185	THR	3.9
1	A	138	LEU	3.8
1	A	158	ILE	3.7
1	A	139	GLY	3.7
1	A	165	GLY	3.6
1	A	156	VAL	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	147	ALA	3.5
1	A	196	ASP	3.5
1	A	146	ARG	3.5
1	B	166	GLY	3.3
1	B	164	GLY	3.3
1	A	187	ALA	3.3
1	A	170	ARG	3.2
1	A	188	GLU	3.2
1	A	184	MET	3.2
1	A	172	VAL	3.2
1	B	172	VAL	3.2
1	A	164	GLY	3.2
1	A	145	ASN	3.2
1	A	150	LYS	3.1
1	A	1	MET	3.1
1	A	169	MET	3.1
1	A	144	LYS	3.0
1	B	182	ILE	3.0
1	B	163	GLY	3.0
1	B	165	GLY	3.0
1	B	174	GLY	3.0
1	A	181	SER	2.9
1	A	197	MET	2.8
1	B	235	ARG	2.7
1	B	167	ARG	2.7
1	A	132	PRO	2.7
1	A	131	VAL	2.7
1	A	153	GLY	2.7
1	B	176	ALA	2.7
1	A	155	PRO	2.6
1	A	194	SER	2.6
1	A	167	ARG	2.6
1	B	193	PHE	2.5
1	A	133	GLY	2.5
1	A	195	ASN	2.5
1	A	142	MET	2.5
1	A	143	ASP	2.4
1	A	183	SER	2.4
1	B	168	GLY	2.4
1	A	182	ILE	2.4
1	A	136	GLY	2.4
1	A	151	ARG	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	444	LEU	2.3
1	B	1	MET	2.3
1	A	186	ARG	2.2
1	A	163	GLY	2.2
1	A	199	TYR	2.2
1	B	147	ALA	2.1
1	A	203	TYR	2.1
1	A	161	SER	2.1
1	B	444	LEU	2.1
1	B	171	VAL	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	MG	B	1003	1/1	0.82	0.10	66,66,66,66	0
4	BTN	B	1005	16/16	0.84	0.13	41,43,45,45	0
2	ADP	B	1001	27/27	0.86	0.12	39,44,50,51	0
2	ADP	A	1000	27/27	0.88	0.10	48,50,51,52	0
4	BTN	A	1004	16/16	0.91	0.10	32,35,39,40	0
3	MG	A	1002	1/1	0.96	0.08	46,46,46,46	0
5	BCT	A	1006	4/4	0.97	0.04	19,20,20,20	0
5	BCT	B	1007	4/4	0.98	0.04	22,23,23,23	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around ADP B 1001:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around ADP A 1000:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.