



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 05:42 PM UTC

PDB ID : 6G9T / pdb_00006g9t
Title : CRYSTAL STRUCTURE OF CMY-136 class C BETA-LACTAMASE
Authors : Zavala, A.; Retailleau, P.; Naas, T.; Iorga, B.
Deposited on : 2018-04-11
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4673 (1.60-1.60)
Clashscore	190562	4931 (1.60-1.60)
Ramachandran outliers	187476	4831 (1.60-1.60)
Sidechain outliers	187428	4830 (1.60-1.60)
RSRZ outliers	180081	4672 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	371	4% 93% • •
1	B	371	9% 92% 5% •

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6105 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-lactamase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	359	Total	C	N	O	S	0	2	0
			2813	1805	489	510	9			
1	B	359	Total	C	N	O	S	0	5	0
			2829	1820	489	511	9			

There are 22 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	221	HIS	TYR	engineered mutation	UNP G3F7G9
A	362	PHE	-	expression tag	UNP G3F7G9
A	363	GLU	-	expression tag	UNP G3F7G9
A	364	HIS	-	expression tag	UNP G3F7G9
A	365	HIS	-	expression tag	UNP G3F7G9
A	366	HIS	-	expression tag	UNP G3F7G9
A	367	HIS	-	expression tag	UNP G3F7G9
A	368	HIS	-	expression tag	UNP G3F7G9
A	369	HIS	-	expression tag	UNP G3F7G9
A	370	HIS	-	expression tag	UNP G3F7G9
A	371	HIS	-	expression tag	UNP G3F7G9
B	221	HIS	TYR	engineered mutation	UNP G3F7G9
B	362	PHE	-	expression tag	UNP G3F7G9
B	363	GLU	-	expression tag	UNP G3F7G9
B	364	HIS	-	expression tag	UNP G3F7G9
B	365	HIS	-	expression tag	UNP G3F7G9
B	366	HIS	-	expression tag	UNP G3F7G9
B	367	HIS	-	expression tag	UNP G3F7G9
B	368	HIS	-	expression tag	UNP G3F7G9
B	369	HIS	-	expression tag	UNP G3F7G9
B	370	HIS	-	expression tag	UNP G3F7G9
B	371	HIS	-	expression tag	UNP G3F7G9

- Molecule 2 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		
2	A	1	Total	O	P	0	0
			5	4	1		
2	A	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is ISOPROPYL ALCOHOL (CCD ID: IPA) (formula: C_3H_8O).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	3	1		
4	A	1	Total	C	O	0	1
			8	6	2		
4	B	1	Total	C	O	0	0
			4	3	1		
4	B	1	Total	C	O	0	0
			4	3	1		
4	B	1	Total	C	O	0	0
			4	3	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	224	Total	O	0	0
			224	224		
5	B	172	Total	O	0	0
			172	172		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Beta-lactamase



• Molecule 1: Beta-lactamase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	60.58Å 58.09Å 100.08Å 90.00° 89.97° 90.00°	Depositor
Resolution (Å)	16.10 – 1.60 16.10 – 1.60	Depositor EDS
% Data completeness (in resolution range)	94.2 (16.10-1.60) 94.1 (16.10-1.60)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.09 (at 1.60Å)	Xtriage
Refinement program	BUSTER 2.10.3	Depositor
R, R_{free}	0.234 , 0.264 (Not available) , 0.270	Depositor DCC
R_{free} test set	4268 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å ²)	14.1	Xtriage
Anisotropy	0.662	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.46 , 59.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.015 for -k,-h,-l 0.014 for k,h,-l 0.369 for h,-k,-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	6105	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 13.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, IPA, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.82	0/2894	1.06	3/3945 (0.1%)
1	B	0.84	0/2920	1.10	2/3981 (0.1%)
All	All	0.83	0/5814	1.08	5/7926 (0.1%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	121	ILE	N-CA-C	-8.98	100.91	108.63
1	B	121	ILE	N-CA-C	-8.93	100.95	108.63
1	B	189	ILE	N-CA-C	-8.88	103.23	111.67
1	A	194	ASN	CA-CB-CG	6.04	118.64	112.60
1	A	47	ASP	CA-CB-CG	5.43	118.03	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2813	0	2793	10	0
1	B	2829	0	2825	13	0
2	A	15	0	0	0	0
2	B	10	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	6	0	8	0	0
3	B	12	0	16	0	0
4	A	12	0	24	0	0
4	B	12	0	24	2	0
5	A	224	0	0	1	0
5	B	172	0	0	4	0
All	All	6105	0	5690	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:67:LYS:NZ	1:A:150:TYR:HE1	1.74	0.85
1:A:67:LYS:NZ	1:A:150:TYR:CE1	2.46	0.83
1:B:347:PRO:O	1:B:351:GLU:HG3	1.88	0.73
1:A:67:LYS:HZ3	1:A:150:TYR:HE1	1.35	0.73
1:B:67:LYS:NZ	1:B:150:TYR:HE1	1.87	0.72
1:B:67:LYS:NZ	1:B:150:TYR:CE1	2.59	0.71
4:B:405:IPA:H13	5:B:546:HOH:O	1.99	0.63
1:B:67:LYS:HZ2	1:B:150:TYR:HE1	1.54	0.56
1:A:67:LYS:HZ1	1:A:150:TYR:HE1	1.52	0.55
1:B:67:LYS:HZ3	1:B:150:TYR:HE1	1.54	0.54
1:B:279:LYS:HG3	1:B:282:SER:HB2	1.91	0.53
1:A:194:ASN:HB2	5:A:705:HOH:O	2.12	0.48
1:A:344:TYR:CE2	1:A:349:ARG:HG2	2.49	0.48
1:B:344:TYR:CE2	1:B:349:ARG:HG2	2.49	0.47
1:A:232:ARG:HH22	1:B:37:LYS:HE3	1.80	0.47
1:B:75:GLY:HA2	1:B:78:ILE:HD12	1.97	0.47
1:A:62:LEU:HB3	1:A:65:VAL:HB	1.99	0.44
1:A:258:ARG:HB2	1:A:305:ALA:HB3	1.99	0.44
4:B:405:IPA:H2	5:B:582:HOH:O	2.18	0.43
1:A:75:GLY:HA2	1:A:78:ILE:HD12	1.99	0.43
1:B:258:ARG:HB2	1:B:305:ALA:HB3	2.00	0.43
1:B:279:LYS:HE3	5:B:583:HOH:O	2.19	0.41
1:B:279:LYS:HB3	5:B:671:HOH:O	2.19	0.41
1:B:149:LEU:HD13	1:B:295:ALA:HB2	2.01	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	359/371 (97%)	354 (99%)	5 (1%)	0	100	100
1	B	361/371 (97%)	356 (99%)	5 (1%)	0	100	100
All	All	720/742 (97%)	710 (99%)	10 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	292/301 (97%)	290 (99%)	2 (1%)	76	63
1	B	295/301 (98%)	293 (99%)	2 (1%)	76	63
All	All	587/602 (98%)	583 (99%)	4 (1%)	81	63

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	242[A]	HIS
1	A	242[B]	HIS
1	B	32	VAL
1	B	331	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	302	ASN
1	B	196	GLN
1	B	242	HIS
1	B	256	GLN
1	B	302	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	IPA	A	406[B]	-	3,3,3	0.66	0	3,3,3	0.49	0
3	GOL	B	403	-	5,5,5	0.10	0	5,5,5	0.57	0
4	IPA	A	405	-	3,3,3	0.65	0	3,3,3	0.52	0
4	IPA	B	405	-	3,3,3	0.81	0	3,3,3	0.31	0
4	IPA	B	406	-	3,3,3	0.56	0	3,3,3	0.60	0
4	IPA	A	406[A]	-	3,3,3	0.59	0	3,3,3	0.51	0
2	PO4	A	402	-	4,4,4	1.87	1 (25%)	6,6,6	0.78	0
2	PO4	B	402	-	4,4,4	1.95	0	6,6,6	0.43	0
3	GOL	A	404	-	5,5,5	0.13	0	5,5,5	0.43	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PO4	B	401	-	4,4,4	2.71	2 (50%)	6,6,6	0.78	0
3	GOL	B	404	-	5,5,5	0.09	0	5,5,5	0.37	0
2	PO4	A	403	-	4,4,4	2.57	1 (25%)	6,6,6	0.80	0
2	PO4	A	401	-	4,4,4	1.86	2 (50%)	6,6,6	0.91	0
4	IPA	B	407	-	3,3,3	0.72	0	3,3,3	0.58	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	404	-	-	2/4/4/4	-
3	GOL	B	403	-	-	1/4/4/4	-
3	GOL	A	404	-	-	2/4/4/4	-

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	401	PO4	P-O1	4.34	1.60	1.50
2	A	403	PO4	P-O1	4.31	1.60	1.50
2	A	401	PO4	P-O4	2.15	1.60	1.54
2	B	401	PO4	P-O2	2.11	1.60	1.54
2	A	402	PO4	P-O2	2.04	1.60	1.54
2	A	401	PO4	P-O2	2.01	1.60	1.54

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	404	GOL	C1-C2-C3-O3
3	A	404	GOL	O1-C1-C2-C3
3	B	404	GOL	O2-C2-C3-O3
3	B	403	GOL	C1-C2-C3-O3
3	A	404	GOL	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	405	IPA	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	359/371 (96%)	0.38	14 (3%) 43 45	7, 15, 34, 65	2 (0%)
1	B	359/371 (96%)	0.89	35 (9%) 13 13	11, 18, 35, 75	4 (1%)
All	All	718/742 (96%)	0.63	49 (6%) 23 24	7, 17, 35, 75	6 (0%)

All (49) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	242[A]	HIS	4.3
1	B	354	TRP	3.9
1	B	4	THR	3.9
1	A	6	GLN	3.7
1	A	361	GLN	3.7
1	B	280	ALA	3.6
1	B	5	GLU	3.5
1	B	361	GLN	3.5
1	B	209	VAL	3.5
1	B	242	HIS	3.4
1	B	213	PRO	3.3
1	B	212	SER	3.2
1	B	276[A]	TRP	3.1
1	B	10	ASP	3.0
1	B	309	LYS	3.0
1	A	213	PRO	2.9
1	A	126	ARG	2.9
1	B	286	GLY	2.9
1	B	6	GLN	2.7
1	A	209	VAL	2.7
1	B	211	VAL	2.7
1	B	279	LYS	2.7
1	B	139	GLN	2.6
1	B	24	ALA	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	12	VAL	2.5
1	B	214	GLY	2.5
1	B	312	TRP	2.5
1	A	244	GLN	2.4
1	A	4	THR	2.4
1	A	7	GLN	2.3
1	B	278	LEU	2.3
1	A	142	TRP	2.2
1	A	354	TRP	2.2
1	B	277	PRO	2.2
1	A	211	VAL	2.2
1	B	14	ARG	2.2
1	B	355	ARG	2.2
1	B	282	SER	2.2
1	B	102	GLN	2.1
1	B	360	LEU	2.1
1	B	94	PRO	2.1
1	B	128	LYS	2.1
1	A	102	GLN	2.1
1	B	239	ASP	2.1
1	B	37	LYS	2.1
1	B	103	GLY	2.0
1	B	243	VAL	2.0
1	A	332	LYS	2.0
1	B	333	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	PO4	A	402	5/5	0.68	0.20	40,42,43,44	0
4	IPA	A	405	4/4	0.70	0.18	35,35,35,35	0
4	IPA	B	405	4/4	0.71	0.17	21,22,24,25	0
2	PO4	B	402	5/5	0.73	0.23	54,56,56,57	0
4	IPA	A	406[B]	4/4	0.78	0.16	24,24,25,25	4
4	IPA	A	406[A]	4/4	0.78	0.16	20,20,20,20	4
4	IPA	B	406	4/4	0.79	0.15	31,31,32,33	0
4	IPA	B	407	4/4	0.79	0.18	20,21,23,29	0
2	PO4	A	403	5/5	0.85	0.13	39,39,40,41	0
3	GOL	B	403	6/6	0.86	0.12	15,17,21,23	0
3	GOL	B	404	6/6	0.86	0.11	26,30,32,33	0
3	GOL	A	404	6/6	0.89	0.12	14,18,19,23	0
2	PO4	B	401	5/5	0.92	0.10	18,24,26,29	0
2	PO4	A	401	5/5	0.94	0.10	14,20,23,25	0

6.5 Other polymers [i](#)

There are no such residues in this entry.