



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 08:41 PM UTC

PDB ID : 8G93 / pdb_00008g93
Title : Crystal structures of 17-beta-hydroxysteroid dehydrogenase 13
Authors : Liu, S.
Deposited on : 2023-02-21
Resolution : 1.91 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.91 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1188 (1.92-1.92)
Clashscore	190562	1209 (1.92-1.92)
Ramachandran outliers	187476	1195 (1.92-1.92)
Sidechain outliers	187428	1195 (1.92-1.92)
RSRZ outliers	180081	1188 (1.92-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	315	8% 73% 8% 18%
1	B	315	15% 77% 9% 14%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4403 atoms, of which 44 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hydroxysteroid 17-beta dehydrogenase 13.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	258	Total	C	N	O	S	0	0	0
			2023	1316	338	361	8			
1	B	271	Total	C	N	O	S	0	0	0
			2119	1380	352	379	8			

There are 40 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	initiating methionine	UNP A0A8C0PP93
A	1	GLY	-	cloning artifact	UNP A0A8C0PP93
A	177	GLU	GLY	engineered mutation	UNP A0A8C0PP93
A	178	GLY	VAL	engineered mutation	UNP A0A8C0PP93
A	205	ALA	THR	engineered mutation	UNP A0A8C0PP93
A	293	VAL	ILE	engineered mutation	UNP A0A8C0PP93
A	301	GLY	-	expression tag	UNP A0A8C0PP93
A	302	SER	-	expression tag	UNP A0A8C0PP93
A	303	GLY	-	expression tag	UNP A0A8C0PP93
A	304	HIS	-	expression tag	UNP A0A8C0PP93
A	305	HIS	-	expression tag	UNP A0A8C0PP93
A	306	HIS	-	expression tag	UNP A0A8C0PP93
A	307	HIS	-	expression tag	UNP A0A8C0PP93
A	308	HIS	-	expression tag	UNP A0A8C0PP93
A	309	HIS	-	expression tag	UNP A0A8C0PP93
A	310	HIS	-	expression tag	UNP A0A8C0PP93
A	311	HIS	-	expression tag	UNP A0A8C0PP93
A	312	HIS	-	expression tag	UNP A0A8C0PP93
A	313	HIS	-	expression tag	UNP A0A8C0PP93
A	314	HIS	-	expression tag	UNP A0A8C0PP93
B	0	MET	-	initiating methionine	UNP A0A8C0PP93
B	1	GLY	-	cloning artifact	UNP A0A8C0PP93
B	177	GLU	GLY	engineered mutation	UNP A0A8C0PP93
B	178	GLY	VAL	engineered mutation	UNP A0A8C0PP93
B	205	ALA	THR	engineered mutation	UNP A0A8C0PP93

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	293	VAL	ILE	engineered mutation	UNP A0A8C0PP93
B	301	GLY	-	expression tag	UNP A0A8C0PP93
B	302	SER	-	expression tag	UNP A0A8C0PP93
B	303	GLY	-	expression tag	UNP A0A8C0PP93
B	304	HIS	-	expression tag	UNP A0A8C0PP93
B	305	HIS	-	expression tag	UNP A0A8C0PP93
B	306	HIS	-	expression tag	UNP A0A8C0PP93
B	307	HIS	-	expression tag	UNP A0A8C0PP93
B	308	HIS	-	expression tag	UNP A0A8C0PP93
B	309	HIS	-	expression tag	UNP A0A8C0PP93
B	310	HIS	-	expression tag	UNP A0A8C0PP93
B	311	HIS	-	expression tag	UNP A0A8C0PP93
B	312	HIS	-	expression tag	UNP A0A8C0PP93
B	313	HIS	-	expression tag	UNP A0A8C0PP93
B	314	HIS	-	expression tag	UNP A0A8C0PP93


- Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (CCD ID: NAD) (formula: C₂₁H₂₇N₇O₁₄P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	B	1	Total	C	N	O	P	0	0
			44	21	7	14	2		

- Molecule 3 is 3-fluoro-N-({(1r,4r)-4-[(2-fluorophenoxy)methyl]-1-hydroxycyclohexyl}methyl)-4-hydroxybenzamide (CCD ID: YXW) (formula: C₂₁H₂₃F₂NO₄) (labeled as "Ligand of

Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	F	H	N	O	22	0
			50	21	2	22	1	4		
3	B	1	Total	C	F	H	N	O	22	0
			50	21	2	22	1	4		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	48	Total	O	0	0
			48	48		
4	B	25	Total	O	0	0
			25	25		

- Molecule 1: Hydroxysteroid 17-beta dehydrogenase 13



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	76.73Å 186.32Å 65.32Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	21.09 – 1.91 21.09 – 1.91	Depositor EDS
% Data completeness (in resolution range)	50.0 (21.09-1.91) 50.0 (21.09-1.91)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.99 (at 1.92Å)	Xtriage
Refinement program	BUSTER	Depositor
R, R_{free}	0.200 , 0.224 0.216 , 0.234	Depositor DCC
R_{free} test set	1875 reflections (2.56%)	wwPDB-VP
Wilson B-factor (Å ²)	29.2	Xtriage
Anisotropy	0.077	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 49.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	4403	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.29% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: YXW, NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.85	0/2065	1.35	4/2800 (0.1%)
1	B	0.89	0/2163	1.37	3/2936 (0.1%)
All	All	0.87	0/4228	1.36	7/5736 (0.1%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	96	ASN	CA-C-N	6.13	128.81	120.54
1	A	96	ASN	C-N-CA	6.13	128.81	120.54
1	A	144	ASN	CA-CB-CG	5.73	118.33	112.60
1	B	96	ASN	CA-C-N	5.48	127.94	120.54
1	B	96	ASN	C-N-CA	5.48	127.94	120.54
1	A	265	ILE	N-CA-CB	5.03	117.39	110.54
1	B	265	ILE	N-CA-CB	5.00	117.34	110.54

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2023	0	2090	10	0
1	B	2119	0	2195	9	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	44	0	26	0	0
2	B	44	0	26	0	0
3	A	28	22	0	0	0
3	B	28	22	0	0	0
4	A	48	0	0	0	0
4	B	25	0	0	0	0
All	All	4359	44	4337	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:123:THR:H	1:A:144:ASN:HD21	1.24	0.86
1:B:176:HIS:HB3	1:B:267:LEU:HD21	1.87	0.57
1:B:42:THR:O	1:B:120:ASN:HB3	2.05	0.56
1:A:176:HIS:HB3	1:A:267:LEU:HD21	1.87	0.55
1:A:191:ALA:HB2	1:B:191:ALA:HB2	1.91	0.53
1:A:221:VAL:HG22	1:A:237:LEU:HD13	1.89	0.53
1:B:135:GLU:H	1:B:135:GLU:CD	2.22	0.48
1:A:135:GLU:CD	1:A:135:GLU:H	2.24	0.45
1:A:208:ILE:HG21	1:A:211:ILE:HD12	2.00	0.44
1:B:208:ILE:HG21	1:B:211:ILE:HD12	1.99	0.44
1:B:124:VAL:HG11	1:B:226:THR:HG23	1.99	0.43
1:A:129:LEU:O	1:B:153:LYS:HD3	2.21	0.41
1:A:163:ASN:O	1:A:210:GLY:HA3	2.21	0.41
1:A:42:THR:O	1:A:120:ASN:HB3	2.20	0.40
1:A:176:HIS:CB	1:A:267:LEU:HD21	2.51	0.40
1:B:62:ARG:NH2	1:B:86:THR:HG21	2.36	0.40
1:B:133:LYS:O	1:B:136:GLU:HB2	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	254/315 (81%)	247 (97%)	6 (2%)	1 (0%)	30	19
1	B	267/315 (85%)	257 (96%)	9 (3%)	1 (0%)	30	19
All	All	521/630 (83%)	504 (97%)	15 (3%)	2 (0%)	30	19

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	219	VAL
1	B	219	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	222/271 (82%)	211 (95%)	11 (5%)	22	8
1	B	234/271 (86%)	222 (95%)	12 (5%)	21	8
All	All	456/542 (84%)	433 (95%)	23 (5%)	22	8

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	LEU
1	A	22	LEU
1	A	26	PHE
1	A	30	LYS
1	A	130	LEU
1	A	156	LEU
1	A	201	LEU
1	A	228	ASN
1	A	237	LEU
1	A	252	THR
1	A	267	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	26	PHE
1	B	30	LYS
1	B	130	LEU
1	B	156	LEU
1	B	161	LYS
1	B	201	LEU
1	B	228	ASN
1	B	231	THR
1	B	237	LEU
1	B	252	THR
1	B	267	LEU
1	B	273	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	144	ASN
1	A	148	HIS
1	A	228	ASN
1	B	88	HIS
1	B	120	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	YXW	A	402	-	30,30,30	1.25	3 (10%)	35,42,42	0.83	1 (2%)
3	YXW	B	402	-	30,30,30	1.26	3 (10%)	35,42,42	0.81	1 (2%)
2	NAD	B	401	-	46,48,48	0.36	0	64,73,73	0.48	0
2	NAD	A	401	-	46,48,48	0.40	0	64,73,73	0.49	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	YXW	A	402	-	-	0/15/27/27	0/3/3/3
3	YXW	B	402	-	-	0/15/27/27	0/3/3/3
2	NAD	B	401	-	-	7/30/62/62	0/5/5/5
2	NAD	A	401	-	-	6/30/62/62	0/5/5/5

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	402	YXW	O28-C25	-4.12	1.28	1.36
3	B	402	YXW	O28-C25	-4.09	1.28	1.36
3	A	402	YXW	C25-C24	2.92	1.41	1.39
3	B	402	YXW	C25-C24	2.79	1.41	1.39
3	B	402	YXW	C20-N19	2.40	1.39	1.33
3	A	402	YXW	C20-N19	2.34	1.38	1.33

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	402	YXW	C27-C26-C25	2.76	123.25	120.50
3	A	402	YXW	C27-C26-C25	2.54	123.03	120.50

There are no chirality outliers.


All (13) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
2	A	401	NAD	C5D-O5D-PN-O3
2	A	401	NAD	C5D-O5D-PN-O1N
2	A	401	NAD	O4D-C1D-N1N-C2N
2	B	401	NAD	C5D-O5D-PN-O3
2	B	401	NAD	C5D-O5D-PN-O1N
2	B	401	NAD	O4D-C1D-N1N-C2N
2	A	401	NAD	C5D-O5D-PN-O2N
2	B	401	NAD	C5B-O5B-PA-O3
2	B	401	NAD	C5D-O5D-PN-O2N
2	A	401	NAD	O4D-C1D-N1N-C6N
2	B	401	NAD	O4D-C1D-N1N-C6N
2	A	401	NAD	PA-O3-PN-O2N
2	B	401	NAD	PA-O3-PN-O2N


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	258/315 (81%)	0.32	26 (10%)	12 16	19, 29, 80, 97	0
1	B	271/315 (86%)	1.06	48 (17%)	4 5	17, 34, 152, 164	0
All	All	529/630 (83%)	0.70	74 (13%)	6 8	17, 31, 107, 164	0

All (74) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	9	LEU	12.7
1	B	7	LEU	11.4
1	B	4	ILE	11.3
1	B	10	LEU	11.1
1	B	14	ILE	10.9
1	B	12	LEU	10.3
1	B	8	LEU	10.0
1	B	3	ILE	9.9
1	B	15	ILE	9.6
1	B	11	LEU	9.5
1	B	5	LEU	9.1
1	B	274	PRO	8.9
1	B	16	TYR	8.7
1	B	232	ARG	8.6
1	B	22	LEU	8.0
1	B	13	THR	7.8
1	B	19	LEU	7.8
1	A	272	PHE	7.4
1	B	6	ASP	6.9
1	B	23	VAL	6.9
1	B	18	TYR	6.8
1	B	273	LEU	6.7
1	A	26	PHE	6.2
1	B	21	ALA	6.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	25	VAL	5.9
1	B	24	LYS	5.7
1	B	27	PHE	5.6
1	B	20	GLU	5.5
1	A	18	TYR	4.9
1	B	26	PHE	4.8
1	A	22	LEU	4.8
1	B	231	THR	4.6
1	A	27	PHE	4.5
1	B	272	PHE	4.5
1	B	2	ASN	4.2
1	A	24	LYS	4.1
1	B	17	SER	4.0
1	A	263	TYR	3.9
1	A	267	LEU	3.8
1	B	28	PRO	3.7
1	A	23	VAL	3.7
1	B	266	TYR	3.6
1	B	270	ASP	3.6
1	B	235	PRO	3.4
1	A	16	TYR	3.3
1	A	14	ILE	3.3
1	B	268	ILE	3.3
1	A	19	LEU	3.2
1	B	269	LEU	3.2
1	A	21	ALA	3.1
1	B	236	ILE	3.1
1	B	263	TYR	2.9
1	A	20	GLU	2.9
1	A	265	ILE	2.8
1	A	30	LYS	2.8
1	B	230	SER	2.7
1	B	267	LEU	2.7
1	A	25	VAL	2.7
1	B	237	LEU	2.6
1	B	265	ILE	2.6
1	B	271	LYS	2.5
1	B	30	LYS	2.5
1	A	8	LEU	2.5
1	B	222	ASN	2.5
1	B	29	ARG	2.5
1	A	12	LEU	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	271	LYS	2.3
1	A	15	ILE	2.3
1	A	228	ASN	2.2
1	A	28	PRO	2.2
1	B	229	PRO	2.2
1	A	266	TYR	2.1
1	A	269	LEU	2.1
1	A	229	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	YXW	A	402	28/28	0.92	0.11	34,50,60,60	22
3	YXW	B	402	28/28	0.93	0.07	32,34,37,47	22
2	NAD	B	401	44/44	0.97	0.06	20,27,32,36	0
2	NAD	A	401	44/44	0.98	0.05	21,26,29,31	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around YXW A 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around YXW B 402:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NAD B 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around NAD A 401:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.