



wwPDB EM Validation Summary Report ⓘ

Mar 9, 2026 – 06:56 AM UTC

PDB ID : 5GAR / pdb_00005gar
EMDB ID : EMD-8016
Title : Thermus thermophilus V/A-ATPase, conformation 1
Authors : Schep, D.G.; Zhao, J.; Rubinstein, J.L.
Deposited on : 2016-02-05
Resolution : 6.40 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 6.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.













Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Q-score	-	25397	544 (5.90 - 6.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	577	
1	B	577	
1	C	577	
2	D	457	
2	E	457	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	457	
3	G	186	
3	H	186	
4	I	105	
4	J	105	
5	K	210	
6	L	100	
7	M	323	
8	N	652	
9	O	99	
9	P	99	
9	Q	99	
9	R	99	
9	S	99	
9	T	99	
9	U	99	
9	V	99	
9	W	99	
9	X	99	
9	Y	99	
9	Z	99	

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 23483 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called V-type ATP synthase alpha chain.

Mol	Chain	Residues	Atoms				AltConf	Trace
1	A	577	Total	C	N	O	0	0
			2307	1154	577	576		
1	B	576	Total	C	N	O	0	0
			2303	1152	576	575		
1	C	577	Total	C	N	O	0	0
			2307	1154	577	576		

- Molecule 2 is a protein called V-type ATP synthase beta chain.

Mol	Chain	Residues	Atoms				AltConf	Trace
2	D	457	Total	C	N	O	0	0
			1827	914	457	456		
2	E	457	Total	C	N	O	0	0
			1827	914	457	456		
2	F	457	Total	C	N	O	0	0
			1827	914	457	456		

- Molecule 3 is a protein called V-type ATP synthase subunit E.

Mol	Chain	Residues	Atoms				AltConf	Trace
3	G	184	Total	C	N	O	0	0
			734	368	184	182		
3	H	184	Total	C	N	O	0	0
			734	368	184	182		

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
G	134	MET	LEU	conflict	UNP P74901
G	171	MET	LEU	conflict	UNP P74901
G	178	MET	LEU	conflict	UNP P74901
H	134	MET	LEU	conflict	UNP P74901
H	171	MET	LEU	conflict	UNP P74901

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
H	178	MET	LEU	conflict	UNP P74901

- Molecule 4 is a protein called V-type ATPase subunit G.

Mol	Chain	Residues	Atoms				AltConf	Trace
4	I	100	Total	C	N	O	0	0
			399	200	100	99		
4	J	100	Total	C	N	O	0	0
			399	200	100	99		

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
I	16	GLY	-	insertion	UNP Q72J66
J	16	GLY	-	insertion	UNP Q72J66

- Molecule 5 is a protein called V-type ATP synthase subunit D.

Mol	Chain	Residues	Atoms				AltConf	Trace
5	K	210	Total	C	N	O	0	0
			839	420	210	209		

- Molecule 6 is a protein called V-type ATP synthase subunit F.

Mol	Chain	Residues	Atoms				AltConf	Trace
6	L	100	Total	C	N	O	0	0
			399	200	100	99		

- Molecule 7 is a protein called V-type ATP synthase subunit C.

Mol	Chain	Residues	Atoms				AltConf	Trace
7	M	320	Total	C	N	O	0	0
			1279	640	320	319		

- Molecule 8 is a protein called Archaeal/vacuolar-type H⁺-ATPase subunit I.

Mol	Chain	Residues	Atoms				AltConf	Trace
8	N	619	Total	C	N	O	0	0
			2474	1238	619	617		

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
N	154	ARG	LYS	conflict	UNP H9ZQR4
N	164	ALA	VAL	conflict	UNP H9ZQR4
N	173	PRO	ALA	conflict	UNP H9ZQR4


- Molecule 9 is a protein called Vacuolar type ATP synthase subunit.

Mol	Chain	Residues	Atoms				AltConf	Trace
9	O	80	Total 319	C 160	N 80	O 79	0	0
9	P	80	Total 319	C 160	N 80	O 79	0	0
9	Q	80	Total 319	C 160	N 80	O 79	0	0
9	R	80	Total 319	C 160	N 80	O 79	0	0
9	S	80	Total 319	C 160	N 80	O 79	0	0
9	T	80	Total 319	C 160	N 80	O 79	0	0
9	U	80	Total 319	C 160	N 80	O 79	0	0
9	V	80	Total 319	C 160	N 80	O 79	0	0
9	W	80	Total 319	C 160	N 80	O 79	0	0
9	X	80	Total 319	C 160	N 80	O 79	0	0
9	Y	80	Total 319	C 160	N 80	O 79	0	0
9	Z	80	Total 319	C 160	N 80	O 79	0	0


3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: V-type ATP synthase alpha chain




• Molecule 1: V-type ATP synthase alpha chain






• Molecule 1: V-type ATP synthase alpha chain




- Molecule 2: V-type ATP synthase beta chain

Chain D:




- Molecule 2: V-type ATP synthase beta chain


Chain E:




- Molecule 2: V-type ATP synthase beta chain




- Molecule 3: V-type ATP synthase subunit E




- Molecule 3: V-type ATP synthase subunit E







• Molecule 4: V-type ATPase subunit G



• Molecule 4: V-type ATPase subunit G




• Molecule 5: V-type ATP synthase subunit D




• Molecule 6: V-type ATP synthase subunit F







• Molecule 7: V-type ATP synthase subunit C




• Molecule 8: Archaeal/vacuolar-type H⁺-ATPase subunit I








• Molecule 9: Vacuolar type ATP synthase subunit





• Molecule 9: Vacuolar type ATP synthase subunit




• Molecule 9: Vacuolar type ATP synthase subunit




• Molecule 9: Vacuolar type ATP synthase subunit




- Molecule 9: Vacuolar type ATP synthase subunit



- Molecule 9: Vacuolar type ATP synthase subunit




- Molecule 9: Vacuolar type ATP synthase subunit






- Molecule 9: Vacuolar type ATP synthase subunit







• Molecule 9: Vacuolar type ATP synthase subunit






• Molecule 9: Vacuolar type ATP synthase subunit



• Molecule 9: Vacuolar type ATP synthase subunit



• Molecule 9: Vacuolar type ATP synthase subunit



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	197178	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TECNAI F20	Depositor
Voltage (kV)	200	Depositor
Electron dose ($e^-/\text{\AA}^2$)	35.7	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	6000	Depositor
Magnification	34483	Depositor
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.152	Depositor
Minimum map value	-0.047	Depositor
Average map value	0.001	Depositor
Map value standard deviation	0.007	Depositor
Recommended contour level	0.0382	Depositor
Map size (Å)	371.2, 371.2, 371.2	wwPDB
Map dimensions	256, 256, 256	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.45, 1.45, 1.45	Depositor

5 Model quality ⓘ

5.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	2.54	109/2306 (4.7%)	2.88	206/2881 (7.2%)
1	B	2.56	119/2302 (5.2%)	2.85	220/2876 (7.6%)
1	C	2.55	126/2306 (5.5%)	2.87	260/2881 (9.0%)
2	D	2.60	105/1826 (5.8%)	2.87	189/2281 (8.3%)
2	E	2.55	83/1826 (4.5%)	2.91	190/2281 (8.3%)
2	F	2.53	102/1826 (5.6%)	2.83	180/2281 (7.9%)
3	G	2.52	35/732 (4.8%)	2.93	83/912 (9.1%)
3	H	2.50	40/732 (5.5%)	2.88	86/912 (9.4%)
4	I	2.55	23/398 (5.8%)	2.84	46/496 (9.3%)
4	J	2.54	17/398 (4.3%)	2.90	48/496 (9.7%)
5	K	2.52	48/838 (5.7%)	3.06	117/1046 (11.2%)
6	L	2.55	19/398 (4.8%)	2.93	46/496 (9.3%)
7	M	2.53	71/1278 (5.6%)	3.13	165/1596 (10.3%)
8	N	2.56	129/2472 (5.2%)	3.03	320/3087 (10.4%)
9	O	2.56	19/318 (6.0%)	3.25	47/396 (11.9%)
9	P	2.52	18/318 (5.7%)	3.18	51/396 (12.9%)
9	Q	2.58	16/318 (5.0%)	3.18	47/396 (11.9%)
9	R	2.53	18/318 (5.7%)	3.06	47/396 (11.9%)
9	S	2.55	18/318 (5.7%)	3.01	38/396 (9.6%)
9	T	2.59	20/318 (6.3%)	3.19	48/396 (12.1%)
9	U	2.60	17/318 (5.3%)	3.36	59/396 (14.9%)
9	V	2.62	21/318 (6.6%)	3.30	52/396 (13.1%)
9	W	2.62	21/318 (6.6%)	3.12	37/396 (9.3%)
9	X	2.47	17/318 (5.3%)	3.17	49/396 (12.4%)
9	Y	2.45	13/318 (4.1%)	3.20	49/396 (12.4%)
9	Z	2.62	17/318 (5.3%)	3.37	57/396 (14.4%)
All	All	2.55	1241/23454 (5.3%)	2.97	2737/29274 (9.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
7	M	0	1

The worst 5 of 1241 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	276	GLU	C-N	12.05	1.43	1.32
1	A	442	ASN	CA-C	-11.65	1.43	1.52
8	N	374	LEU	C-N	11.64	1.48	1.33
1	A	60	VAL	CA-C	-11.22	1.43	1.52
1	A	192	ALA	CA-C	-10.96	1.39	1.52

The worst 5 of 2737 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	K	108	PHE	N-CA-C	-13.37	96.17	108.07
2	E	100	ILE	N-CA-C	-13.26	100.54	113.53
1	B	150	PRO	O-C-N	12.47	127.01	121.15
7	M	78	GLY	CA-C-N	12.30	137.24	120.38
7	M	78	GLY	C-N-CA	12.30	137.24	120.38

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
7	M	3	ASP	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2307	0	654	1	0
1	B	2303	0	650	0	0
1	C	2307	0	654	0	0
2	D	1827	0	510	0	0
2	E	1827	0	510	0	0
2	F	1827	0	510	0	0
3	G	734	0	191	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	H	734	0	191	0	0
4	I	399	0	100	0	0
4	J	399	0	100	0	0
5	K	839	0	230	0	0
6	L	399	0	119	0	0
7	M	1279	0	357	0	0
8	N	2474	0	691	0	0
9	O	319	0	109	0	0
9	P	319	0	109	0	0
9	Q	319	0	109	0	0
9	R	319	0	109	0	0
9	S	319	0	109	0	0
9	T	319	0	109	0	0
9	U	319	0	109	0	0
9	V	319	0	109	0	0
9	W	319	0	109	0	0
9	X	319	0	109	0	0
9	Y	319	0	109	0	0
9	Z	319	0	109	0	0
All	All	23483	0	6775	1	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All (1) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:80:LEU:C	1:A:82:GLY:H	2.29	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	575/577 (100%)	504 (88%)	56 (10%)	15 (3%)	4	25
1	B	574/577 (100%)	477 (83%)	73 (13%)	24 (4%)	2	17
1	C	575/577 (100%)	499 (87%)	56 (10%)	20 (4%)	3	20
2	D	455/457 (100%)	370 (81%)	65 (14%)	20 (4%)	2	17
2	E	455/457 (100%)	369 (81%)	66 (14%)	20 (4%)	2	17
2	F	455/457 (100%)	377 (83%)	57 (12%)	21 (5%)	2	16
3	G	180/186 (97%)	162 (90%)	10 (6%)	8 (4%)	2	17
3	H	180/186 (97%)	167 (93%)	9 (5%)	4 (2%)	5	29
4	I	98/105 (93%)	93 (95%)	4 (4%)	1 (1%)	12	48
4	J	98/105 (93%)	94 (96%)	3 (3%)	1 (1%)	12	48
5	K	208/210 (99%)	172 (83%)	25 (12%)	11 (5%)	1	14
6	L	98/100 (98%)	78 (80%)	15 (15%)	5 (5%)	1	15
7	M	318/323 (98%)	307 (96%)	7 (2%)	4 (1%)	9	42
8	N	615/652 (94%)	559 (91%)	43 (7%)	13 (2%)	5	30
9	O	78/99 (79%)	76 (97%)	2 (3%)	0	100	100
9	P	78/99 (79%)	74 (95%)	2 (3%)	2 (3%)	4	25
9	Q	78/99 (79%)	72 (92%)	4 (5%)	2 (3%)	4	25
9	R	78/99 (79%)	73 (94%)	4 (5%)	1 (1%)	9	42
9	S	78/99 (79%)	74 (95%)	4 (5%)	0	100	100
9	T	78/99 (79%)	75 (96%)	2 (3%)	1 (1%)	9	42
9	U	78/99 (79%)	75 (96%)	3 (4%)	0	100	100
9	V	78/99 (79%)	72 (92%)	6 (8%)	0	100	100
9	W	78/99 (79%)	74 (95%)	1 (1%)	3 (4%)	2	19
9	X	78/99 (79%)	74 (95%)	1 (1%)	3 (4%)	2	19
9	Y	78/99 (79%)	74 (95%)	3 (4%)	1 (1%)	9	42
9	Z	78/99 (79%)	74 (95%)	3 (4%)	1 (1%)	9	42
All	All	5820/6157 (94%)	5115 (88%)	524 (9%)	181 (3%)	5	22

5 of 181 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	134	VAL
1	B	258	ARG
1	B	303	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	305	VAL
1	B	466	GLU

5.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-8016. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y



Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)

6.2.1 Primary map



X Index: 128



Y Index: 128



Z Index: 128

The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)

6.3.1 Primary map



X Index: 128



Y Index: 130




Z Index: 178


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)

6.4.1 Primary map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)


6.5.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.0382. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 507 nm^3 ; this corresponds to an approximate mass of 458 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ



*Reported resolution corresponds to spatial frequency of 0.156 Å⁻¹


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-8016 and PDB model 5GAR. Per-residue inclusion information can be found in [section 3](#) on [page 7](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.0382 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.0382).

9.4 Atom inclusion [i](#)



At the recommended contour level, 90% of all backbone atoms, 90% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.0382) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.8970	 0.2510
A	 0.9610	 0.2860
B	 0.9410	 0.2780
C	 0.9570	 0.2790
D	 0.9390	 0.2780
E	 0.9390	 0.2770
F	 0.9350	 0.2770
G	 0.9020	 0.2610
H	 0.8990	 0.2510
I	 0.8220	 0.2220
J	 0.9070	 0.2500
K	 0.8150	 0.2160
L	 0.8150	 0.2420
M	 0.8610	 0.2610
N	 0.8430	 0.2140
O	 0.8400	 0.2010
P	 0.8120	 0.1950
Q	 0.8620	 0.2270
R	 0.8930	 0.2450
S	 0.9250	 0.2260
T	 0.8400	 0.2010
U	 0.7840	 0.1870
V	 0.6330	 0.1060
W	 0.8530	 0.2070
X	 0.8250	 0.1760
Y	 0.8780	 0.1830
Z	 0.6520	 0.1330

