



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:55 PM UTC

PDB ID : 6GB6 / pdb_00006gb6
Title : Structure of H-2Kb with dipeptide GL
Authors : Hafstrand, I.; Sandalova, T.; Achour, A.
Deposited on : 2018-04-13
Resolution : 1.78 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.78 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1365 (1.78-1.78)
Clashscore	190562	1395 (1.78-1.78)
Ramachandran outliers	187476	1382 (1.78-1.78)
Sidechain outliers	187428	1382 (1.78-1.78)
RSRZ outliers	180081	1365 (1.78-1.78)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	276	26% 94% . ..
1	D	276	25% 96% .
2	B	100	% 97% .
2	E	100	2% 97% .

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6719 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called H-2 class I histocompatibility antigen, K-B alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	273	Total	C	N	O	S	0	3	0
			2192	1391	377	414	10			
1	D	276	Total	C	N	O	S	0	2	0
			2199	1390	381	418	10			

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	100	Total	C	N	O	S	0	0	0
			821	523	138	153	7			
2	B	100	Total	C	N	O	S	0	0	0
			821	523	138	153	7			

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
E	0	GLY	ALA	conflict	UNP P01887
E	85	ASP	ALA	variant	UNP P01887
B	0	GLY	ALA	conflict	UNP P01887
B	85	ASP	ALA	variant	UNP P01887

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		
3	E	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is GLYCINE (CCD ID: GLY) (formula: $C_2H_5NO_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			4	2	1	1		
4	D	1	Total	C	N	O	0	0
			4	2	1	1		

- Molecule 5 is LEUCINE (CCD ID: LEU) (formula: C₆H₁₃NO₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			9	6	1	2		
5	D	1	Total	C	N	O	0	0
			9	6	1	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	191	Total	O	0	0
			191	191		
6	D	184	Total	O	0	0
			184	184		
6	E	134	Total	O	0	0
			134	134		
6	B	127	Total	O	0	0
			127	127		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: H-2 class I histocompatibility antigen, K-B alpha chain




- Molecule 1: H-2 class I histocompatibility antigen, K-B alpha chain



- Molecule 2: Beta-2-microglobulin



- Molecule 2: Beta-2-microglobulin



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	66.63Å 82.24Å 89.55Å 90.00° 111.29° 90.00°	Depositor
Resolution (Å)	83.44 – 1.78 83.44 – 1.78	Depositor EDS
% Data completeness (in resolution range)	99.9 (83.44-1.78) 99.9 (83.44-1.78)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.65 (at 1.78Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.207 , 0.232 0.213 , 0.240	Depositor DCC
R_{free} test set	4328 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	22.0	Xtriage
Anisotropy	0.742	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 45.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.53$, $\langle L^2 \rangle = 0.37$	Xtriage
Estimated twinning fraction	0.000 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6719	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 47.23 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.0216e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.61	0/2259	0.78	2/3074 (0.1%)
1	D	0.59	0/2264	0.78	0/3085
2	B	0.62	0/847	0.76	0/1149
2	E	0.59	0/847	0.72	0/1149
All	All	0.60	0/6217	0.77	2/8457 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	178	THR	N-CA-C	5.68	122.91	110.80
1	A	170	ARG	NE-CZ-NH2	5.13	123.81	119.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2192	0	2043	8	0
1	D	2199	0	2037	5	0
2	B	821	0	788	2	0
2	E	821	0	788	2	0
3	A	6	0	8	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	D	12	0	16	0	0
3	E	6	0	8	0	0
4	A	4	0	2	0	0
4	D	4	0	2	0	0
5	A	9	0	11	0	0
5	D	9	0	11	0	0
6	A	191	0	0	1	0
6	B	127	0	0	0	0
6	D	184	0	0	0	0
6	E	134	0	0	0	0
All	All	6719	0	5714	15	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (15) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:7:ILE:HD12	2:B:91:LYS:HE3	1.78	0.66
1:D:51:TRP:CD1	1:D:178:THR:HG1	2.16	0.63
1:A:173:LYS:HG3	1:A:178:THR:HG22	1.84	0.60
1:A:171:TYR:O	1:A:173:LYS:HD2	2.07	0.54
1:D:127:ASN:OD1	1:D:132:THR:HG23	2.08	0.54
1:A:127:ASN:OD1	1:A:132:THR:HG23	2.07	0.53
1:A:72:GLN:HG3	6:A:546:HOH:O	2.10	0.52
1:A:28:VAL:HG23	1:A:33:PHE:CD1	2.46	0.50
2:E:40:LEU:HD23	2:E:45:LYS:HA	1.95	0.48
1:A:126:LEU:HD22	1:A:156:LEU:HD13	1.98	0.44
1:A:117:ALA:HB2	2:B:60:TRP:CE2	2.52	0.44
1:D:260:HIS:CE1	1:D:271:THR:HG22	2.55	0.42
1:D:55:GLU:OE2	1:D:170:ARG:NH1	2.53	0.41
1:D:117:ALA:HB2	2:E:60:TRP:CE2	2.56	0.41
1:A:170:ARG:HH21	1:A:170:ARG:HG3	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	272/276 (99%)	267 (98%)	4 (2%)	1 (0%)	30	16
1	D	276/276 (100%)	270 (98%)	6 (2%)	0	100	100
2	B	98/100 (98%)	96 (98%)	2 (2%)	0	100	100
2	E	98/100 (98%)	96 (98%)	2 (2%)	0	100	100
All	All	744/752 (99%)	729 (98%)	14 (2%)	1 (0%)	48	33

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	178	THR

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	221/234 (94%)	217 (98%)	4 (2%)	51	33
1	D	222/234 (95%)	220 (99%)	2 (1%)	70	59
2	B	93/94 (99%)	93 (100%)	0	100	100
2	E	93/94 (99%)	93 (100%)	0	100	100
All	All	629/656 (96%)	623 (99%)	6 (1%)	68	55

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	45	TYR
1	A	170	ARG
1	A	173	LYS
1	A	178	THR
1	D	45	TYR
1	D	58	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	72	GLN
2	E	29	GLN
2	E	38	GLN
2	B	29	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	D	302	-	5,5,5	0.38	0	5,5,5	0.68	0
3	GOL	A	301	-	5,5,5	0.40	0	5,5,5	0.48	0
3	GOL	E	101	-	5,5,5	0.36	0	5,5,5	0.23	0
3	GOL	D	301	-	5,5,5	0.21	0	5,5,5	0.25	0
5	LEU	A	303	-	6,8,8	0.98	1 (16%)	5,10,10	1.23	1 (20%)
5	LEU	D	304	-	6,8,8	0.94	1 (16%)	5,10,10	1.13	1 (20%)
4	GLY	D	303	-	3,3,4	0.85	0	1,2,4	0.27	0
4	GLY	A	302	-	3,3,4	0.85	0	1,2,4	0.34	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	D	302	-	-	2/4/4/4	-
3	GOL	A	301	-	-	4/4/4/4	-
3	GOL	E	101	-	-	0/4/4/4	-
3	GOL	D	301	-	-	2/4/4/4	-
5	LEU	A	303	-	-	0/8/8/8	-
5	LEU	D	304	-	-	0/8/8/8	-
4	GLY	D	303	-	-	0/0/1/2	-
4	GLY	A	302	-	-	0/0/1/2	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	303	LEU	OXT-C	-2.30	1.23	1.30
5	D	304	LEU	OXT-C	-2.19	1.23	1.30

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	303	LEU	OXT-C-O	-2.61	118.15	124.08
5	D	304	LEU	OXT-C-O	-2.38	118.67	124.08

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	301	GOL	O1-C1-C2-C3
3	D	302	GOL	C1-C2-C3-O3
3	D	301	GOL	O1-C1-C2-O2
3	A	301	GOL	C1-C2-C3-O3
3	D	301	GOL	O1-C1-C2-C3
3	A	301	GOL	O1-C1-C2-O2
3	A	301	GOL	O2-C2-C3-O3
3	D	302	GOL	O2-C2-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	273/276 (98%)	1.18	72 (26%) 1 1	12, 32, 67, 89	3 (1%)
1	D	276/276 (100%)	1.07	70 (25%) 1 1	10, 33, 58, 71	2 (0%)
2	B	100/100 (100%)	0.12	1 (1%) 79 85	14, 22, 35, 50	0
2	E	100/100 (100%)	0.07	2 (2%) 65 73	14, 24, 38, 45	0
All	All	749/752 (99%)	0.85	145 (19%) 3 3	10, 28, 59, 89	5 (0%)

All (145) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	172	LEU	10.4
1	A	179	LEU	8.1
1	A	177	ALA	8.0
1	A	178	THR	6.2
1	D	179	LEU	5.1
2	E	0	GLY	4.8
1	A	180	LEU	4.6
1	A	51	TRP	4.6
1	A	171	TYR	4.6
1	A	2	PRO	4.5
1	D	178	THR	4.3
1	D	142	ILE	4.3
1	A	110	LEU	4.2
1	D	156	LEU	4.1
1	A	132	THR	4.0
1	D	160	LEU	4.0
1	D	130	LEU	3.9
1	A	153	ALA	3.9
1	D	132	THR	3.8
1	A	173	LYS	3.7
1	D	1	GLY	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	49	ALA	3.7
1	A	158	ALA	3.7
1	D	113	TYR	3.6
1	D	133	TRP	3.6
1	A	168	LEU	3.6
1	D	180	LEU	3.6
1	A	195	PRO	3.5
1	D	151	GLY	3.5
1	A	60	TRP	3.5
1	D	158	ALA	3.4
1	D	225	ILE	3.4
1	D	172	LEU	3.4
1	A	150	ALA	3.4
1	D	110	LEU	3.4
1	D	50	ARG	3.4
1	D	103	VAL	3.3
1	A	156	LEU	3.3
1	A	103	VAL	3.3
1	A	129	ASP	3.3
1	A	109	LEU	3.2
1	A	160	LEU	3.2
1	A	50	ARG	3.2
1	A	101	CYS	3.2
1	A	159	TYR	3.2
1	A	62	ARG	3.2
1	D	169	ARG	3.2
1	A	220	ASN	3.1
1	D	16	GLY	3.1
1	D	226	GLN	3.1
1	A	102	GLU	3.1
1	A	113	TYR	3.1
1	A	141	LEU	3.0
1	A	149	GLN	3.0
1	D	51	TRP	3.0
1	D	147	TRP	3.0
1	A	225	ILE	3.0
1	D	129	ASP	3.0
1	D	164	CYS	3.0
1	D	138	MET	3.0
1	D	149	GLN	3.0
1	D	49	ALA	2.9
1	A	57	PRO	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	2	PRO	2.9
1	A	59	TYR	2.9
1	D	60	TRP	2.9
1	D	177	ALA	2.8
1	D	153	ALA	2.8
1	D	126	LEU	2.8
1	A	169	ARG	2.8
1	D	86	ASN	2.8
1	D	148	GLU	2.8
1	A	227	ASP	2.7
1	D	276	PRO	2.7
1	D	165	VAL	2.7
1	A	130	LEU	2.7
1	D	17	LEU	2.7
1	D	176	ASN	2.7
1	A	111	ARG	2.6
1	D	145	HIS	2.6
2	B	29	GLN	2.6
1	A	17	LEU	2.6
1	D	101	CYS	2.6
1	D	128	GLU	2.6
1	A	1	GLY	2.6
1	A	28	VAL	2.6
1	D	150	ALA	2.6
1	D	157	ARG	2.6
1	D	146	LYS	2.5
1	D	171	TYR	2.5
1	A	222	GLU	2.5
1	A	16	GLY	2.5
1	D	162	GLY	2.5
1	D	104	GLY	2.5
1	D	84	TYR	2.4
1	D	227	ASP	2.4
1	A	157	ARG	2.4
1	A	52	MET	2.4
1	D	168	LEU	2.4
1	D	54	GLN	2.4
1	A	142	ILE	2.4
1	A	155	ARG	2.4
1	D	127	ASN	2.3
1	A	128	GLU	2.3
1	A	163	THR	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	159	TYR	2.3
1	D	154	GLU	2.3
1	A	167	TRP	2.3
1	A	193	SER	2.3
1	A	165	VAL	2.3
1	D	28	VAL	2.3
1	A	126	LEU	2.3
1	D	109	LEU	2.3
1	D	59	TYR	2.3
1	A	162	GLY	2.3
2	E	1	ILE	2.3
1	D	52	MET	2.3
1	A	47	PRO	2.2
1	A	230	LEU	2.2
1	D	3	HIS	2.2
1	A	46	GLU	2.2
1	D	40	ALA	2.2
1	D	175	GLY	2.2
1	A	131	LYS	2.2
1	D	193	SER	2.2
1	D	136	ALA	2.2
1	A	133	TRP	2.2
1	A	147	TRP	2.2
1	A	154	GLU	2.2
1	A	151	GLY	2.2
1	A	170	ARG	2.2
1	D	230	LEU	2.1
1	D	144	LYS	2.1
1	D	174	ASN	2.1
1	D	135	ALA	2.1
1	D	134	THR	2.1
1	A	145	HIS	2.1
1	D	56	GLY	2.0
1	A	61	GLU	2.0
1	A	127	ASN	2.0
1	A	79	ARG	2.0
1	A	224	LEU	2.0
1	A	3	HIS	2.0
1	A	196	GLU	2.0
1	A	115	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	GLY	D	303	4/5	0.76	0.16	42,45,45,46	0
3	GOL	D	301	6/6	0.81	0.18	51,52,54,54	0
3	GOL	D	302	6/6	0.84	0.14	46,53,55,57	0
4	GLY	A	302	4/5	0.86	0.12	39,39,41,42	0
3	GOL	A	301	6/6	0.87	0.14	40,45,48,51	0
5	LEU	D	304	9/9	0.89	0.12	40,41,42,43	0
3	GOL	E	101	6/6	0.94	0.08	29,31,33,33	0
5	LEU	A	303	9/9	0.95	0.09	32,35,36,36	0

6.5 Other polymers [i](#)

There are no such residues in this entry.