



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 12:31 PM UTC

PDB ID : 8GB8 / pdb_00008gb8
Title : Crystal structure of SARS-CoV-2 BA.2 receptor binding domain in complex with neutralizing antibody 20A7
Authors : Yuan, M.; Wilson, I.A.
Deposited on : 2023-02-24
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	E	231	30% 76% 9% 16%
2	H	224	3% 90% 7% .
3	L	214	2% 89% 9% .

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 5044 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Spike protein S1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	E	195	Total	C	N	O	S	0	0	0
			1555	1004	262	281	8			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	339	ASP	GLY	variant	UNP P0DTC2
E	371	PHE	SER	variant	UNP P0DTC2
E	373	PRO	SER	variant	UNP P0DTC2
E	375	PHE	SER	variant	UNP P0DTC2
E	376	ALA	THR	variant	UNP P0DTC2
E	405	ASN	ASP	variant	UNP P0DTC2
E	408	SER	ARG	variant	UNP P0DTC2
E	417	ASN	LYS	variant	UNP P0DTC2
E	440	LYS	ASN	variant	UNP P0DTC2
E	477	ASN	SER	variant	UNP P0DTC2
E	478	LYS	THR	variant	UNP P0DTC2
E	484	ALA	GLU	variant	UNP P0DTC2
E	493	ARG	GLN	variant	UNP P0DTC2
E	498	ARG	GLN	variant	UNP P0DTC2
E	501	TYR	ASN	variant	UNP P0DTC2
E	505	HIS	TYR	variant	UNP P0DTC2
E	542	SER	-	expression tag	UNP P0DTC2
E	543	GLY	-	expression tag	UNP P0DTC2
E	544	HIS	-	expression tag	UNP P0DTC2
E	545	HIS	-	expression tag	UNP P0DTC2
E	546	HIS	-	expression tag	UNP P0DTC2
E	547	HIS	-	expression tag	UNP P0DTC2
E	548	HIS	-	expression tag	UNP P0DTC2
E	549	HIS	-	expression tag	UNP P0DTC2


- Molecule 2 is a protein called 20A7 Heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	217	Total	C	N	O	S	0	0	0
			1628	1024	279	319	6			

- Molecule 3 is a protein called 20A7 Light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	L	211	Total	C	N	O	S	0	1	0
			1624	1018	277	325	4			

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	E	1	Total	C	N	O		0	0
			14	8	1	5			

- Molecule 5 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	E	1	Total	C	O	0	0
			4	2	2		
5	L	1	Total	C	O	0	0
			4	2	2		
5	L	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	L	1	Total	C	O	0	0
			6	3	3		

- Molecule 7 is GLYCINE (CCD ID: GLY) (formula: $C_2H_5NO_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	L	1	Total	C	N	O	0	0
			4	2	1	1		

- Molecule 8 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	L	1	Total	C	O	0	0
			7	4	3		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	E	27	Total 27	O 27	0	0
9	H	88	Total 88	O 88	0	0
9	L	79	Total 79	O 79	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Spike protein S1



- Molecule 2: 20A7 Heavy chain



- Molecule 3: 20A7 Light chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	44.54Å 132.18Å 170.47Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.10 – 2.30 43.10 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.3 (43.10-2.30) 99.6 (43.10-2.30)	Depositor EDS
R_{merge}	0.99	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.29 (at 2.29Å)	Xtriage
Refinement program	PHENIX 1.19.2	Depositor
R, R_{free}	0.200 , 0.228 0.196 , 0.226	Depositor DCC
R_{free} test set	2321 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	44.8	Xtriage
Anisotropy	0.449	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 39.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5044	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.85% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, PEG, NAG, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	E	0.10	0/1602	0.31	1/2180 (0.0%)
2	H	0.10	0/1665	0.32	0/2270
3	L	0.10	0/1661	0.33	0/2260
All	All	0.10	0/4928	0.32	1/6710 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	527	PRO	N-CA-CB	6.89	110.58	103.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	1555	0	1473	11	0
2	H	1628	0	1594	8	0
3	L	1624	0	1580	11	0
4	E	14	0	13	0	0
5	E	4	0	6	0	0
5	L	8	0	12	1	0
6	L	6	0	8	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	L	4	0	2	0	0
8	L	7	0	10	0	0
9	E	27	0	0	2	0
9	H	88	0	0	2	0
9	L	79	0	0	5	0
All	All	5044	0	4698	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:373:PRO:O	9:E:701:HOH:O	1.95	0.83
3:L:72:THR:OG1	9:L:401:HOH:O	2.07	0.73
1:E:514:SER:OG	9:E:702:HOH:O	2.06	0.72
3:L:199:GLN:OE1	9:L:403:HOH:O	2.15	0.63
3:L:2:ILE:HG12	3:L:27:GLN:HG2	1.82	0.61
2:H:83:MET:HE2	2:H:86:LEU:HD21	1.84	0.58
5:L:304:EDO:O2	9:L:404:HOH:O	2.18	0.57
2:H:67:ARG:NH2	2:H:90:ASP:OD2	2.35	0.56
3:L:70:ASP:OD1	9:L:405:HOH:O	2.18	0.54
2:H:127:PRO:HB3	2:H:153:TYR:HB3	1.88	0.54
3:L:185:ASP:HA	3:L:188:LYS:HD3	1.89	0.54
3:L:37:GLN:HB2	3:L:47:LEU:HD11	1.89	0.54
1:E:393:THR:HA	1:E:522:ALA:HA	1.94	0.49
3:L:31:GLY:HA3	3:L:91:TYR:O	2.12	0.49
1:E:425:LEU:HD21	1:E:512:VAL:HG11	1.94	0.49
2:H:32:SER:HA	9:H:304:HOH:O	2.12	0.49
3:L:161:GLU:HG3	9:L:415:HOH:O	2.12	0.48
3:L:142[B]:ARG:HE	3:L:163:VAL:HG11	1.78	0.47
1:E:376:ALA:O	1:E:434:ILE:HA	2.14	0.47
1:E:383:SER:HB2	1:E:386:LYS:HG3	1.97	0.46
2:H:89:ASP:OD1	2:H:89:ASP:N	2.48	0.46
1:E:439:ASN:O	1:E:443:SER:OG	2.34	0.45
2:H:99:TYR:CE2	2:H:101:TRP:HB3	2.52	0.45
1:E:358:ILE:HB	1:E:395:VAL:HG22	2.00	0.44
3:L:190:LYS:HG2	3:L:191:VAL:HG23	2.00	0.44
1:E:366:SER:HA	1:E:369:TYR:HD2	1.84	0.43
2:H:209:LYS:HE3	9:H:334:HOH:O	2.17	0.43
1:E:518:LEU:O	1:E:520:ALA:N	2.53	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:100:ASN:HB3	2:H:105:VAL:HB	2.02	0.41
1:E:366:SER:HA	1:E:369:TYR:CD2	2.56	0.40
3:L:145:LYS:HB3	3:L:197:THR:HB	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	193/231 (84%)	185 (96%)	8 (4%)	0	100	100
2	H	213/224 (95%)	211 (99%)	2 (1%)	0	100	100
3	L	210/214 (98%)	205 (98%)	5 (2%)	0	100	100
All	All	616/669 (92%)	601 (98%)	15 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	166/202 (82%)	165 (99%)	1 (1%)	78	89
2	H	184/191 (96%)	182 (99%)	2 (1%)	65	81
3	L	182/183 (100%)	180 (99%)	2 (1%)	65	81

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	532/576 (92%)	527 (99%)	5 (1%)	70	84

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	333	THR
2	H	21	SER
2	H	96	CYS
3	L	7	SER
3	L	14	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	E	417	ASN
2	H	39	GLN
3	L	27	GLN
3	L	38	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	E	601	1	14,14,15	0.28	0	17,19,21	0.41	0
7	GLY	L	302	-	3,3,4	0.59	0	1,2,4	0.31	0
6	GOL	L	301	-	5,5,5	0.91	0	5,5,5	1.09	0
5	EDO	L	304	-	3,3,3	0.42	0	2,2,2	0.41	0
5	EDO	L	305	-	3,3,3	0.42	0	2,2,2	0.37	0
5	EDO	E	602	-	3,3,3	0.44	0	2,2,2	0.39	0
8	PEG	L	303	-	6,6,6	0.11	0	5,5,5	0.10	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	E	601	1	-	2/6/23/26	0/1/1/1
7	GLY	L	302	-	-	0/0/1/2	-
6	GOL	L	301	-	-	2/4/4/4	-
5	EDO	L	304	-	-	0/1/1/1	-
5	EDO	L	305	-	-	0/1/1/1	-
5	EDO	E	602	-	-	0/1/1/1	-
8	PEG	L	303	-	-	1/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	L	301	GOL	O1-C1-C2-O2
4	E	601	NAG	O5-C5-C6-O6
6	L	301	GOL	O1-C1-C2-C3
8	L	303	PEG	O2-C3-C4-O4
4	E	601	NAG	C4-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	L	304	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	E	195/231 (84%)	1.66	69 (35%) 1 1	35, 67, 132, 146	0
2	H	217/224 (96%)	0.12	7 (3%) 50 52	29, 43, 66, 95	0
3	L	211/214 (98%)	0.22	5 (2%) 59 62	20, 43, 67, 74	1 (0%)
All	All	623/669 (93%)	0.64	81 (13%) 7 8	20, 47, 121, 146	1 (0%)

All (81) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	527	PRO	7.1
1	E	368	LEU	6.8
1	E	371	PHE	6.5
1	E	518	LEU	6.4
1	E	373	PRO	6.3
1	E	521	PRO	6.1
1	E	362	VAL	5.7
2	H	141	GLY	5.4
1	E	520	ALA	5.2
1	E	372	ALA	5.0
1	E	392	PHE	4.7
1	E	369	TYR	4.6
1	E	517	LEU	4.5
1	E	365	TYR	4.4
1	E	335	LEU	4.4
1	E	385	THR	4.3
1	E	361	CYS	4.1
2	H	135	SER	4.1
1	E	523	THR	4.0
1	E	478	LYS	3.9
1	E	378	LYS	3.9
1	E	363	ALA	3.9
1	E	522	ALA	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	E	519	HIS	3.9
1	E	333	THR	3.8
2	H	89	ASP	3.8
1	E	367	VAL	3.8
1	E	336	CYS	3.8
1	E	524	VAL	3.8
1	E	387	LEU	3.7
1	E	382	VAL	3.7
1	E	479	PRO	3.6
1	E	486	PHE	3.6
1	E	386	LYS	3.6
1	E	391	CYS	3.6
1	E	432	CYS	3.6
2	H	222	LYS	3.5
1	E	376	ALA	3.5
1	E	390	LEU	3.5
1	E	525	CYS	3.5
1	E	377	PHE	3.3
1	E	489	TYR	3.3
1	E	374	PHE	3.2
1	E	379	CYS	3.1
1	E	358	ILE	3.1
2	H	142	GLY	3.1
1	E	346	ARG	2.9
1	E	413	GLY	2.9
1	E	334	ASN	2.8
1	E	515	PHE	2.8
3	L	7	SER	2.7
3	L	51	ALA	2.7
2	H	96	CYS	2.6
1	E	384	PRO	2.6
3	L	181	LEU	2.6
1	E	526	GLY	2.6
1	E	395	VAL	2.6
1	E	359	SER	2.5
1	E	383	SER	2.5
1	E	354	ASN	2.5
1	E	394	ASN	2.4
1	E	389	ASP	2.4
1	E	388	ASN	2.4
1	E	480	CYS	2.3
1	E	375	PHE	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	E	477	ASN	2.3
1	E	476	GLY	2.3
1	E	514	SER	2.3
1	E	360	ASN	2.3
1	E	415	THR	2.3
1	E	434	ILE	2.3
1	E	485	GLY	2.2
2	H	199	THR	2.2
3	L	190	LYS	2.2
1	E	430	THR	2.2
1	E	490	PHE	2.2
1	E	339	ASP	2.1
1	E	380	TYR	2.1
3	L	143	GLU	2.1
1	E	472	ILE	2.1
1	E	340	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	E	601	14/15	0.36	0.22	64,130,136,140	0
7	GLY	L	302	4/5	0.40	0.39	75,79,86,91	0
5	EDO	L	305	4/4	0.86	0.18	53,68,73,75	0
5	EDO	L	304	4/4	0.86	0.17	45,63,64,65	0
6	GOL	L	301	6/6	0.87	0.17	46,50,61,68	0
5	EDO	E	602	4/4	0.87	0.17	52,54,55,63	0
8	PEG	L	303	7/7	0.92	0.12	37,46,50,66	0

6.5 Other polymers [i](#)

There are no such residues in this entry.