



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 24, 2026 – 10:47 AM UTC

PDB ID : 9GBO / pdb_00009gbo
Title : Human Angiotensin-1 converting enzyme C-domain in complex with a diprolyl inhibitor- SG16
Authors : Gregory, K.S.; Cozier, G.E.; Acharya, K.R.
Deposited on : 2024-07-31
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	597	% 73% 20% • •
2	B	8	25% 25% 50%

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Angiotensin-converting enzyme.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	577	Total	C	N	O	S	0	8	0
			4759	3046	812	877	24			

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	64	GLY	GLU	engineered mutation	UNP P12821
A	90	GLN	ASN	engineered mutation	UNP P12821
A	155	GLN	ASN	engineered mutation	UNP P12821
A	337	GLN	ASN	engineered mutation	UNP P12821
A	586	GLN	ASN	engineered mutation	UNP P12821

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	8	Total	C	N	O	0	0	0
			99	56	4	39			

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Zn 1 1	0	0

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	Cl	0	0
			2	2		

- Molecule 6 is (2 {S})-1-[(2 {S})-2-[(1 {S})-1-[(2 {S})-1-[(2 {S})-2-azanyl-4-oxidanyl-4-oxidanylidene-butanoyl]pyrrolidin-2-yl]-2-oxidanyl-2-oxidanylidene-ethyl]amino]propanoyl]pyrrolidine-2-carboxylic acid (CCD ID: BJ2) (formula: $C_{18}H_{28}N_4O_8$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	O	0	0
			30	18	4	8		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	245	Total	O	0	0
			245	245		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Angiotensin-converting enzyme



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	56.78Å 84.92Å 135.06Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	67.62 – 2.00 67.62 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.8 (67.62-2.00) 99.7 (67.62-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.39 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.8.0425	Depositor
R, R_{free}	0.196 , 0.255 0.200 , 0.260	Depositor DCC
R_{free} test set	2100 reflections (4.67%)	wwPDB-VP
Wilson B-factor (Å ²)	27.3	Xtriage
Anisotropy	0.409	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 34.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5150	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.55% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, BMA, MAN, ZN, NAG, BJ2, FUC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.87	1/4896 (0.0%)	1.60	76/6652 (1.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	209	ARG	NE-CZ	6.42	1.40	1.33

All (76) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	166	THR	CA-CB-OG1	-10.37	94.05	109.60
1	A	609	LEU	N-CA-CB	-9.85	95.05	110.22
1	A	342	GLU	CB-CA-C	-9.53	89.10	110.07
1	A	166	THR	OG1-CB-CG2	-8.96	91.39	109.30
1	A	372	THR	CA-CB-OG1	-8.86	96.31	109.60
1	A	53	ARG	CB-CA-C	-8.79	97.08	110.88
1	A	82	LEU	N-CA-CB	8.58	123.52	110.30
1	A	239	GLU	CB-CG-CD	-8.50	98.15	112.60
1	A	98	GLN	N-CA-CB	8.30	122.33	110.12
1	A	324	THR	CA-CB-OG1	-8.03	97.56	109.60
1	A	585	PRO	N-CA-CB	-7.95	94.90	103.25
1	A	258	HIS	CA-CB-CG	-7.94	105.86	113.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	90	GLN	CB-CA-C	-7.82	97.81	110.79
1	A	75	THR	CA-CB-OG1	-7.80	97.90	109.60
1	A	586	GLN	N-CA-CB	-7.80	97.98	110.46
1	A	345	THR	CA-CB-OG1	-7.51	98.33	109.60
1	A	79	LYS	N-CA-CB	7.28	120.55	109.91
1	A	300	ASP	CB-CA-C	7.16	121.55	109.80
1	A	102	PHE	CA-CB-CG	-7.14	106.66	113.80
1	A	53	ARG	CG-CD-NE	-6.95	96.72	112.00
1	A	94	LYS	CD-CE-NZ	-6.89	89.85	111.90
1	A	359	PHE	CA-CB-CG	-6.88	106.92	113.80
1	A	340	MET	CG-SD-CE	6.85	115.96	100.90
1	A	199	LYS	CB-CG-CD	6.77	126.86	111.30
1	A	572	ARG	O-C-N	-6.71	117.41	121.71
1	A	171	THR	CA-CB-OG1	-6.67	99.60	109.60
1	A	315	MET	CG-SD-CE	6.63	115.49	100.90
1	A	81	LEU	N-CA-CB	6.58	119.55	110.01
1	A	95	TYR	N-CA-CB	6.49	119.77	110.16
1	A	191	ARG	NE-CZ-NH1	-6.48	115.02	121.50
1	A	235	ARG	CA-CB-CG	-6.07	101.96	114.10
1	A	305	MET	CG-SD-CE	-5.96	87.79	100.90
1	A	302	THR	CA-CB-OG1	-5.96	100.67	109.60
1	A	68	ASN	CA-CB-CG	-5.86	106.74	112.60
1	A	353	HIS	CA-CB-CG	-5.82	107.98	113.80
1	A	257	ARG	N-CA-CB	-5.78	101.60	110.16
1	A	168	VAL	N-CA-CB	5.78	117.31	110.55
1	A	396	ASP	CB-CA-C	5.77	119.69	109.65
1	A	225	GLU	CB-CA-C	-5.76	103.39	111.80
1	A	86	MET	CG-SD-CE	5.76	113.57	100.90
1	A	313	ARG	CB-CA-C	5.71	121.63	110.67
1	A	480	ASN	CA-CB-CG	-5.68	106.92	112.60
1	A	561	ARG	CG-CD-NE	-5.66	99.55	112.00
1	A	155	GLN	CB-CA-C	5.62	120.88	112.06
1	A	433	LEU	CA-C-O	-5.58	114.74	120.99
1	A	68	ASN	CB-CA-C	-5.57	100.32	110.63
1	A	480	ASN	CB-CA-C	5.48	119.73	111.18
1	A	117	LYS	N-CA-CB	5.47	118.25	110.16
1	A	154	PRO	N-CA-C	-5.45	102.64	111.19
1	A	165	LEU	N-CA-CB	5.40	118.15	110.16
1	A	572	ARG	CB-CA-C	-5.39	100.10	109.32
1	A	260	GLY	CA-C-N	5.36	127.72	120.38
1	A	260	GLY	C-N-CA	5.36	127.72	120.38
1	A	348	ARG	CB-CA-C	-5.35	100.92	109.75

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	188	LYS	CG-CD-CE	5.35	123.61	111.30
1	A	394	TYR	CA-CB-CG	5.35	123.53	113.90
1	A	191	ARG	NE-CZ-NH2	5.35	124.01	119.20
1	A	191	ARG	CA-CB-CG	-5.32	103.47	114.10
1	A	440	ASP	CB-CA-C	-5.32	101.64	110.68
1	A	449	LYS	CB-CA-C	5.31	119.61	110.79
1	A	75	THR	CA-C-N	5.31	127.71	120.54
1	A	75	THR	C-N-CA	5.31	127.71	120.54
1	A	391	PHE	CB-CA-C	-5.29	102.01	110.79
1	A	343	LYS	CB-CG-CD	-5.27	99.17	111.30
1	A	103	ASP	CA-CB-CG	5.26	117.86	112.60
1	A	83	GLN	CB-CA-C	5.20	119.43	110.79
1	A	123	GLU	N-CA-CB	-5.20	101.70	110.49
1	A	145	THR	CA-CB-OG1	-5.14	101.89	109.60
1	A	143	GLU	CB-CA-C	-5.10	102.32	110.79
1	A	79	LYS	CB-CA-C	-5.09	103.12	110.96
1	A	364	ASP	CA-CB-CG	5.07	117.67	112.60
1	A	96	GLY	CA-C-N	5.05	127.01	120.44
1	A	96	GLY	C-N-CA	5.05	127.01	120.44
1	A	307	LYS	N-CA-CB	5.01	118.05	110.14
1	A	413	ILE	CA-C-N	5.00	125.53	119.98
1	A	413	ILE	C-N-CA	5.00	125.53	119.98

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	191	ARG	Sidechain
1	A	235	ARG	Sidechain
1	A	252	ARG	Sidechain
1	A	314	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4759	0	4586	55	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	99	0	85	2	0
3	A	1	0	0	0	0
4	A	14	0	13	0	0
5	A	2	0	0	0	0
6	A	30	0	0	3	0
7	A	245	0	0	3	0
All	All	5150	0	4684	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:705:BJ2:C27	6:A:705:BJ2:N23	1.69	1.21
1:A:83:GLN:HB2	7:A:1017:HOH:O	1.64	0.95
1:A:66:ASN:O	1:A:70:ASN:ND2	2.13	0.81
2:B:3:BMA:H2	2:B:4:MAN:O5	1.84	0.78
2:B:6:MAN:O6	2:B:7:NAG:H82	1.86	0.75
1:A:301:THR:HG21	1:A:375:LEU:HD13	1.75	0.67
1:A:71:THR:O	1:A:72:ASN:HB2	1.96	0.65
1:A:187:ASP:HA	1:A:191:ARG:HD2	1.78	0.64
1:A:75:THR:HG22	1:A:79:LYS:NZ	2.13	0.64
1:A:67:TRP:CD1	1:A:366:ARG:NH1	2.66	0.63
1:A:57:VAL:O	1:A:61:GLU:HG3	2.00	0.62
1:A:455:ILE:HD13	1:A:592:MET:HE2	1.81	0.60
1:A:335:PHE:C	1:A:335:PHE:CD1	2.81	0.58
1:A:470:ARG:HH22	1:A:484[B]:GLU:CD	2.11	0.58
1:A:590:SER:HB2	7:A:967:HOH:O	2.03	0.58
1:A:562:LEU:HB3	1:A:566:MET:HE2	1.85	0.56
1:A:312:PRO:HG2	1:A:342:GLU:HG3	1.88	0.55
1:A:267:GLU:OE1	1:A:613:LYS:HD3	2.07	0.54
1:A:457:PHE:CE2	1:A:461:SER:HB3	2.44	0.53
6:A:705:BJ2:N23	6:A:705:BJ2:C28	2.64	0.53
1:A:433:LEU:HD23	1:A:434:SER:N	2.23	0.52
6:A:705:BJ2:C27	6:A:705:BJ2:C21	2.81	0.51
1:A:448:MET:HE2	1:A:599:LEU:HD11	1.92	0.51
1:A:344:PRO:CG	1:A:350:VAL:HG11	2.40	0.51
1:A:161:LEU:O	1:A:166:THR:HG23	2.11	0.51
1:A:75:THR:HG22	1:A:79:LYS:HZ3	1.74	0.50
1:A:600:LEU:HD21	1:A:604:ARG:NH1	2.27	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:433:LEU:HD22	1:A:435:SER:H	1.77	0.50
1:A:613:LYS:NZ	1:A:617:PRO:HG3	2.27	0.49
1:A:67:TRP:NE1	1:A:366:ARG:HH11	2.12	0.48
1:A:433:LEU:HD23	1:A:434:SER:H	1.79	0.47
1:A:560:GLN:HE21	1:A:564:THR:HG23	1.79	0.47
1:A:480:ASN:O	1:A:481:TYR:C	2.55	0.47
1:A:76:GLU:HA	1:A:79:LYS:HE2	1.97	0.47
1:A:613:LYS:HE2	1:A:613:LYS:HB3	1.28	0.46
1:A:169:MET:O	1:A:276:GLY:HA2	2.17	0.45
1:A:574:TRP:N	1:A:575:PRO:CD	2.80	0.45
1:A:118:LYS:HB2	1:A:118:LYS:HE2	1.46	0.45
1:A:143:GLU:O	1:A:147[B]:SER:HB2	2.17	0.45
1:A:594[B]:SER:O	1:A:595:TYR:C	2.60	0.44
1:A:326:LEU:O	1:A:559:GLY:HA3	2.18	0.43
1:A:398:PRO:HG2	1:A:401:LEU:HD12	2.00	0.43
1:A:67:TRP:CD1	1:A:366:ARG:HH11	2.37	0.43
1:A:496:CYS:HB2	1:A:497:PRO:HD2	2.01	0.43
1:A:59:TRP:CD1	1:A:88:ILE:HD11	2.54	0.42
1:A:361:ASN:OD1	1:A:361:ASN:C	2.61	0.42
1:A:433:LEU:CD2	1:A:435:SER:H	2.32	0.42
1:A:62:TYR:CD1	1:A:62:TYR:C	2.97	0.42
1:A:368:LYS:NZ	7:A:801:HOH:O	2.34	0.42
1:A:511:LYS:O	1:A:515:PRO:HD2	2.20	0.41
1:A:341:LEU:O	1:A:371:THR:OG1	2.22	0.41
1:A:243:LEU:HD13	1:A:600:LEU:HB2	2.03	0.41
1:A:153:HIS:HB2	1:A:156:GLY:HA3	2.03	0.41
1:A:480:ASN:O	1:A:484[A]:GLU:HG2	2.21	0.41
1:A:568:LEU:HD11	1:A:580:LEU:CD1	2.50	0.41
1:A:606:GLU:O	1:A:609:LEU:HB3	2.21	0.41
1:A:613:LYS:HZ2	1:A:617:PRO:HG3	1.85	0.40
1:A:537:LEU:HD11	1:A:562:LEU:HG	2.02	0.40
1:A:67:TRP:NE1	1:A:366:ARG:NH1	2.68	0.40
1:A:560:GLN:HE21	1:A:564:THR:CG2	2.34	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	582/597 (98%)	563 (97%)	17 (3%)	2 (0%)	36	35

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	72	ASN
1	A	353	HIS

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	511/520 (98%)	489 (96%)	22 (4%)	26	25

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	43	GLU
1	A	48	VAL
1	A	86	MET
1	A	104	VAL
1	A	118	LYS
1	A	130	GLN
1	A	133	GLU
1	A	138	ILE
1	A	155	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	188	LYS
1	A	235	ARG
1	A	243	LEU
1	A	275	LEU
1	A	312	PRO
1	A	335	PHE
1	A	345	THR
1	A	394	TYR
1	A	399	VAL
1	A	440	ASP
1	A	539	GLN
1	A	585	PRO
1	A	613	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	60	ASN
1	A	153	HIS
1	A	241	GLN
1	A	560	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

8 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	B	1	1,2	14,14,15	0.37	0	17,19,21	1.26	3 (17%)
2	NAG	B	2	2	14,14,15	0.43	0	17,19,21	1.42	2 (11%)
2	BMA	B	3	2	11,11,12	1.24	1 (9%)	15,15,17	1.82	4 (26%)
2	MAN	B	4	2	11,11,12	0.92	1 (9%)	15,15,17	2.34	4 (26%)
2	NAG	B	5	2	14,14,15	0.30	0	17,19,21	0.70	0
2	MAN	B	6	2	11,11,12	1.54	2 (18%)	15,15,17	1.50	2 (13%)
2	NAG	B	7	2	14,14,15	0.48	0	17,19,21	1.34	3 (17%)
2	FUC	B	8	2	10,10,11	0.40	0	14,14,16	0.66	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	2/6/23/26	0/1/1/1
2	BMA	B	3	2	-	2/2/19/22	0/1/1/1
2	MAN	B	4	2	-	2/2/19/22	0/1/1/1
2	NAG	B	5	2	-	4/6/23/26	0/1/1/1
2	MAN	B	6	2	-	1/2/19/22	0/1/1/1
2	NAG	B	7	2	-	2/6/23/26	0/1/1/1
2	FUC	B	8	2	-	-	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	6	MAN	O5-C5	3.39	1.50	1.43
2	B	3	BMA	O5-C5	2.91	1.49	1.43
2	B	6	MAN	C2-C3	2.65	1.56	1.52
2	B	4	MAN	O5-C5	2.06	1.47	1.43

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	4	MAN	C1-O5-C5	5.59	119.67	112.19
2	B	4	MAN	C1-C2-C3	4.62	116.37	109.64

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	6	MAN	C1-O5-C5	4.32	117.98	112.19
2	B	2	NAG	C2-N2-C7	-4.03	117.49	122.90
2	B	7	NAG	C1-O5-C5	3.79	117.26	112.19
2	B	3	BMA	O3-C3-C4	3.50	118.63	110.38
2	B	4	MAN	C3-C4-C5	-3.01	104.78	110.23
2	B	3	BMA	C3-C4-C5	-2.91	104.96	110.23
2	B	1	NAG	C1-C2-N2	2.88	114.97	110.43
2	B	3	BMA	C1-O5-C5	2.85	116.00	112.19
2	B	2	NAG	O3-C3-C2	-2.65	103.89	109.40
2	B	7	NAG	O3-C3-C2	2.60	114.79	109.40
2	B	3	BMA	O4-C4-C3	2.52	116.31	110.38
2	B	1	NAG	O3-C3-C2	-2.36	104.49	109.40
2	B	6	MAN	O2-C2-C3	2.34	115.00	110.15
2	B	4	MAN	O4-C4-C5	2.26	114.90	109.32
2	B	7	NAG	C1-C2-N2	2.17	113.86	110.43
2	B	1	NAG	O5-C5-C4	-2.16	105.57	110.83

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	2	NAG	C8-C7-N2-C2
2	B	2	NAG	O7-C7-N2-C2
2	B	5	NAG	C8-C7-N2-C2
2	B	5	NAG	O7-C7-N2-C2
2	B	4	MAN	O5-C5-C6-O6
2	B	4	MAN	C4-C5-C6-O6
2	B	3	BMA	C4-C5-C6-O6
2	B	7	NAG	C8-C7-N2-C2
2	B	7	NAG	O7-C7-N2-C2
2	B	3	BMA	O5-C5-C6-O6
2	B	6	MAN	C4-C5-C6-O6
2	B	5	NAG	C4-C5-C6-O6
2	B	5	NAG	O5-C5-C6-O6

There are no ring outliers.

4 monomers are involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	7	NAG	1	0
2	B	6	MAN	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	3	BMA	1	0
2	B	4	MAN	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	BJ2	A	705	3	31,31,31	3.86	13 (41%)	37,44,44	2.34	10 (27%)
4	NAG	A	702	1	14,14,15	0.49	0	17,19,21	1.16	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	BJ2	A	705	3	-	5/36/56/56	0/2/2/2
4	NAG	A	702	1	-	1/6/23/26	0/1/1/1

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	705	BJ2	C27-N23	11.53	1.69	1.47
6	A	705	BJ2	C08-N12	-8.85	1.36	1.46
6	A	705	BJ2	C21-N23	7.17	1.50	1.34
6	A	705	BJ2	C26-C27	-5.59	1.40	1.53
6	A	705	BJ2	C09-C08	5.55	1.62	1.53
6	A	705	BJ2	C04-N03	5.15	1.52	1.46
6	A	705	BJ2	C13-N12	4.67	1.45	1.34
6	A	705	BJ2	C11-N12	4.60	1.56	1.47
6	A	705	BJ2	C10-C09	-3.95	1.35	1.51
6	A	705	BJ2	C24-N23	-3.68	1.40	1.47
6	A	705	BJ2	C25-C24	3.03	1.62	1.51
6	A	705	BJ2	C25-C26	2.58	1.62	1.51
6	A	705	BJ2	C04-C08	-2.18	1.51	1.54

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	705	BJ2	O29-C28-O30	-6.97	108.27	124.08
6	A	705	BJ2	O29-C28-C27	6.56	134.76	113.34
6	A	705	BJ2	O19-C18-C17	4.63	128.41	114.00
6	A	705	BJ2	C25-C26-C27	4.12	112.72	104.14
4	A	702	NAG	C1-O5-C5	3.67	117.11	112.19
6	A	705	BJ2	C10-C09-C08	-2.88	99.39	104.21
6	A	705	BJ2	O19-C18-O20	-2.64	116.55	123.33
6	A	705	BJ2	O20-C18-C17	-2.55	115.00	122.84
6	A	705	BJ2	C11-N12-C08	-2.53	107.70	111.69

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	705	BJ2	C13-C15-N16	2.30	117.58	109.20
6	A	705	BJ2	C26-C27-C28	2.16	115.86	111.26

There are no chirality outliers.

All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	A	705	BJ2	C05-C04-C08-C09
6	A	705	BJ2	N03-C04-C08-C09
6	A	705	BJ2	N16-C15-C17-C18
6	A	705	BJ2	C05-C04-C08-N12
6	A	705	BJ2	O14-C13-C15-N16
4	A	702	NAG	O7-C7-N2-C2

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	705	BJ2	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	577/597 (96%)	-0.14	8 (1%) 73 73	7, 32, 61, 91	8 (1%)

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	82	LEU	3.9
1	A	438	GLY	3.1
1	A	86	MET	3.0
1	A	344	PRO	2.6
1	A	345	THR	2.4
1	A	73	ILE	2.3
1	A	351	VAL	2.3
1	A	297	PRO	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	B	5	14/15	0.67	0.15	64,88,94,101	0
2	MAN	B	6	11/12	0.76	0.13	57,67,78,79	0
2	BMA	B	3	11/12	0.78	0.10	67,74,90,93	0
2	NAG	B	7	14/15	0.80	0.14	33,56,65,67	14

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	MAN	B	4	11/12	0.82	0.12	51,76,94,102	0
2	NAG	B	2	14/15	0.86	0.10	50,57,63,70	0
2	FUC	B	8	10/11	0.90	0.09	42,51,61,68	0
2	NAG	B	1	14/15	0.92	0.08	44,49,64,71	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	A	702	14/15	0.72	0.14	71,86,97,97	0
6	BJ2	A	705	30/30	0.91	0.10	27,39,50,58	0
5	CL	A	704	1/1	0.99	0.06	24,24,24,24	0
3	ZN	A	701	1/1	1.00	0.01	25,25,25,25	0
5	CL	A	703	1/1	1.00	0.03	23,23,23,23	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around BJ2 A 705:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around ZN A 701:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.