



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 24, 2026 – 10:15 PM EDT

PDB ID : 1GC8 / pdb_00001gc8
Title : THE CRYSTAL STRUCTURE OF THERMUS THERMOPHILUS 3-ISOPROPYLMALATE DEHYDROGENASE MUTATED AT 172TH FROM ALA TO PHE
Authors : Qu, C.; Akanuma, S.; Tanaka, N.; Moriyama, H.; Oshima, T.
Deposited on : 2000-07-27
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	345	 74% 23% ..
1	B	345	 71% 27% .

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5275 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3-ISOPROPYLMALATE DEHYDROGENASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	345	Total	C	N	O	S	0	0	0
			2601	1660	452	483	6			
1	B	345	Total	C	N	O	S	0	0	0
			2601	1660	452	483	6			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1085	ARG	SER	SEE REMARK 999	UNP Q5SIY4
A	1172	PHE	ALA	engineered mutation	UNP Q5SIY4
B	85	ARG	SER	SEE REMARK 999	UNP Q5SIY4
B	172	PHE	ALA	engineered mutation	UNP Q5SIY4


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	38	Total	O	0	0
			38	38		
2	B	35	Total	O	0	0
			35	35		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: 3-ISOPROPYLMALATE DEHYDROGENASE



• Molecule 1: 3-ISOPROPYLMALATE DEHYDROGENASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	55.26Å 87.83Å 71.19Å 90.00° 100.02° 90.00°	Depositor
Resolution (Å)	10.00 – 2.50 10.00 – 2.50	Depositor EDS
% Data completeness (in resolution range)	89.4 (10.00-2.50) 88.3 (10.00-2.50)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.89 (at 2.50Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.170 , 0.258 0.155 , 0.231	Depositor DCC
R_{free} test set	2024 reflections (9.84%)	wwPDB-VP
Wilson B-factor (Å ²)	38.8	Xtriage
Anisotropy	0.080	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.23 , 56.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5275	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.69% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.46	0/2657	1.01	15/3605 (0.4%)
1	B	0.47	0/2657	0.98	12/3605 (0.3%)
All	All	0.47	0/5314	0.99	27/7210 (0.4%)

There are no bond length outliers.

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	135	THR	N-CA-C	8.86	120.94	111.28
1	B	257	LEU	N-CA-C	7.41	120.30	109.04
1	A	1006	LEU	CA-C-N	7.05	126.58	119.24
1	A	1006	LEU	C-N-CA	7.05	126.58	119.24
1	A	1284	ILE	N-CA-C	6.77	123.43	109.34
1	A	1257	LEU	N-CA-C	6.63	119.47	110.31
1	A	1206	TYR	CA-C-N	6.32	125.94	119.56
1	A	1206	TYR	C-N-CA	6.32	125.94	119.56
1	A	1283	GLY	N-CA-C	-6.28	100.27	110.95
1	A	1257	LEU	CA-C-N	6.07	127.43	119.84
1	A	1257	LEU	C-N-CA	6.07	127.43	119.84
1	B	134	LEU	N-CA-C	5.96	120.62	112.04
1	A	1324	PRO	CA-C-N	5.69	125.81	119.32
1	A	1324	PRO	C-N-CA	5.69	125.81	119.32
1	B	141	GLY	N-CA-C	5.67	117.99	111.36
1	A	1051	GLU	CA-C-N	5.63	125.47	119.28
1	A	1051	GLU	C-N-CA	5.63	125.47	119.28
1	B	12	GLY	CA-C-N	5.62	125.08	119.24
1	B	12	GLY	C-N-CA	5.62	125.08	119.24
1	B	74	GLY	CA-C-N	5.53	125.88	119.47
1	B	74	GLY	C-N-CA	5.53	125.88	119.47
1	A	1134	LEU	N-CA-C	5.33	118.11	111.24
1	B	154	THR	N-CA-C	5.32	117.06	108.34
1	A	1178	LYS	N-CA-C	5.24	118.62	111.39
1	B	6	LEU	CA-C-N	5.16	124.61	119.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	6	LEU	C-N-CA	5.16	124.61	119.24
1	B	279	ILE	N-CA-C	-5.10	108.00	112.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2601	0	2634	62	0
1	B	2601	0	2637	64	0
2	A	38	0	0	0	0
2	B	35	0	0	2	0
All	All	5275	0	5271	117	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (117) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1133:GLU:HG2	1:A:1165:VAL:HG11	1.46	0.96
1:A:1083:LYS:HD2	1:A:1083:LYS:H	1.35	0.89
1:A:1153:ASN:HD22	1:B:191:VAL:HG23	1.45	0.80
1:B:115:LEU:HD21	1:B:327:LEU:HD22	1.66	0.78
1:A:1055:GLU:HA	1:A:1058:ARG:NH1	2.01	0.75
1:A:1116:SER:O	1:B:119:LYS:HE2	1.89	0.72
1:B:219:MET:HE3	1:B:223:LEU:HG	1.72	0.70
1:B:99:LEU:HA	1:B:263:GLY:HA3	1.74	0.69
1:A:1030:GLU:HB2	1:A:1032:LEU:HD13	1.75	0.69
1:B:319:LEU:HD22	1:B:327:LEU:HD11	1.76	0.67
1:A:1082:ARG:NE	1:A:1082:ARG:H	1.92	0.66
1:A:1055:GLU:HA	1:A:1058:ARG:HH12	1.60	0.66
1:A:1153:ASN:ND2	1:B:191:VAL:HG23	2.11	0.65
1:B:246:LEU:O	1:B:249:VAL:HG22	1.97	0.63

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1118:LEU:HG	1:A:1122:ILE:HD11	1.81	0.63
1:A:1250:LEU:HB2	1:A:1251:PRO:HD3	1.80	0.62
1:B:183:VAL:HG21	1:B:219:MET:HG2	1.80	0.62
1:B:164:ARG:HH11	1:B:164:ARG:HG3	1.64	0.62
1:A:1008:GLY:HA3	1:A:1071:SER:O	2.01	0.61
1:B:94:ARG:HG2	1:B:94:ARG:HH11	1.66	0.60
1:B:82:ARG:HG2	1:B:85:ARG:NH2	2.18	0.59
1:A:1191:VAL:HG23	1:B:153:ASN:HD22	1.68	0.58
1:A:1292:LEU:HD21	1:A:1315:VAL:HG21	1.85	0.58
1:B:306:GLU:H	1:B:306:GLU:CD	2.12	0.57
1:A:1126:VAL:HG11	1:A:1250:LEU:HD23	1.87	0.57
1:A:1246:LEU:O	1:A:1249:VAL:HG22	2.05	0.57
1:B:69:LEU:HD23	1:B:70:GLY:N	2.20	0.56
1:B:1:MET:HE2	1:B:34:LEU:HD13	1.88	0.56
1:B:81:PRO:HD2	1:B:84:ILE:HD11	1.88	0.56
1:A:1185:LYS:NZ	1:A:1187:ASN:HD21	2.04	0.55
1:B:179:HIS:HB2	2:B:749:HOH:O	2.05	0.55
1:A:1006:LEU:O	1:A:1069:LEU:HD23	2.07	0.55
1:A:1286:ASN:ND2	1:A:1288:THR:H	2.03	0.55
1:A:1156:ARG:HG3	1:B:152:TRP:HB3	1.87	0.55
1:A:1313:ASP:O	1:A:1317:LYS:HG2	2.07	0.54
1:B:2:LYS:HA	1:B:35:ALA:O	2.08	0.53
1:B:55:GLU:N	1:B:56:PRO:HD2	2.24	0.53
1:B:69:LEU:HB3	1:B:270:GLU:HB3	1.90	0.52
1:B:8:GLY:HA3	1:B:71:SER:O	2.09	0.52
1:B:323:PRO:O	1:B:329:GLY:HA3	2.09	0.52
1:A:1081:PRO:HG2	1:A:1084:ILE:HG22	1.90	0.52
1:B:123:ALA:O	1:B:126:VAL:HG23	2.09	0.52
1:B:225:ARG:HB3	1:B:225:ARG:NH1	2.25	0.52
1:A:1249:VAL:HG21	1:B:224:VAL:HG11	1.91	0.52
1:B:54:PRO:HB2	1:B:56:PRO:HD2	1.91	0.51
1:B:167:ARG:HH21	1:B:202:VAL:HG22	1.74	0.51
1:B:109:PHE:HB2	1:B:112:LEU:HD23	1.91	0.51
1:A:1083:LYS:HD2	1:A:1083:LYS:N	2.14	0.51
1:A:1205:GLY:C	1:A:1207:PRO:HD3	2.35	0.51
1:B:184:ASP:O	1:B:215:TYR:HA	2.10	0.51
1:B:45:ALA:HB1	1:B:54:PRO:HD3	1.94	0.50
1:B:313:ASP:O	1:B:317:LYS:HG2	2.11	0.50
1:A:1319:LEU:HD22	1:A:1327:LEU:HD22	1.92	0.50
1:B:238:ILE:HG22	2:B:701:HOH:O	2.10	0.50
1:A:1102:ASN:HB2	1:A:1134:LEU:HG	1.94	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:87:GLU:H	1:B:87:GLU:CD	2.19	0.50
1:A:1141:GLY:HA3	1:A:1154:THR:O	2.12	0.50
1:B:74:GLY:HA3	1:B:77:TRP:HZ3	1.77	0.49
1:A:1057:THR:O	1:A:1061:VAL:HG13	2.13	0.49
1:A:1106:ALA:HB3	1:A:1128:VAL:HG13	1.94	0.49
1:A:1185:LYS:HZ2	1:A:1187:ASN:HD21	1.61	0.49
1:A:1020:LEU:O	1:A:1024:ARG:HG3	2.13	0.49
1:A:1279:ILE:HB	1:A:1284:ILE:HD11	1.95	0.49
1:A:1226:SER:O	1:A:1229:ARG:HG2	2.13	0.48
1:B:167:ARG:O	1:B:171:GLU:HB2	2.13	0.48
1:B:69:LEU:HD23	1:B:70:GLY:H	1.79	0.47
1:B:90:LEU:O	1:B:94:ARG:HG3	2.14	0.47
1:A:1082:ARG:HA	1:A:1085:ARG:HD2	1.95	0.47
1:A:1155:GLU:O	1:B:152:TRP:HA	2.14	0.47
1:A:1054:PRO:HB2	1:A:1056:PRO:HD2	1.97	0.47
1:B:2:LYS:HG3	1:B:64:ALA:HA	1.96	0.46
1:B:74:GLY:HA3	1:B:77:TRP:CZ3	2.51	0.45
1:A:1312:GLU:O	1:A:1315:VAL:HG22	2.17	0.45
1:A:1339:THR:O	1:A:1343:HIS:HD2	2.00	0.45
1:B:113:GLU:HB3	1:B:120:GLU:HG3	1.99	0.45
1:A:1153:ASN:HD21	1:B:191:VAL:H	1.63	0.45
1:A:1054:PRO:HG2	1:A:1057:THR:OG1	2.16	0.45
1:A:1275:SER:O	1:A:1276:ALA:C	2.60	0.45
1:A:1055:GLU:N	1:A:1056:PRO:HD2	2.32	0.44
1:B:174:ARG:HA	1:B:178:LYS:HD3	1.98	0.44
1:A:1005:VAL:C	1:A:1006:LEU:HD12	2.42	0.44
1:B:266:THR:HA	1:B:267:PRO:HD3	1.86	0.44
1:B:225:ARG:HB3	1:B:225:ARG:CZ	2.47	0.44
1:B:286:ASN:HD22	1:B:287:PRO:HD2	1.81	0.44
1:A:1271:PRO:HB2	1:A:1273:HIS:CD2	2.53	0.43
1:B:59:LYS:HD3	1:B:59:LYS:HA	1.81	0.43
1:A:1183:VAL:HG11	1:A:1243:LEU:HD11	2.00	0.43
1:B:20:LEU:HD23	1:B:23:LEU:HD12	2.01	0.43
1:B:159:LYS:N	1:B:160:PRO:HD2	2.32	0.43
1:B:211:LEU:HD22	1:B:212:GLU:H	1.83	0.43
1:B:216:VAL:HG13	1:B:217:ASP:N	2.34	0.43
1:A:1071:SER:HB3	1:A:1275:SER:HB3	2.00	0.42
1:A:1023:LEU:HD12	1:A:1023:LEU:HA	1.84	0.42
1:A:1159:LYS:O	1:A:1163:GLU:HG3	2.19	0.42
1:A:1247:ALA:HA	1:A:1250:LEU:HD13	2.01	0.42
1:B:248:SER:HB2	1:B:257:LEU:HD13	2.01	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1188:VAL:HG23	1:A:1189:LEU:HG	2.00	0.42
1:A:1081:PRO:O	1:A:1084:ILE:HG22	2.19	0.42
1:A:1196:ARG:HG3	1:A:1213:HIS:CE1	2.54	0.42
1:A:1082:ARG:HA	1:A:1085:ARG:CD	2.50	0.42
1:B:166:ALA:O	1:B:170:PHE:HD1	2.02	0.42
1:B:223:LEU:O	1:B:227:PRO:HB3	2.20	0.42
1:A:1030:GLU:HB2	1:A:1032:LEU:CD1	2.48	0.41
1:B:100:PHE:CE2	1:B:164:ARG:HD2	2.55	0.41
1:B:325:PRO:HD3	1:B:331:ALA:O	2.20	0.41
1:B:254:LEU:HA	1:B:257:LEU:HD12	2.03	0.41
1:B:322:THR:HB	1:B:331:ALA:HB2	2.02	0.41
1:A:1094:ARG:HA	1:A:1099:LEU:HD12	2.02	0.41
1:A:1153:ASN:HD22	1:B:191:VAL:CG2	2.25	0.41
1:B:134:LEU:O	1:B:237:ASN:ND2	2.54	0.41
1:B:141:GLY:HA3	1:B:154:THR:O	2.21	0.41
1:B:86:PRO:HB2	1:B:87:GLU:OE1	2.21	0.40
1:A:1183:VAL:HG22	1:A:1219:MET:HG3	2.03	0.40
1:A:1083:LYS:H	1:A:1083:LYS:CD	2.18	0.40
1:A:1185:LYS:HD2	1:A:1185:LYS:HA	1.93	0.40
1:A:1248:SER:O	1:A:1251:PRO:HD2	2.22	0.40
1:A:1306:GLU:H	1:A:1306:GLU:CD	2.29	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	343/345 (99%)	331 (96%)	10 (3%)	2 (1%)	21	38
1	B	343/345 (99%)	326 (95%)	17 (5%)	0	100	100
All	All	686/690 (99%)	657 (96%)	27 (4%)	2 (0%)	36	55

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1284	ILE
1	A	1082	ARG

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	267/267 (100%)	253 (95%)	14 (5%)	21	42
1	B	267/267 (100%)	257 (96%)	10 (4%)	30	57
All	All	534/534 (100%)	510 (96%)	24 (4%)	24	49

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1014	GLU
1	A	1023	LEU
1	A	1028	GLU
1	A	1032	LEU
1	A	1061	VAL
1	A	1069	LEU
1	A	1082	ARG
1	A	1155	GLU
1	A	1187	ASN
1	A	1214	GLN
1	A	1223	LEU
1	A	1241	ASP
1	A	1304	LEU
1	A	1311	VAL
1	B	26	LEU
1	B	90	LEU
1	B	93	LEU
1	B	95	LYS
1	B	155	GLU
1	B	211	LEU
1	B	214	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	250	LEU
1	B	307	LEU
1	B	344	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1153	ASN
1	A	1187	ASN
1	A	1214	GLN
1	A	1286	ASN
1	A	1343	HIS
1	B	102	ASN
1	B	153	ASN
1	B	286	ASN
1	B	343	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	345/345 (100%)	-0.81	0 100 100	12, 29, 55, 83	0
1	B	345/345 (100%)	-0.93	0 100 100	11, 27, 52, 68	0
All	All	690/690 (100%)	-0.87	0 100 100	11, 28, 53, 83	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.