



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 06:19 PM UTC

PDB ID : 4GD9 / pdb_00004gd9
Title : Circular Permuted Streptavidin N49/G48
Authors : Le Trong, I.; Chu, V.; Xing, Y.; Lybrand, T.P.; Stayton, P.S.; Stenkamp, R.E.
Deposited on : 2012-07-31
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4037 (1.50-1.50)
Clashscore	190562	4235 (1.50-1.50)
Ramachandran outliers	187476	4153 (1.50-1.50)
Sidechain outliers	187428	4150 (1.50-1.50)
RSRZ outliers	180081	4039 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	132	12% 83% 8% 9%
1	B	132	16% 94% 5% ..
1	C	132	17% 85% 8% 7%
1	D	132	17% 88% 5% 8%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4154 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Streptavidin.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	120	Total	C	N	O	0	4	0
			909	565	159	185			
1	B	131	Total	C	N	O	0	2	0
			964	597	168	199			
1	C	123	Total	C	N	O	0	5	0
			934	577	168	189			
1	D	122	Total	C	N	O	0	0	0
			911	568	158	185			

There are 20 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	48	MET	-	expression tag	UNP P22629
A	140	GLY	-	linker	UNP P22629
A	141	GLY	-	linker	UNP P22629
A	142	GLY	-	linker	UNP P22629
A	143	SER	-	linker	UNP P22629
B	48	MET	-	expression tag	UNP P22629
B	140	GLY	-	linker	UNP P22629
B	141	GLY	-	linker	UNP P22629
B	142	GLY	-	linker	UNP P22629
B	143	SER	-	linker	UNP P22629
C	48	MET	-	expression tag	UNP P22629
C	140	GLY	-	linker	UNP P22629
C	141	GLY	-	linker	UNP P22629
C	142	GLY	-	linker	UNP P22629
C	143	SER	-	linker	UNP P22629
D	48	MET	-	expression tag	UNP P22629
D	140	GLY	-	linker	UNP P22629
D	141	GLY	-	linker	UNP P22629
D	142	GLY	-	linker	UNP P22629
D	143	SER	-	linker	UNP P22629

- Molecule 2 is BIOTIN (CCD ID: BTN) (formula: $C_{10}H_{16}N_2O_3S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			16	10	2	3	1		
2	B	1	Total	C	N	O	S	0	0
			16	10	2	3	1		
2	C	1	Total	C	N	O	S	0	0
			16	10	2	3	1		
2	D	1	Total	C	N	O	S	0	0
			16	10	2	3	1		

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	C	1	Total O S 5 4 1	0	0
3	D	1	Total S 1 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	103	Total O 103 103	0	0
4	B	114	Total O 114 114	0	0
4	C	72	Total O 72 72	0	0
4	D	77	Total O 77 77	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Streptavidin




- Molecule 1: Streptavidin



- Molecule 1: Streptavidin



- Molecule 1: Streptavidin



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	54.72Å 85.89Å 100.14Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.94 – 1.50 36.94 – 1.50	Depositor EDS
% Data completeness (in resolution range)	100.0 (36.94-1.50) 87.9 (36.94-1.50)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.71 (at 1.50Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.201 , 0.230 0.208 , 0.236	Depositor DCC
R_{free} test set	3343 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	19.6	Xtriage
Anisotropy	0.203	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 36.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4154	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.95% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, BTN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.93	1/952 (0.1%)	0.84	0/1299
1	B	0.87	0/997	0.86	0/1362
1	C	0.91	2/985 (0.2%)	0.87	0/1342
1	D	0.82	0/932	0.83	0/1272
All	All	0.88	3/3866 (0.1%)	0.85	0/5275

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	87	HIS	CG-CD2	5.46	1.41	1.35
1	C	87	HIS	ND1-CE1	5.41	1.38	1.32
1	A	127	HIS	ND1-CE1	5.11	1.37	1.32

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	909	0	846	12	0
1	B	964	0	900	9	0
1	C	934	0	869	9	0
1	D	911	0	851	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	16	0	15	0	0
2	B	16	0	15	0	0
2	C	16	0	15	3	0
2	D	16	0	15	0	0
3	C	5	0	0	1	0
3	D	1	0	0	0	0
4	A	103	0	0	2	0
4	B	114	0	0	7	0
4	C	72	0	0	1	0
4	D	77	0	0	3	0
All	All	4154	0	3526	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:213:ALA:HB3	4:B:5199:HOH:O	1.72	0.88
1:D:213:ALA:HB3	4:D:5170:HOH:O	1.78	0.83
1:B:120:TRP:NE1	1:C:247:VAL:HG21	2.04	0.73
1:B:242[A]:THR:HG21	4:B:5112:HOH:O	1.89	0.72
1:C:248:GLY:HA3	2:C:5001:BTN:H92	1.74	0.68
1:A:132:LYS:O	1:A:133:VAL:HG22	1.93	0.68
1:B:242[A]:THR:HG23	4:B:5114:HOH:O	1.96	0.65
1:A:214:GLU:N	4:A:6094:HOH:O	2.28	0.64
1:A:110[A]:LEU:HD22	4:A:6030:HOH:O	1.96	0.64
1:A:132:LYS:O	1:A:133:VAL:CG2	2.47	0.63
1:A:110[A]:LEU:HD13	1:A:124:LEU:HD12	1.83	0.61
1:C:66:THR:HG23	3:C:5002:SO4:O1	2.01	0.60
1:D:105:ASN:OD1	1:D:129:THR:HG23	2.01	0.59
1:B:213:ALA:CB	4:B:5199:HOH:O	2.43	0.53
1:C:133:VAL:HG12	1:C:220:THR:O	2.09	0.53
1:A:80:LYS:HB2	4:B:5131:HOH:O	2.08	0.53
1:A:105:ASN:OD1	1:A:129:THR:HG23	2.09	0.53
1:C:248:GLY:HA3	2:C:5001:BTN:C9	2.41	0.50
2:C:5001:BTN:H102	4:C:5162:HOH:O	2.10	0.49
1:B:61:ASP:HA	4:B:5131:HOH:O	2.13	0.48
1:D:248:GLY:C	4:D:5176:HOH:O	2.56	0.47
1:A:80:LYS:CB	4:B:5131:HOH:O	2.63	0.46
1:C:91:THR:HB	1:D:91:THR:HB	1.98	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:49:ASN:N	4:D:5172:HOH:O	2.51	0.44
1:C:60:TYR:OH	1:C:214:GLU:HG2	2.17	0.44
1:A:132:LYS:C	1:A:133:VAL:CG2	2.91	0.44
1:B:120:TRP:NE1	1:C:247:VAL:CG2	2.76	0.42
1:A:91:THR:HB	1:B:91:THR:HB	2.04	0.40
1:B:120:TRP:HE1	1:C:247:VAL:HG21	1.81	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	120/132 (91%)	117 (98%)	3 (2%)	0	100	100
1	B	131/132 (99%)	128 (98%)	3 (2%)	0	100	100
1	C	124/132 (94%)	122 (98%)	2 (2%)	0	100	100
1	D	118/132 (89%)	116 (98%)	2 (2%)	0	100	100
All	All	493/528 (93%)	483 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	92/94 (98%)	89 (97%)	3 (3%)	33	8
1	B	95/94 (101%)	91 (96%)	4 (4%)	26	4
1	C	95/94 (101%)	92 (97%)	3 (3%)	34	8
1	D	89/94 (95%)	89 (100%)	0	100	100
All	All	371/376 (99%)	361 (97%)	10 (3%)	50	12

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	103	ARG
1	A	110[A]	LEU
1	A	110[B]	LEU
1	B	51	GLU
1	B	143	SER
1	B	242[A]	THR
1	B	242[B]	THR
1	C	52	SER
1	C	242[A]	THR
1	C	242[B]	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 1 is modelled with single atom - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	BTN	D	5001	-	17,17,17	0.91	1 (5%)	23,23,23	1.18	3 (13%)
2	BTN	A	5001	-	17,17,17	0.67	0	23,23,23	1.12	2 (8%)
3	SO4	C	5002	-	4,4,4	0.41	0	6,6,6	0.39	0
2	BTN	C	5001	-	17,17,17	0.72	0	23,23,23	1.11	1 (4%)
2	BTN	B	5001	-	17,17,17	0.76	0	23,23,23	0.94	1 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BTN	A	5001	-	-	0/7/28/28	0/2/2/2
2	BTN	C	5001	-	-	0/7/28/28	0/2/2/2
2	BTN	D	5001	-	-	0/7/28/28	0/2/2/2
2	BTN	B	5001	-	-	0/7/28/28	0/2/2/2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	5001	BTN	C2-S1	-2.38	1.78	1.82

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	5001	BTN	O12-C11-C10	2.57	122.13	114.00
2	D	5001	BTN	C8-C7-C2	-2.35	108.64	114.04
2	C	5001	BTN	C9-C10-C11	2.33	120.60	114.51
2	D	5001	BTN	O12-C11-C10	2.21	120.98	114.00
2	D	5001	BTN	C6-C5-N1	2.14	115.94	113.18
2	A	5001	BTN	O11-C11-C10	-2.12	116.37	123.09
2	B	5001	BTN	O12-C11-C10	2.12	120.69	114.00

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	5002	SO4	1	0
2	C	5001	BTN	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	120/132 (90%)	1.13	16 (13%)	7 7	14, 21, 44, 66	4 (3%)
1	B	131/132 (99%)	1.11	21 (16%)	5 5	14, 22, 43, 67	2 (1%)
1	C	123/132 (93%)	1.21	22 (17%)	3 3	17, 23, 45, 63	5 (4%)
1	D	122/132 (92%)	1.15	22 (18%)	3 3	17, 22, 51, 78	0
All	All	496/528 (93%)	1.15	81 (16%)	4 4	14, 22, 45, 78	11 (2%)

All (81) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	248	GLY	6.6
1	A	215	ALA	6.2
1	A	248	GLY	6.0
1	A	133	VAL	5.2
1	C	248	GLY	5.0
1	B	222	TYR	4.9
1	C	247	VAL	4.6
1	C	215	ALA	4.3
1	A	222	TYR	4.2
1	C	222	TYR	4.2
1	B	248	GLY	4.2
1	D	133	VAL	3.9
1	D	222	TYR	3.8
1	B	102	ALA	3.7
1	A	54	TYR	3.6
1	D	213	ALA	3.6
1	C	213	ALA	3.5
1	A	49	ASN	3.5
1	C	131	THR	3.3
1	B	142	GLY	3.3
1	D	216	GLY	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	103[A]	ARG	3.1
1	D	236	ASP	3.1
1	B	49	ASN	3.0
1	D	134	LYS	3.0
1	C	143	SER	3.0
1	D	129	THR	3.0
1	D	247	VAL	2.9
1	C	229	PHE	2.9
1	C	49	ASN	2.9
1	D	215	ALA	2.9
1	A	102	ALA	2.8
1	B	82	ASN	2.8
1	B	103	ARG	2.7
1	B	133	VAL	2.7
1	A	236	ASP	2.7
1	D	235	ALA	2.7
1	C	133	VAL	2.7
1	A	214	GLU	2.6
1	B	213	ALA	2.6
1	B	235	ALA	2.6
1	C	246	ALA	2.6
1	D	130	PHE	2.6
1	B	247	VAL	2.6
1	B	50	ALA	2.5
1	A	52[A]	SER	2.4
1	C	120	TRP	2.4
1	C	130	PHE	2.4
1	B	138	ALA	2.4
1	C	214	GLU	2.4
1	B	73	LEU	2.4
1	C	66	THR	2.4
1	A	116	GLU	2.4
1	D	97	VAL	2.3
1	B	53	ARG	2.3
1	D	83	TYR	2.3
1	B	141	GLY	2.3
1	A	230	ILE	2.3
1	B	104	ILE	2.3
1	D	100	ALA	2.3
1	B	143	SER	2.3
1	A	97	VAL	2.3
1	A	103	ARG	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	65	ALA	2.2
1	D	49	ASN	2.2
1	A	84	ARG	2.2
1	D	66	THR	2.2
1	C	82	ASN	2.2
1	A	246	ALA	2.2
1	B	246	ALA	2.2
1	C	92	TRP	2.2
1	C	239	LEU	2.1
1	D	50	ALA	2.1
1	C	230	ILE	2.1
1	C	216	GLY	2.1
1	D	54	TYR	2.1
1	B	217	ILE	2.1
1	C	84[A]	ARG	2.0
1	D	103	ARG	2.0
1	D	96	TYR	2.0
1	D	68	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	SO4	C	5002	5/5	0.88	0.17	16,17,20,22	5
2	BTN	C	5001	16/16	0.94	0.10	17,20,33,40	0
2	BTN	A	5001	16/16	0.96	0.08	17,19,23,24	0
2	BTN	D	5001	16/16	0.96	0.08	17,18,23,24	0
2	BTN	B	5001	16/16	0.96	0.08	16,20,23,25	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	SO4	D	5002	1/5	0.96	0.07	26,26,26,26	0

6.5 Other polymers [i](#)

There are no such residues in this entry.