



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 01:46 PM EDT

PDB ID : 4GDM / pdb_00004gdm
Title : Crystal Structure of E.coli MenH
Authors : Johnston, J.M.; Baker, E.N.; Guo, Z.; Jiang, M.
Deposited on : 2012-07-31
Resolution : 2.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1009 (2.76-2.76)
Clashscore	190562	1044 (2.76-2.76)
Ramachandran outliers	187476	1024 (2.76-2.76)
Sidechain outliers	187428	1024 (2.76-2.76)
RSRZ outliers	180081	1009 (2.76-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	268	 4% 83% 11% • 6%
1	B	268	 3% 80% 14% • •
1	C	268	 3% 82% 13% • •

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6043 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 2-succinyl-6-hydroxy-2,4-cyclohexadiene-1-carboxylate synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	253	Total	C	N	O	S	0	0	0
			1959	1236	356	358	9			
1	B	258	Total	C	N	O	S	0	0	0
			1990	1257	361	363	9			
1	C	259	Total	C	N	O	S	0	0	0
			2000	1262	365	364	9			

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-15	MET	-	expression tag	UNP P37355
A	-14	HIS	-	expression tag	UNP P37355
A	-13	HIS	-	expression tag	UNP P37355
A	-12	HIS	-	expression tag	UNP P37355
A	-11	HIS	-	expression tag	UNP P37355
A	-10	HIS	-	expression tag	UNP P37355
A	-9	HIS	-	expression tag	UNP P37355
A	-8	SER	-	expression tag	UNP P37355
A	-7	SER	-	expression tag	UNP P37355
A	-6	GLY	-	expression tag	UNP P37355
A	-5	LEU	-	expression tag	UNP P37355
A	-4	VAL	-	expression tag	UNP P37355
A	-3	PRO	-	expression tag	UNP P37355
A	-2	ARG	-	expression tag	UNP P37355
A	-1	GLY	-	expression tag	UNP P37355
A	0	SER	-	expression tag	UNP P37355
B	-15	MET	-	expression tag	UNP P37355
B	-14	HIS	-	expression tag	UNP P37355
B	-13	HIS	-	expression tag	UNP P37355
B	-12	HIS	-	expression tag	UNP P37355
B	-11	HIS	-	expression tag	UNP P37355
B	-10	HIS	-	expression tag	UNP P37355

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-9	HIS	-	expression tag	UNP P37355
B	-8	SER	-	expression tag	UNP P37355
B	-7	SER	-	expression tag	UNP P37355
B	-6	GLY	-	expression tag	UNP P37355
B	-5	LEU	-	expression tag	UNP P37355
B	-4	VAL	-	expression tag	UNP P37355
B	-3	PRO	-	expression tag	UNP P37355
B	-2	ARG	-	expression tag	UNP P37355
B	-1	GLY	-	expression tag	UNP P37355
B	0	SER	-	expression tag	UNP P37355
C	-15	MET	-	expression tag	UNP P37355
C	-14	HIS	-	expression tag	UNP P37355
C	-13	HIS	-	expression tag	UNP P37355
C	-12	HIS	-	expression tag	UNP P37355
C	-11	HIS	-	expression tag	UNP P37355
C	-10	HIS	-	expression tag	UNP P37355
C	-9	HIS	-	expression tag	UNP P37355
C	-8	SER	-	expression tag	UNP P37355
C	-7	SER	-	expression tag	UNP P37355
C	-6	GLY	-	expression tag	UNP P37355
C	-5	LEU	-	expression tag	UNP P37355
C	-4	VAL	-	expression tag	UNP P37355
C	-3	PRO	-	expression tag	UNP P37355
C	-2	ARG	-	expression tag	UNP P37355
C	-1	GLY	-	expression tag	UNP P37355
C	0	SER	-	expression tag	UNP P37355

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Cl	0	0
			1	1		
3	B	1	Total	Cl	0	0
			1	1		
3	C	1	Total	Cl	0	0
			1	1		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	29	Total	O	0	0
			29	29		
5	B	21	Total	O	0	0
			21	21		
5	C	15	Total	O	0	0
			15	15		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 2-succinyl-6-hydroxy-2,4-cyclohexadiene-1-carboxylate synthase



- Molecule 1: 2-succinyl-6-hydroxy-2,4-cyclohexadiene-1-carboxylate synthase



- Molecule 1: 2-succinyl-6-hydroxy-2,4-cyclohexadiene-1-carboxylate synthase



4 Data and refinement statistics

Property	Value	Source
Space group	I 41 2 2	Depositor
Cell constants a, b, c, α , β , γ	116.66Å 116.66Å 476.95Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.77 – 2.75 45.77 – 2.75	Depositor EDS
% Data completeness (in resolution range)	99.9 (45.77-2.75) 99.9 (45.77-2.75)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.00 (at 2.77Å)	Xtriage
Refinement program	BUSTER-TNT, BUSTER 2.10.0	Depositor
R, R_{free}	0.192 , 0.223 0.194 , 0.225	Depositor DCC
R_{free} test set	2182 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	53.6	Xtriage
Anisotropy	0.167	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 55.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6043	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, CL, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.87	0/2005	1.35	12/2720 (0.4%)
1	B	0.90	0/2037	1.39	10/2765 (0.4%)
1	C	0.85	0/2047	1.34	8/2777 (0.3%)
All	All	0.87	0/6089	1.36	30/8262 (0.4%)

There are no bond length outliers.

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	63	ASP	CA-CB-CG	6.92	119.52	112.60
1	C	39	ASP	CA-CB-CG	6.52	119.12	112.60
1	C	214	ARG	CA-C-N	6.42	128.89	120.28
1	C	214	ARG	C-N-CA	6.42	128.89	120.28
1	A	37	PHE	CA-C-N	6.16	128.82	120.38
1	A	37	PHE	C-N-CA	6.16	128.82	120.38
1	B	67	LEU	CA-C-N	5.80	127.98	120.44
1	B	67	LEU	C-N-CA	5.80	127.98	120.44
1	A	223	ASP	CA-CB-CG	5.76	118.36	112.60
1	B	157	ASN	CA-C-N	5.74	127.90	120.44
1	B	157	ASN	C-N-CA	5.74	127.90	120.44
1	A	39	ASP	CA-CB-CG	5.73	118.33	112.60
1	A	162	ARG	CA-C-N	5.73	127.88	120.44
1	A	162	ARG	C-N-CA	5.73	127.88	120.44
1	A	91	VAL	N-CA-CB	5.69	119.11	110.58
1	B	213	PHE	CA-CB-CG	-5.59	108.21	113.80
1	C	91	VAL	N-CA-CB	5.47	118.79	110.58
1	C	178	ALA	CA-C-N	5.45	127.58	120.28
1	C	178	ALA	C-N-CA	5.45	127.58	120.28
1	B	223	ASP	CA-CB-CG	5.44	118.04	112.60
1	B	73	VAL	N-CA-CB	5.38	117.86	110.54

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	101	ALA	N-CA-C	5.34	117.54	111.02
1	A	213	PHE	CA-CB-CG	-5.26	108.54	113.80
1	C	-3	PRO	CA-C-N	5.24	127.30	120.28
1	C	-3	PRO	C-N-CA	5.24	127.30	120.28
1	A	159	ASP	CA-C-N	5.21	127.26	120.28
1	A	159	ASP	C-N-CA	5.21	127.26	120.28
1	B	210	ASP	CA-CB-CG	5.19	117.79	112.60
1	A	32	GLU	CA-C-N	5.10	126.94	120.72
1	A	32	GLU	C-N-CA	5.10	126.94	120.72

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1959	0	1901	8	0
1	B	1990	0	1933	14	0
1	C	2000	0	1947	13	0
2	A	5	0	0	0	0
2	B	10	0	0	0	0
2	C	5	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
4	B	6	0	8	1	0
5	A	29	0	0	0	0
5	B	21	0	0	0	0
5	C	15	0	0	1	0
All	All	6043	0	5789	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:94:MET:HE1	1:A:188:GLN:HG3	1.57	0.84
1:C:106:VAL:HG13	1:C:202:PHE:HD1	1.63	0.63
1:A:205:LEU:HD23	1:A:225:HIS:HB2	1.85	0.59
1:C:235:HIS:HA	1:C:242:VAL:HG21	1.87	0.57
1:B:132:VAL:HG21	1:B:181:GLU:HG2	1.87	0.56
1:A:132:VAL:HG21	1:A:181:GLU:HG2	1.89	0.55
1:C:69:ARG:O	1:C:73:VAL:HG23	2.11	0.51
1:A:94:MET:CE	1:A:188:GLN:HG3	2.37	0.51
1:C:132:VAL:HG13	1:C:177:ALA:HB1	1.95	0.49
1:B:184:SER:O	1:B:187:VAL:HG12	2.13	0.48
1:C:20:LEU:HD13	1:C:91:VAL:HG22	1.96	0.46
1:B:29:GLU:OE1	1:B:235:HIS:NE2	2.34	0.46
1:A:184:SER:O	1:A:187:VAL:HG12	2.15	0.46
1:B:61:PHE:CG	1:B:94:MET:HG2	2.51	0.46
1:C:235:HIS:CA	1:C:242:VAL:HG21	2.47	0.45
1:B:145:ALA:HA	1:B:165:VAL:HG11	1.99	0.45
1:C:17:LEU:HB3	1:C:19:PHE:CE2	2.51	0.45
1:B:161:ARG:O	1:B:165:VAL:HG13	2.17	0.45
1:B:109:GLU:OE1	1:B:235:HIS:ND1	2.50	0.44
1:B:198:ARG:NH1	4:B:304:GOL:H11	2.33	0.44
1:B:106:VAL:HG12	1:B:108:VAL:HG23	2.00	0.44
1:C:86:SER:HB3	5:C:409:HOH:O	2.17	0.44
1:C:16:TRP:CD2	1:C:41:SER:HB3	2.54	0.43
1:B:-2:ARG:HA	1:B:55:ALA:HB1	2.00	0.43
1:A:161:ARG:O	1:A:165:VAL:HG23	2.19	0.42
1:B:93:MET:HG3	1:B:108:VAL:HG21	2.00	0.42
1:B:132:VAL:HG13	1:B:177:ALA:HB1	2.02	0.42
1:C:2:ILE:HG23	1:C:71:THR:HG23	2.02	0.42
1:A:208:GLU:CD	1:C:198:ARG:HH12	2.27	0.42
1:B:78:LEU:O	1:B:102:GLY:HA3	2.21	0.41
1:A:31:GLN:O	1:A:35:GLU:HG2	2.21	0.41
1:C:95:ALA:HB1	1:C:100:LEU:HD11	2.02	0.41
1:C:144:PHE:HA	1:C:147:TRP:HB3	2.04	0.40
1:B:86:SER:OG	1:B:232:HIS:NE2	2.53	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	251/268 (94%)	241 (96%)	9 (4%)	1 (0%)	30	50
1	B	256/268 (96%)	247 (96%)	9 (4%)	0	100	100
1	C	257/268 (96%)	246 (96%)	11 (4%)	0	100	100
All	All	764/804 (95%)	734 (96%)	29 (4%)	1 (0%)	48	71

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	86	SER

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	197/210 (94%)	186 (94%)	11 (6%)	19	36
1	B	200/210 (95%)	183 (92%)	17 (8%)	10	18
1	C	201/210 (96%)	185 (92%)	16 (8%)	11	20
All	All	598/630 (95%)	554 (93%)	44 (7%)	13	24

All (44) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	17	LEU
1	A	20	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	76	ASN
1	A	79	ASP
1	A	91	VAL
1	A	129	ARG
1	A	149	GLN
1	A	161	ARG
1	A	176	LEU
1	A	196	SER
1	A	199	THR
1	B	2	ILE
1	B	14	LEU
1	B	20	LEU
1	B	57	SER
1	B	62	ASP
1	B	67	LEU
1	B	73	VAL
1	B	78	LEU
1	B	79	ASP
1	B	100	LEU
1	B	119	GLU
1	B	149	GLN
1	B	158	ASP
1	B	181	GLU
1	B	199	THR
1	B	223	ASP
1	B	237	GLU
1	C	-4	VAL
1	C	1	MET
1	C	17	LEU
1	C	39	ASP
1	C	66	ASP
1	C	68	LEU
1	C	86	SER
1	C	91	VAL
1	C	125	GLN
1	C	141	THR
1	C	155	SER
1	C	175	THR
1	C	176	LEU
1	C	181	GLU
1	C	187	VAL
1	C	199	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	125	GLN
1	A	130	GLN
1	A	171	ASN
1	A	248	GLN
1	B	170	ASN
1	C	130	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 3 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GOL	B	304	-	5,5,5	0.21	0	5,5,5	0.80	0
2	SO4	A	301	-	4,4,4	0.34	0	6,6,6	0.21	0
2	SO4	B	302	-	4,4,4	0.32	0	6,6,6	0.27	0
2	SO4	C	301	-	4,4,4	0.29	0	6,6,6	0.21	0
2	SO4	B	301	-	4,4,4	0.21	0	6,6,6	0.41	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	B	304	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	304	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	253/268 (94%)	-0.03	10 (3%) 42 40	30, 49, 80, 97	0
1	B	258/268 (96%)	0.09	7 (2%) 56 54	30, 48, 78, 110	0
1	C	259/268 (96%)	0.24	7 (2%) 56 54	36, 55, 81, 95	0
All	All	770/804 (95%)	0.10	24 (3%) 51 49	30, 52, 80, 110	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	-5	LEU	7.4
1	B	-3	PRO	4.4
1	B	-4	VAL	4.2
1	C	252	PHE	4.2
1	B	-2	ARG	3.7
1	B	1	MET	3.6
1	B	-1	GLY	3.3
1	C	1	MET	3.0
1	B	158	ASP	2.7
1	C	-2	ARG	2.6
1	A	0	SER	2.6
1	A	159	ASP	2.6
1	C	11	LYS	2.6
1	A	199	THR	2.5
1	A	119	GLU	2.3
1	A	118	ALA	2.3
1	C	118	ALA	2.2
1	A	141	THR	2.1
1	A	123	GLU	2.1
1	A	229	ARG	2.1
1	C	221	ALA	2.1
1	A	252	PHE	2.1
1	A	196	SER	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	251	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	GOL	B	304	6/6	0.70	0.22	57,60,62,62	0
2	SO4	B	302	5/5	0.89	0.22	98,101,101,104	0
2	SO4	B	301	5/5	0.93	0.11	71,72,73,74	0
2	SO4	C	301	5/5	0.94	0.08	67,70,74,75	0
2	SO4	A	301	5/5	0.95	0.10	70,72,74,74	0
3	CL	B	303	1/1	0.96	0.09	48,48,48,48	0
3	CL	A	302	1/1	0.96	0.15	61,61,61,61	0
3	CL	C	302	1/1	0.98	0.06	49,49,49,49	0

6.5 Other polymers [i](#)

There are no such residues in this entry.