



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 18, 2026 – 12:36 PM UTC

PDB ID : 2GFE / pdb_00002gfe
Title : Crystal structure of the GluR2 A476E S673D Ligand Binding Core Mutant at 1.54 Angstroms Resolution
Authors : Mayer, M.L.
Deposited on : 2006-03-21
Resolution : 1.54 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.54 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1003 (1.54-1.54)
Clashscore	190562	1025 (1.54-1.54)
Ramachandran outliers	187476	1007 (1.54-1.54)
Sidechain outliers	187428	1007 (1.54-1.54)
RSRZ outliers	180081	1002 (1.54-1.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	262	6% 84% 15% ..
1	B	262	8% 87% 12% .
1	C	262	19% 82% 16% ..

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6989 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutamate receptor 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	260	Total	C	N	O	S	28	9	0
			2104	1336	348	404	16			
1	B	260	Total	C	N	O	S	27	2	0
			2051	1304	341	391	15			
1	C	260	Total	C	N	O	S	37	2	0
			2053	1307	341	390	15			


There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	cloning artifact	UNP P19491
A	2	ALA	-	cloning artifact	UNP P19491
A	66	GLU	ALA	engineered mutation	UNP P19491
A	118	GLY	-	linker	UNP P19491
A	119	THR	-	linker	UNP P19491
A	140	ASP	SER	engineered mutation	UNP P19491
B	1	GLY	-	cloning artifact	UNP P19491
B	2	ALA	-	cloning artifact	UNP P19491
B	66	GLU	ALA	engineered mutation	UNP P19491
B	118	GLY	-	linker	UNP P19491
B	119	THR	-	linker	UNP P19491
B	140	ASP	SER	engineered mutation	UNP P19491
C	1	GLY	-	cloning artifact	UNP P19491
C	2	ALA	-	cloning artifact	UNP P19491
C	66	GLU	ALA	engineered mutation	UNP P19491
C	118	GLY	-	linker	UNP P19491
C	119	THR	-	linker	UNP P19491
C	140	ASP	SER	engineered mutation	UNP P19491

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total 1	Zn 1	0	0
2	B	2	Total 2	Zn 2	0	0
2	C	2	Total 2	Zn 2	0	0

- Molecule 3 is GLUTAMIC ACID (CCD ID: GLU) (formula: C₅H₉NO₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total 10	C 5	N 1	O 4	0	0
3	B	1	Total 10	C 5	N 1	O 4	0	0
3	C	1	Total 10	C 5	N 1	O 4	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	253	Total 253	O 253	0	0
4	B	299	Total 299	O 299	0	0
4	C	194	Total 194	O 194	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Glutamate receptor 2



• Molecule 1: Glutamate receptor 2



• Molecule 1: Glutamate receptor 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	114.32Å 163.30Å 47.56Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.42 – 1.54 39.42 – 1.54	Depositor EDS
% Data completeness (in resolution range)	98.0 (39.42-1.54) 98.2 (39.42-1.54)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.61 (at 1.54Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.212 , 0.227 0.219 , 0.233	Depositor DCC
R_{free} test set	6532 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å ²)	14.6	Xtriage
Anisotropy	0.017	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 40.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6989	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 13.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.31	0/2140	0.80	3/2877 (0.1%)
1	B	0.32	0/2087	0.81	4/2806 (0.1%)
1	C	0.30	0/2089	0.78	2/2809 (0.1%)
All	All	0.31	0/6316	0.80	9/8492 (0.1%)

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	116	LYS	N-CA-C	-6.74	99.55	110.20
1	B	116	LYS	N-CA-C	-6.45	100.00	110.20
1	A	104	LYS	N-CA-C	-5.44	102.12	110.01
1	B	104	LYS	N-CA-C	-5.38	102.21	110.01
1	A	103	SER	N-CA-C	-5.38	103.59	110.53
1	B	257	ASP	N-CA-C	-5.32	106.05	112.54
1	C	116	LYS	N-CA-C	-5.30	101.34	109.76
1	C	57	GLY	N-CA-C	5.23	118.96	112.64
1	B	184	SER	N-CA-C	5.08	119.12	113.02

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2104	0	2116	24	0
1	B	2051	0	2073	22	0
1	C	2053	0	2079	34	0
2	A	1	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
3	A	10	0	5	0	0
3	B	10	0	5	0	0
3	C	10	0	5	0	0
4	A	253	0	0	6	0
4	B	299	0	0	9	0
4	C	194	0	0	3	0
All	All	6989	0	6283	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (79) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:203:ARG:HH21	1:C:259:GLY:HA2	1.11	1.12
1:C:22:ASN:HB2	1:C:25:MET:HE3	1.33	1.06
1:C:203:ARG:NH2	1:C:259:GLY:HA2	1.82	0.94
1:B:183:LYS:HG3	4:B:1297:HOH:O	1.87	0.74
1:A:140[B]:ASP:HB3	4:A:1189:HOH:O	1.90	0.71
1:A:4:LYS:HE3	4:A:1197:HOH:O	1.92	0.70
1:C:39:LEU:HG	1:C:43[B]:ILE:HD11	1.73	0.70
1:C:194:SER:O	1:C:198:GLU:HG2	1.91	0.69
1:B:81:GLY:HA2	4:B:1228:HOH:O	1.93	0.67
1:C:88:ALA:HB1	1:C:89:PRO:HD2	1.76	0.67
1:A:42:GLU:HG3	1:A:246:LEU:HD21	1.77	0.66
1:A:81:GLY:HA2	4:A:1181:HOH:O	1.94	0.66
1:B:140:ASP:HB3	4:B:1198:HOH:O	1.96	0.66
1:A:43[B]:ILE:HD13	1:A:222[B]:ILE:HD13	1.80	0.62
1:C:166:GLU:HA	1:C:167:PRO:C	2.24	0.62
1:A:39:LEU:HG	1:A:43[B]:ILE:HD11	1.82	0.62
1:A:22:ASN:HB2	1:A:25:MET:HE3	1.82	0.61
1:A:210:LYS:HE2	4:A:1190:HOH:O	2.03	0.59
1:B:253:LYS:O	1:B:258:LYS:HG2	2.03	0.59
1:C:43[B]:ILE:HD13	1:C:222:ILE:HD13	1.85	0.59
1:A:52:LYS:NZ	1:A:52:LYS:HB2	2.19	0.58
1:C:22:ASN:CB	1:C:25:MET:HE3	2.22	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:88:ALA:HB1	1:A:89:PRO:HD2	1.87	0.56
1:B:210:LYS:HE2	4:B:1211:HOH:O	2.05	0.56
1:B:88:ALA:HB1	1:B:89:PRO:HD2	1.87	0.55
1:C:137:THR:HG22	1:C:191:LEU:HB2	1.89	0.54
1:A:65:ASP:OD1	1:A:67:ASP:HB2	2.08	0.54
1:C:45:LYS:NZ	1:C:45:LYS:HB2	2.22	0.54
1:C:81:GLY:HA2	4:C:1144:HOH:O	2.08	0.53
1:B:184:SER:HB2	4:B:1079:HOH:O	2.08	0.53
1:C:149:ARG:NH1	4:C:1013:HOH:O	2.41	0.53
1:B:50:LYS:HB3	1:B:50:LYS:NZ	2.24	0.53
1:B:23:HIS:CD2	1:B:26:LEU:HD12	2.45	0.52
1:C:242:ASN:HB2	1:C:247:LEU:HD12	1.93	0.51
1:B:107:MET:HE2	1:B:251:LYS:HD3	1.93	0.50
1:C:225:PRO:HG2	1:C:228:SER:HB2	1.93	0.50
1:A:204:LYS:HG3	1:A:260:GLU:HB3	1.94	0.50
1:C:240:LYS:O	1:C:244:GLN:HG3	2.10	0.50
1:B:183:LYS:HG2	4:B:1291:HOH:O	2.12	0.50
1:C:159:TRP:O	1:C:163:ARG:HG2	2.11	0.50
1:C:246:LEU:C	1:C:246:LEU:HD13	2.38	0.49
1:A:145:GLU:HG2	4:A:1158:HOH:O	2.13	0.48
1:A:137:THR:HG22	1:A:191:LEU:HB2	1.95	0.48
1:A:39:LEU:HG	1:A:43[B]:ILE:CD1	2.45	0.46
1:A:251:LYS:O	1:A:255:TRP:HB2	2.16	0.45
1:C:64:ARG:CZ	1:C:69:LYS:HD2	2.46	0.45
1:B:154:VAL:HG23	4:B:1107:HOH:O	2.16	0.45
1:C:39:LEU:HG	1:C:43[B]:ILE:CD1	2.45	0.45
1:B:204:LYS:HG3	1:B:260:GLU:HB3	1.99	0.44
1:A:140[A]:ASP:HB3	4:A:1189:HOH:O	2.17	0.44
1:C:15:PRO:HB3	1:C:199:TYR:CE1	2.53	0.44
1:C:109:LEU:HD22	1:C:109:LEU:C	2.43	0.43
1:B:183:LYS:CG	4:B:1297:HOH:O	2.55	0.43
1:C:39:LEU:O	1:C:43[B]:ILE:HG13	2.18	0.43
1:A:241:LEU:HD22	1:A:246:LEU:HD23	2.00	0.43
1:A:179:ALA:O	1:A:183:LYS:HG2	2.19	0.42
1:A:125:GLU:CG	1:A:129:LYS:HE3	2.50	0.42
1:C:45:LYS:HB2	1:C:45:LYS:HZ2	1.84	0.42
1:C:135:TYR:HA	1:C:189:ALA:O	2.19	0.42
1:A:109:LEU:HB3	1:A:195:THR:HG23	2.02	0.42
1:B:26:LEU:O	1:B:31:ARG:HD3	2.20	0.42
1:C:27:GLU:HG3	4:C:1040:HOH:O	2.20	0.42
1:B:199:TYR:O	1:B:203:ARG:HG2	2.19	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:179:ALA:O	1:C:183:LYS:HG2	2.19	0.42
1:A:244:GLN:OE1	1:C:151:LYS:HD3	2.20	0.41
1:B:137:THR:HG22	1:B:191:LEU:HB2	2.02	0.41
1:C:136:GLY:HA3	1:C:169:VAL:O	2.20	0.41
1:C:215:LEU:HD12	1:C:215:LEU:N	2.35	0.41
1:B:100:ILE:HD12	1:B:223:ALA:HB1	2.02	0.41
1:A:41:ALA:O	1:A:45:LYS:HG3	2.21	0.41
1:B:4:LYS:HD3	4:B:1203:HOH:O	2.19	0.41
1:A:136:GLY:HA3	1:A:169:VAL:O	2.21	0.41
1:C:4:LYS:H	1:C:4:LYS:HD3	1.85	0.41
1:C:199:TYR:CE1	1:C:203:ARG:NH1	2.89	0.41
1:C:109:LEU:HD13	1:C:109:LEU:N	2.35	0.40
1:B:3:ASN:C	1:B:4:LYS:HD2	2.46	0.40
1:B:246:LEU:HD13	1:B:246:LEU:C	2.46	0.40
1:B:3:ASN:O	1:B:4:LYS:HD2	2.21	0.40
1:C:22:ASN:O	1:C:25:MET:HG2	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	267/262 (102%)	261 (98%)	3 (1%)	3 (1%)	11	2
1	B	260/262 (99%)	257 (99%)	3 (1%)	0	100	100
1	C	260/262 (99%)	256 (98%)	4 (2%)	0	100	100
All	All	787/786 (100%)	774 (98%)	10 (1%)	3 (0%)	36	12

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	139[A]	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	139[B]	ASP
1	A	67	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	227/219 (104%)	223 (98%)	4 (2%)	51	21
1	B	221/219 (101%)	219 (99%)	2 (1%)	70	46
1	C	221/219 (101%)	218 (99%)	3 (1%)	59	29
All	All	669/657 (102%)	660 (99%)	9 (1%)	59	31

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	15	PRO
1	A	27	GLU
1	A	52	LYS
1	A	109	LEU
1	B	97	GLU
1	B	109	LEU
1	C	21	LYS
1	C	45	LYS
1	C	109	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	242	ASN
1	C	244	GLN
1	C	252	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 8 ligands modelled in this entry, 5 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GLU	C	998	-	8,9,9	1.01	0	8,11,11	0.78	0
3	GLU	A	997	-	8,9,9	1.02	0	8,11,11	0.81	0
3	GLU	B	999	-	8,9,9	1.04	0	8,11,11	0.79	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GLU	C	998	-	-	1/9/9/9	-
3	GLU	A	997	-	-	1/9/9/9	-
3	GLU	B	999	-	-	1/9/9/9	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	997	GLU	O-C-CA-N
3	B	999	GLU	O-C-CA-N
3	C	998	GLU	O-C-CA-CB

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	260/262 (99%)	0.36	17 (6%)	25 31	5, 15, 27, 43	17 (6%)
1	B	260/262 (99%)	0.29	20 (7%)	19 24	5, 14, 28, 36	11 (4%)
1	C	260/262 (99%)	1.07	49 (18%)	3 4	7, 19, 38, 44	12 (4%)
All	All	780/786 (99%)	0.57	86 (11%)	10 14	5, 15, 32, 44	40 (5%)

All (86) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2	ALA	6.8
1	C	159	TRP	5.9
1	C	154	VAL	5.4
1	C	153	ALA	5.1
1	C	256	TYR	4.9
1	C	149	ARG	4.9
1	C	203	ARG	4.9
1	C	131	THR	4.8
1	A	66	GLU	4.8
1	C	2	ALA	4.7
1	B	3	ASN	4.7
1	B	2	ALA	4.7
1	C	150	SER	4.6
1	B	26	LEU	4.6
1	C	155	PHE	4.4
1	C	160	THR	4.2
1	C	152	ILE	4.1
1	C	165	ALA	4.0
1	A	68	THR	4.0
1	A	27	GLU	4.0
1	C	128	SER	3.9
1	C	147	PHE	3.7
1	B	27	GLU	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	140[A]	ASP	3.6
1	B	256	TYR	3.5
1	C	129	LYS	3.5
1	B	140	ASP	3.4
1	A	4	LYS	3.4
1	C	259	GLY	3.4
1	C	227	GLY	3.3
1	B	67	ASP	3.3
1	C	148	ARG	3.3
1	B	66	GLU	3.2
1	C	166	GLU	3.2
1	C	184	SER	3.2
1	B	68	THR	3.1
1	C	164	SER	3.0
1	C	161	TYR	3.0
1	C	167	PRO	2.9
1	B	227	GLY	2.9
1	B	258	LYS	2.9
1	C	258	LYS	2.9
1	B	229	SER	2.8
1	C	27	GLU	2.8
1	C	257	ASP	2.7
1	A	227	GLY	2.7
1	C	132	GLU	2.7
1	B	4	LYS	2.6
1	C	133	ILE	2.6
1	C	146	PHE	2.6
1	C	127	LEU	2.6
1	C	156	ASP	2.6
1	C	229	SER	2.6
1	C	151	LYS	2.5
1	A	3	ASN	2.5
1	B	257	ASP	2.5
1	C	246	LEU	2.5
1	C	163	ARG	2.5
1	B	152	ILE	2.5
1	C	4	LYS	2.4
1	B	25	MET	2.4
1	A	69	LYS	2.4
1	C	145	GLU	2.4
1	C	199	TYR	2.4
1	A	64	ARG	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	261	CYS	2.4
1	C	24	GLU	2.4
1	A	67	ASP	2.3
1	A	25	MET	2.3
1	C	25	MET	2.3
1	B	145	GLU	2.3
1	C	228	SER	2.3
1	C	158	MET	2.2
1	A	148	ARG	2.2
1	A	152	ILE	2.2
1	B	153	ALA	2.2
1	C	169	VAL	2.2
1	A	139[A]	ASP	2.1
1	B	65	ASP	2.1
1	C	252	ASN	2.1
1	A	163	ARG	2.1
1	C	250	LEU	2.1
1	C	126	ASP	2.1
1	A	226	LYS	2.0
1	C	170	PHE	2.0
1	B	261	CYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	ZN	C	873	1/1	0.95	0.21	49,49,49,49	0
2	ZN	C	871	1/1	0.96	0.07	22,22,22,22	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	GLU	C	998	10/10	0.96	0.06	10,12,14,14	0
2	ZN	B	870	1/1	0.98	0.04	19,19,19,19	0
3	GLU	A	997	10/10	0.98	0.04	9,10,12,13	0
3	GLU	B	999	10/10	0.98	0.04	8,8,10,10	0
2	ZN	A	869	1/1	0.98	0.03	14,14,14,14	0
2	ZN	B	872	1/1	0.99	0.03	15,15,15,15	0

6.5 Other polymers [i](#)

There are no such residues in this entry.