



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 06:41 PM UTC

PDB ID : 1GGI / pdb_00001ggi
Title : CRYSTAL STRUCTURE OF AN HIV-1 NEUTRALIZING ANTIBODY 50.1
IN COMPLEX WITH ITS V3 LOOP PEPTIDE ANTIGEN
Authors : Stanfield, R.L.; Rini, J.M.; Wilson, I.A.
Deposited on : 1993-04-02
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	L	218	38% 40% 17% ..
1	M	218	41% 43% 13% ..
2	H	222	37% 42% 13% 5% .
2	J	222	38% 42% 15% ..
3	P	16	19% 25% 12% 44%
3	Q	16	19% 25% 12% 44%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6712 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called IGG2A 50.1 FAB (LIGHT CHAIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	215	Total	C	N	O	S	0	0	0
			1662	1031	283	343	5			
1	M	215	Total	C	N	O	S	0	0	0
			1662	1031	283	343	5			

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
L	4	LEU	MET	conflict	EMBL AJ131289
L	7	SER	THR	conflict	EMBL AJ131289
L	9	GLY	ALA	conflict	EMBL AJ131289
L	27A	SER	ASN	conflict	EMBL AJ131289
L	27C	ASP	ARG	conflict	EMBL AJ131289
L	28	ASP	TYR	conflict	EMBL AJ131289
L	33	LEU	MET	conflict	EMBL AJ131289
L	40	PRO	ALA	conflict	EMBL AJ131289
L	51	SER	ALA	conflict	EMBL AJ131289
L	55	ILE	GLU	conflict	EMBL AJ131289
L	60	ASP	ALA	conflict	EMBL AJ131289
L	87	TYR	PHE	conflict	EMBL AJ131289
L	90	GLN	ARG	conflict	EMBL AJ131289
L	94	ASP	VAL	conflict	EMBL AJ131289
L	96	LEU	TRP	conflict	EMBL AJ131289
L	100	ALA	GLY	conflict	EMBL AJ131289
M	4	LEU	MET	conflict	EMBL AJ131289
M	7	SER	THR	conflict	EMBL AJ131289
M	9	GLY	ALA	conflict	EMBL AJ131289
M	27A	SER	ASN	conflict	EMBL AJ131289
M	27C	ASP	ARG	conflict	EMBL AJ131289
M	28	ASP	TYR	conflict	EMBL AJ131289
M	33	LEU	MET	conflict	EMBL AJ131289
M	40	PRO	ALA	conflict	EMBL AJ131289
M	51	SER	ALA	conflict	EMBL AJ131289

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
M	55	ILE	GLU	conflict	EMBL AJ131289
M	60	ASP	ALA	conflict	EMBL AJ131289
M	87	TYR	PHE	conflict	EMBL AJ131289
M	90	GLN	ARG	conflict	EMBL AJ131289
M	94	ASP	VAL	conflict	EMBL AJ131289
M	96	LEU	TRP	conflict	EMBL AJ131289
M	100	ALA	GLY	conflict	EMBL AJ131289

- Molecule 2 is a protein called IGG2A 50.1 FAB (HEAVY CHAIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	215	Total	C	N	O	S	0	0	0
			1627	1032	266	323	6			
2	J	215	Total	C	N	O	S	0	0	0
			1627	1032	266	323	6			

- Molecule 3 is a protein called HIV-1 V3 LOOP PEPTIDE ANTIGEN.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	P	9	Total	C	N	O	S	0	0	0
			67	42	15	9	1			
3	Q	9	Total	C	N	O	S	0	0	0
			67	42	15	9	1			


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


• Molecule 1: IGG2A 50.1 FAB (LIGHT CHAIN)






• Molecule 2: IGG2A 50.1 FAB (HEAVY CHAIN)

Chain J: 38% 42% 15% . .




• Molecule 3: HIV-1 V3 LOOP PEPTIDE ANTIGEN

Chain P: 19% 25% 12% 44%



• Molecule 3: HIV-1 V3 LOOP PEPTIDE ANTIGEN

Chain Q: 19% 25% 12% 44%



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	130.33Å 52.57Å 82.04Å 90.00° 97.50° 90.00°	Depositor
Resolution (Å)	(Not available) – 2.80	Depositor
% Data completeness (in resolution range)	(Not available) ((Not available)-2.80)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR	Depositor
R, R_{free}	0.188 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	6712	wwPDB-VP
Average B, all atoms (Å ²)	12.0	wwPDB-VP

5 Model quality ⓘ

5.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	L	1.22	6/1698 (0.4%)	2.31	94/2309 (4.1%)
1	M	1.25	5/1698 (0.3%)	2.41	103/2309 (4.5%)
2	H	1.27	10/1669 (0.6%)	2.36	90/2281 (3.9%)
2	J	1.23	8/1669 (0.5%)	2.36	100/2281 (4.4%)
3	P	1.76	2/68 (2.9%)	2.71	6/89 (6.7%)
3	Q	1.66	1/68 (1.5%)	2.61	6/89 (6.7%)
All	All	1.25	32/6870 (0.5%)	2.37	399/9358 (4.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	L	0	2
1	M	0	1
All	All	0	3

The worst 5 of 32 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	M	34	HIS	CD2-NE2	-8.86	1.28	1.37
2	H	50	HIS	CD2-NE2	-8.45	1.28	1.37
2	H	50	HIS	CG-ND1	-7.39	1.30	1.38
1	L	34	HIS	CD2-NE2	-7.31	1.29	1.37
2	J	212	HIS	CD2-NE2	-7.04	1.30	1.37

The worst 5 of 399 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	56	ASP	CA-CB-CG	16.80	129.40	112.60
1	L	28	ASP	CA-CB-CG	-16.39	96.21	112.60
2	H	94	GLN	OE1-CD-NE2	-13.78	108.82	122.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	J	178	LEU	N-CA-C	13.09	130.92	108.02
1	M	167	ASP	CA-CB-CG	11.40	124.00	112.60

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	L	192	TYR	Sidechain
1	L	203	SER	Peptide
1	M	186	TYR	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1662	0	1592	51	0
1	M	1662	0	1592	40	0
2	H	1627	0	1601	53	0
2	J	1627	0	1601	44	0
3	P	67	0	72	1	0
3	Q	67	0	72	3	0
All	All	6712	0	6530	183	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 183 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:119:PRO:HB3	2:H:147:TYR:HB3	1.44	0.98
1:M:150:ILE:HD11	1:M:179:LEU:HD21	1.64	0.79
1:L:7:SER:HB3	1:L:24:ARG:HH22	1.50	0.76
2:H:6:GLU:HB2	2:H:107:THR:HG23	1.69	0.72
1:L:115:VAL:HG22	1:L:136:LEU:HG	1.72	0.71

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	213/218 (98%)	194 (91%)	12 (6%)	7 (3%)	3	11
1	M	213/218 (98%)	191 (90%)	14 (7%)	8 (4%)	2	9
2	H	213/222 (96%)	183 (86%)	24 (11%)	6 (3%)	4	14
2	J	213/222 (96%)	188 (88%)	15 (7%)	10 (5%)	2	6
3	P	7/16 (44%)	6 (86%)	1 (14%)	0	100	100
3	Q	7/16 (44%)	6 (86%)	1 (14%)	0	100	100
All	All	866/912 (95%)	768 (89%)	67 (8%)	31 (4%)	2	10

5 of 31 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	L	28	ASP
1	L	170	ASP
2	H	128	CYS
2	H	215	SER
2	J	64	LYS

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	191/194 (98%)	149 (78%)	42 (22%)	1	3
1	M	191/194 (98%)	167 (87%)	24 (13%)	4	15
2	H	189/195 (97%)	152 (80%)	37 (20%)	1	5

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	J	189/195 (97%)	158 (84%)	31 (16%)	2	8
3	P	7/13 (54%)	5 (71%)	2 (29%)	0	1
3	Q	7/13 (54%)	5 (71%)	2 (29%)	0	1
All	All	774/804 (96%)	636 (82%)	138 (18%)	2	6

5 of 138 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	J	123	PRO
2	J	137	SER
2	J	203	GLN
2	H	38	ARG
2	H	34	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
2	J	16	GLN
1	M	145	ASN
2	H	203	GLN
1	M	137	ASN
2	H	76	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.