



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 01:46 PM UTC

PDB ID : 3GGF / pdb_00003ggf
Title : Crystal structure of human Serine/threonine-protein kinase MST4 in complex with an quinazolin
Authors : Chaikuad, A.; Rellos, P.; Eidarus, S.; Das, S.; Pike, A.C.W.; Sethi, R.; Fedorov, O.; Savitsky, P.; Roos, A.K.; Filippakopoulos, P.; Arrowsmith, C.H.; Edwards, A.M.; Weigelt, J.; Bountra, C.; von Delft, F.; Knapp, S.; Structural Genomics Consortium (SGC)
Deposited on : 2009-02-27
Resolution : 2.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1596 (2.36-2.36)
Clashscore	190562	1663 (2.36-2.36)
Ramachandran outliers	187476	1646 (2.36-2.36)
Sidechain outliers	187428	1646 (2.36-2.36)
RSRZ outliers	180081	1598 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	301	9% 81% 10% 9%
1	B	301	11% 79% 11% 10%

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4506 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine/threonine-protein kinase MST4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	275	Total	C	N	O	S	0	1	0
			2171	1397	356	411	7			
1	B	270	Total	C	N	O	S	0	0	0
			2124	1371	346	400	7			


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP Q9P289
B	0	SER	-	expression tag	UNP Q9P289

- Molecule 2 is CADMIUM ION (CCD ID: CD) (formula: Cd).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	4	Total	Cd	0	0
			4	4		
2	B	3	Total	Cd	0	0
			3	3		

- Molecule 3 is [4-({4-[(5-CYCLOPROPYL-1H-PYRAZOL-3-YL)AMINO]QUINAZOLIN-2-YL}IMINO)CYCLOHEXA-2,5-DIEN-1-YL]ACETONITRILE (CCD ID: GVD) (formula: C₂₂H₁₉N₇).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total 29	C 22	N 7	0	0
3	B	1	Total 29	C 22	N 7	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	79	Total O 79 79	0	0
4	B	67	Total O 67 67	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Serine/threonine-protein kinase MST4



- Molecule 1: Serine/threonine-protein kinase MST4



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	65.92Å 91.25Å 108.97Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.78 – 2.35 46.78 – 2.35	Depositor EDS
% Data completeness (in resolution range)	99.9 (46.78-2.35) 99.9 (46.78-2.35)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.03 (at 2.34Å)	Xtriage
Refinement program	REFMAC 5.5.0085	Depositor
R, R_{free}	0.219 , 0.277 0.234 , 0.286	Depositor DCC
R_{free} test set	1408 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	32.5	Xtriage
Anisotropy	0.148	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 40.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	4506	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 33.50 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 7.9252e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CD, GVD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.92	0/2216	0.93	0/2991
1	B	0.91	2/2168 (0.1%)	0.93	1/2928 (0.0%)
All	All	0.92	2/4384 (0.0%)	0.93	1/5919 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	236	PRO	C-O	-5.67	1.20	1.25
1	B	236	PRO	CA-C	5.13	1.54	1.51

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	64	ILE	N-CA-C	6.17	116.28	110.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2171	0	2160	36	2
1	B	2124	0	2119	22	2
2	A	4	0	0	0	1

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	3	0	0	0	1
3	A	29	0	18	2	0
3	B	29	0	18	2	0
4	A	79	0	0	7	0
4	B	67	0	0	1	2
All	All	4506	0	4315	52	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:222:MET:HE1	1:A:230:LEU:HD12	1.59	0.82
1:A:180:VAL:HG13	1:B:225:MET:HE2	1.62	0.81
1:B:222:MET:HE1	1:B:230:LEU:HD12	1.69	0.74
1:A:120:GLN:CD	1:A:287:LEU:HD11	2.14	0.72
1:A:33:GLY:CA	4:A:384:HOH:O	2.37	0.71
1:B:84:LYS:NZ	4:B:370:HOH:O	2.24	0.71
1:A:33:GLY:HA3	4:A:384:HOH:O	1.91	0.70
1:A:157:ASP:OD2	4:A:357:HOH:O	2.11	0.69
1:A:120:GLN:NE2	1:A:287:LEU:CD1	2.57	0.67
1:A:99:MET:HE3	3:A:305:GVD:HAM1	1.82	0.62
1:A:299:HIS:CG	1:A:299:HIS:O	2.52	0.62
1:A:33:GLY:N	4:A:384:HOH:O	2.33	0.61
1:B:114:GLY:O	1:B:290:ARG:NH2	2.33	0.60
1:A:222:MET:HE1	1:A:230:LEU:CD1	2.32	0.60
3:B:304:GVD:N1	3:B:304:GVD:HAK	2.18	0.59
1:A:295:LYS:HE2	4:A:348:HOH:O	2.02	0.59
1:A:61:GLU:CB	4:A:374:HOH:O	2.52	0.58
1:A:180:VAL:HG13	1:B:225:MET:CE	2.31	0.58
1:B:140:LYS:HB3	1:B:166:ALA:HB3	1.85	0.58
1:A:99:MET:CE	3:A:305:GVD:HAM1	2.36	0.56
3:B:304:GVD:N1	3:B:304:GVD:CAK	2.71	0.53
1:B:67:ILE:O	1:B:71:ILE:HG12	2.10	0.52
1:B:110:LEU:HD13	1:B:287:LEU:HB3	1.92	0.52
1:A:120:GLN:CD	1:A:287:LEU:CD1	2.84	0.51
1:A:225:MET:HE1	1:B:186:MET:HE1	1.93	0.50
1:A:59:GLU:OE2	1:B:112:ARG:NH2	2.45	0.49
1:A:35:PHE:HB2	1:A:55:ILE:HG22	1.96	0.48
1:A:180:VAL:HG21	1:B:228:LEU:HD22	1.96	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:56:ASP:OD1	1:B:58:GLU:HB2	2.13	0.48
1:A:31:GLY:HA2	4:A:335:HOH:O	2.14	0.46
1:A:225:MET:SD	1:B:186:MET:HE1	2.56	0.46
1:A:58:GLU:HA	1:A:93:SER:CB	2.45	0.46
1:B:152:LEU:N	1:B:152:LEU:HD12	2.33	0.44
1:A:117:ASP:OD1	1:A:117:ASP:C	2.61	0.43
1:A:133:ASP:OD1	1:A:266:LYS:HE3	2.18	0.43
1:B:25:THR:HB	1:B:42:ILE:CG2	2.49	0.43
1:B:120:GLN:CD	1:B:287:LEU:HD11	2.43	0.43
1:A:216:GLU:HB2	1:A:217:PRO:HD2	2.00	0.43
1:A:225:MET:HE2	1:B:180:VAL:HG13	2.00	0.43
1:B:186:MET:CE	1:B:190:VAL:HG12	2.49	0.43
1:A:180:VAL:CG1	1:B:225:MET:HE2	2.41	0.42
1:A:116:PHE:CE2	1:A:212:LEU:HD22	2.55	0.42
1:A:58:GLU:HA	1:A:93:SER:HB3	2.00	0.42
1:B:25:THR:HB	1:B:42:ILE:HG22	2.01	0.42
1:A:56:ASP:OD1	1:A:58:GLU:HB2	2.20	0.41
1:B:45:ARG:HG2	1:B:46:THR:HG23	2.02	0.41
1:A:120:GLN:NE2	1:A:287:LEU:HD11	2.29	0.41
1:A:133:ASP:O	1:A:133:ASP:CG	2.64	0.41
1:A:25:THR:HB	1:A:42:ILE:HG22	2.03	0.41
1:B:160:LEU:HB3	1:B:162:ASP:OD1	2.20	0.40
1:A:35:PHE:HB2	1:A:55:ILE:CG2	2.51	0.40
1:A:190:VAL:HG22	1:A:196:TYR:CG	2.56	0.40

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:78:ASP:OD2	4:B:369:HOH:O[4_457]	2.04	0.16
1:A:242:ASP:OD2	2:A:302:CD:CD[4_447]	2.11	0.09
1:A:249:GLU:OE2	4:B:356:HOH:O[4_547]	2.14	0.06
1:B:242:ASP:OD2	2:B:302:CD:CD[4_557]	2.17	0.03

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	272/301 (90%)	260 (96%)	12 (4%)	0	100	100
1	B	266/301 (88%)	254 (96%)	11 (4%)	1 (0%)	30	34
All	All	538/602 (89%)	514 (96%)	23 (4%)	1 (0%)	43	52

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	33	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	233/261 (89%)	232 (100%)	1 (0%)	84	91
1	B	227/261 (87%)	225 (99%)	2 (1%)	70	82
All	All	460/522 (88%)	457 (99%)	3 (1%)	76	86

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	266	LYS
1	B	140	LYS
1	B	266	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	68	GLN
1	A	299	HIS
1	B	256	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 9 ligands modelled in this entry, 7 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GVD	A	305	2	32,33,33	2.46	12 (37%)	35,46,46	2.82	10 (28%)
3	GVD	B	304	2	32,33,33	2.56	10 (31%)	35,46,46	3.12	15 (42%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GVD	A	305	2	-	3/12/27/27	0/5/5/5
3	GVD	B	304	2	-	5/12/27/27	0/5/5/5

All (22) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	304	GVD	CAT-CAF	-6.40	1.38	1.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	305	GVD	CAT-CAF	-5.92	1.39	1.50
3	A	305	GVD	NAS-NAN	-5.86	1.24	1.36
3	A	305	GVD	CAT-CAE	-5.72	1.39	1.50
3	B	304	GVD	CAT-CAE	-5.57	1.40	1.50
3	B	304	GVD	NAS-NAN	-5.29	1.25	1.36
3	B	304	GVD	CAG-CAE	4.38	1.41	1.33
3	B	304	GVD	CAK-CAV	-4.20	1.32	1.40
3	A	305	GVD	CAG-CAE	3.73	1.40	1.33
3	A	305	GVD	CAH-CAF	3.65	1.40	1.33
3	B	304	GVD	C2-NAQ	-3.38	1.30	1.37
3	A	305	GVD	C6-C5	-3.31	1.40	1.44
3	B	304	GVD	CAK-CAX	-3.02	1.32	1.37
3	A	305	GVD	CAK-CAV	-3.00	1.35	1.40
3	B	304	GVD	CAH-CAF	2.87	1.38	1.33
3	B	304	GVD	C4-N3	-2.85	1.33	1.37
3	A	305	GVD	CAG-CAU	-2.37	1.38	1.44
3	A	305	GVD	C4-N3	-2.22	1.34	1.37
3	A	305	GVD	CAK-CAX	-2.09	1.34	1.37
3	B	304	GVD	CAH-CAU	-2.09	1.38	1.44
3	A	305	GVD	CAX-NAS	-2.03	1.32	1.34
3	A	305	GVD	CAH-CAU	-2.00	1.39	1.44

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	305	GVD	CAV-NAN-NAS	8.33	109.77	103.71
3	A	305	GVD	C6-C5-C4	7.80	120.47	115.86
3	B	304	GVD	C6-C5-C4	7.75	120.44	115.86
3	B	304	GVD	CA0-CAT-CAE	7.14	125.07	111.86
3	B	304	GVD	CBB-CAX-CAK	-6.59	120.47	131.09
3	B	304	GVD	CAV-NAN-NAS	6.54	108.47	103.71
3	A	305	GVD	CAK-CAV-NAN	-5.81	107.99	112.14
3	A	305	GVD	CA0-CAT-CAF	5.58	122.19	111.86
3	B	304	GVD	CBB-CAX-NAS	4.59	129.63	122.62
3	B	304	GVD	CAK-CAX-NAS	4.43	109.88	106.19
3	B	304	GVD	CAK-CAV-NAN	-4.09	109.22	112.14
3	A	305	GVD	CA0-CAT-CAE	3.81	118.91	111.86
3	A	305	GVD	CAK-CAX-NAS	3.75	109.31	106.19
3	A	305	GVD	C5-C6-N1	-3.71	116.51	122.07
3	B	304	GVD	C5-C6-N1	-3.71	116.51	122.07
3	B	304	GVD	NAR-CAV-NAN	3.11	124.90	117.81
3	B	304	GVD	C5-C4-N3	-3.10	119.52	122.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	304	GVD	CAI-C4-C5	3.08	122.45	119.13
3	B	304	GVD	C2-N3-C4	2.90	119.59	115.23
3	A	305	GVD	C5-C4-N3	-2.54	120.11	122.80
3	A	305	GVD	CAJ-C5-C6	-2.35	120.77	124.16
3	A	305	GVD	C2-N3-C4	2.35	118.76	115.23
3	B	304	GVD	C5-C6-NAR	2.27	122.05	119.78
3	B	304	GVD	CA0-CAT-CAF	2.15	115.84	111.86
3	B	304	GVD	CAM-CBB-CAL	2.10	62.03	58.80

There are no chirality outliers.

All (8) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	305	GVD	CAB-CA0-CAT-CAF
3	B	304	GVD	CAG-CAU-NAQ-C2
3	B	304	GVD	CAH-CAU-NAQ-C2
3	B	304	GVD	CAB-CA0-CAT-CAE
3	A	305	GVD	CAG-CAU-NAQ-C2
3	A	305	GVD	CAH-CAU-NAQ-C2
3	B	304	GVD	CAK-CAX-CBB-CAM
3	B	304	GVD	CAK-CAX-CBB-CAL


There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	305	GVD	2	0
3	B	304	GVD	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	275/301 (91%)	0.46	28 (10%) 12 13	4, 19, 46, 83	1 (0%)
1	B	270/301 (89%)	0.49	33 (12%) 8 10	4, 19, 43, 73	0
All	All	545/602 (90%)	0.47	61 (11%) 10 12	4, 19, 44, 83	1 (0%)

All (61) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	33	GLY	6.1
1	A	35	PHE	5.3
1	A	32	LYS	5.3
1	B	114	GLY	5.0
1	B	34	SER	4.9
1	A	166	ALA	4.5
1	B	296	ALA	4.5
1	B	33	GLY	4.4
1	B	195	ALA	4.4
1	B	35	PHE	4.4
1	B	115	PRO	4.3
1	B	294	TRP	4.3
1	B	295	LYS	4.2
1	B	291	PHE	4.2
1	A	34	SER	4.2
1	A	62[A]	ASP	4.2
1	B	179	PHE	4.0
1	A	260	SER	3.8
1	A	18	ALA	3.8
1	A	195	ALA	3.7
1	A	66	ASP	3.4
1	B	177	ASN	3.4
1	B	65	GLU	3.4
1	A	283	TYR	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	298	GLY	3.1
1	B	18	ALA	3.1
1	B	36	GLY	3.1
1	A	300	SER	3.1
1	A	17	ILE	3.0
1	B	166	ALA	3.0
1	A	296	ALA	3.0
1	A	31	GLY	3.0
1	B	282	SER	2.9
1	B	297	GLU	2.9
1	A	165	VAL	2.9
1	B	56	ASP	2.8
1	A	76	GLN	2.7
1	B	32	LYS	2.7
1	B	66	ASP	2.7
1	A	178	THR	2.5
1	B	293	ARG	2.5
1	B	283	TYR	2.5
1	A	197	ASP	2.5
1	A	196	TYR	2.4
1	A	261	PHE	2.4
1	A	293	ARG	2.4
1	A	177	ASN	2.3
1	A	160	LEU	2.3
1	B	196	TYR	2.3
1	B	113	ALA	2.2
1	A	27	LEU	2.2
1	B	62	ASP	2.2
1	B	93	SER	2.2
1	A	29	ARG	2.1
1	B	64	ILE	2.1
1	A	299	HIS	2.1
1	B	59	GLU	2.1
1	B	288	ILE	2.1
1	B	61	GLU	2.0
1	B	58	GLU	2.0
1	B	112	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CD	B	303	1/1	0.81	0.13	59,59,59,59	1
3	GVD	B	304	29/29	0.93	0.12	5,22,44,55	0
3	GVD	A	305	29/29	0.94	0.11	7,20,32,41	0
2	CD	A	303	1/1	0.97	0.06	68,68,68,68	0
2	CD	B	302	1/1	0.99	0.07	22,22,22,22	0
2	CD	A	301	1/1	0.99	0.04	43,43,43,43	1
2	CD	A	304	1/1	0.99	0.03	39,39,39,39	0
2	CD	B	301	1/1	0.99	0.03	48,48,48,48	0
2	CD	A	302	1/1	1.00	0.08	23,23,23,23	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around GVD B 304:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.