



Full wwPDB NMR Structure Validation Report ⓘ

Mar 8, 2026 – 07:00 AM UTC

PDB ID : 1GHU / pdb_00001ghu
Title : NMR solution structure of growth factor receptor-bound protein 2 (GRB2) SH2 domain, 24 structures
Authors : Thornton, K.H.; Mueller, W.T.; McConnell, P.; Zhu, G.; Saltiel, A.R.; Thanabal, V.
Deposited on : 1996-08-05

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	107	

2 Ensemble composition and analysis

This entry contains 24 models. Model 24 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:60-A:152 (93)	0.94	24

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters and 9 single-model clusters were found.

Cluster number	Models
1	2, 7, 9, 19, 24
2	4, 14, 15, 21
3	11, 12, 16, 22
4	13, 23
Single-model clusters	1; 3; 5; 6; 8; 10; 17; 18; 20

3 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 1538 atoms, of which 760 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called GRB2.


Mol	Chain	Residues	Atoms						Trace
1	A	95	Total	C	H	N	O	S	0
			1538	498	760	139	140	1	

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: GRB2




4.2 Scores per residue for each member of the ensemble

Colouring as in section [4.1](#) above.


4.2.1 Score per residue for model 1

- Molecule 1: GRB2



4.2.2 Score per residue for model 2

- Molecule 1: GRB2



4.2.3 Score per residue for model 3

- Molecule 1: GRB2

Chain A:  72% 15% • 11%



4.2.4 Score per residue for model 4

- Molecule 1: GRB2

Chain A:  78% 8% • 11%



4.2.5 Score per residue for model 5

- Molecule 1: GRB2

Chain A:  79% 8% • 11%



4.2.6 Score per residue for model 6

- Molecule 1: GRB2

Chain A:  76% 11% • 11%



4.2.7 Score per residue for model 7

- Molecule 1: GRB2

Chain A:  74% 13% • 11%



4.2.8 Score per residue for model 8

- Molecule 1: GRB2

Chain A:  77% 10% • 11%



4.2.9 Score per residue for model 9

- Molecule 1: GRB2

Chain A:  73% 14% • 11%



4.2.10 Score per residue for model 10

- Molecule 1: GRB2

Chain A:  74% 12% •• 11%



4.2.11 Score per residue for model 11

- Molecule 1: GRB2

Chain A:  79% 8% • 11%



4.2.12 Score per residue for model 12


- Molecule 1: GRB2

Chain A:  77% 9% •• 11%




4.2.13 Score per residue for model 13

- Molecule 1: GRB2




4.2.14 Score per residue for model 14

- Molecule 1: GRB2




4.2.15 Score per residue for model 15

- Molecule 1: GRB2




4.2.16 Score per residue for model 16

- Molecule 1: GRB2



4.2.17 Score per residue for model 17

- Molecule 1: GRB2



4.2.18 Score per residue for model 18

- Molecule 1: GRB2

Chain A:  70% 15% 11%



4.2.19 Score per residue for model 19

- Molecule 1: GRB2


Chain A:  74% 12% 11%



4.2.20 Score per residue for model 20

- Molecule 1: GRB2


Chain A:  73% 13% 11%



4.2.21 Score per residue for model 21

- Molecule 1: GRB2

Chain A:  73% 13% 11%



4.2.22 Score per residue for model 22

- Molecule 1: GRB2

Chain A:  76% 11% 11%



4.2.23 Score per residue for model 23

- Molecule 1: GRB2


Chain A:  76% 10% .. 11%



4.2.24 Score per residue for model 24 (medoid)

- Molecule 1: GRB2

Chain A:  75% 11% .. 11%



5 Refinement protocol and experimental data overview

Of the ? calculated structures, 24 were deposited, based on the following criterion: ?.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
DGII	refinement	

No chemical shift data was provided.

6 Model quality

6.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.70±0.01	2±0/787 (0.3± 0.1%)	1.73±0.03	15±2/1056 (1.5± 0.2%)
All	All	1.70	52/18888 (0.3%)	1.73	368/25344 (1.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	0.0±0.2
All	All	0	1

All unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	A	65	ILE	CA-C	6.86	1.58	1.52	15	4
1	A	62	PHE	CA-C	6.39	1.56	1.52	20	2
1	A	107	HIS	ND1-CE1	5.83	1.38	1.32	18	24
1	A	79	HIS	ND1-CE1	5.33	1.37	1.32	8	21
1	A	135	HIS	ND1-CE1	5.00	1.37	1.32	5	1

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	107	HIS	ND1-CE1-NE2	12.34	120.74	108.40	19	24
1	A	135	HIS	ND1-CE1-NE2	12.34	120.73	108.40	8	24
1	A	79	HIS	ND1-CE1-NE2	12.09	120.49	108.40	14	24
1	A	125	PHE	CA-CB-CG	11.23	125.03	113.80	17	1
1	A	127	SER	N-CA-CB	-10.88	97.84	110.35	17	1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	127	SER	N-CA-C	10.67	124.09	110.65	17	1
1	A	107	HIS	CE1-NE2-CD2	-9.65	99.35	109.00	8	24
1	A	135	HIS	CE1-NE2-CD2	-9.55	99.45	109.00	17	24
1	A	79	HIS	CE1-NE2-CD2	-9.38	99.62	109.00	1	24
1	A	101	PHE	CA-CB-CG	9.30	123.10	113.80	2	6
1	A	113	ASP	CA-CB-CG	7.73	120.33	112.60	9	23
1	A	129	ASN	CA-CB-CG	7.53	120.13	112.60	1	5
1	A	133	ASP	CA-CB-CG	7.15	119.75	112.60	9	4
1	A	95	PHE	CA-CB-CG	7.09	120.89	113.80	19	4
1	A	147	PHE	CA-CB-CG	6.67	120.47	113.80	3	10
1	A	119	PHE	CA-CB-CG	6.66	120.46	113.80	16	23
1	A	65	ILE	CB-CA-C	6.42	117.24	110.37	18	1
1	A	126	ASN	N-CA-C	6.40	118.79	111.11	5	8
1	A	107	HIS	CG-ND1-CE1	-6.31	98.57	109.30	19	24
1	A	135	HIS	CG-ND1-CE1	-6.31	98.57	109.30	19	24
1	A	79	HIS	CG-ND1-CE1	-6.26	98.66	109.30	1	24
1	A	143	ASN	CA-CB-CG	6.21	118.81	112.60	18	5
1	A	129	ASN	N-CA-CB	6.19	120.95	110.49	10	5
1	A	126	ASN	CA-CB-CG	6.13	118.73	112.60	17	6
1	A	78	ARG	N-CA-C	6.11	120.20	112.87	8	10
1	A	142	ARG	N-CA-C	5.90	117.71	111.28	18	1
1	A	74	LEU	N-CA-C	5.81	117.41	111.14	18	2
1	A	88	SER	N-CA-C	5.79	120.07	113.19	9	1
1	A	60	TRP	N-CA-C	5.77	117.57	111.28	9	2
1	A	132	VAL	N-CA-CB	5.72	117.24	110.55	15	1
1	A	146	ILE	CA-CB-CG1	5.68	120.06	110.40	4	3
1	A	80	ASP	CA-CB-CG	5.60	118.20	112.60	21	10
1	A	104	ASP	CA-CB-CG	5.56	118.16	112.60	16	3
1	A	83	PHE	N-CA-C	5.56	118.14	109.52	18	1
1	A	67	ARG	N-CA-C	5.47	117.33	111.36	18	1
1	A	89	GLU	N-CA-C	5.44	116.90	110.97	20	1
1	A	79	HIS	CB-CG-CD2	-5.34	124.25	131.20	1	2
1	A	75	SER	N-CA-C	5.32	117.08	111.28	24	1
1	A	135	HIS	CB-CG-CD2	-5.21	124.43	131.20	5	1
1	A	66	PRO	N-CA-C	5.10	119.22	111.11	13	1
1	A	65	ILE	CA-CB-CG1	5.10	119.06	110.40	1	1
1	A	103	ASN	CA-CB-CG	5.09	117.69	112.60	19	2
1	A	105	VAL	CB-CA-C	5.07	117.01	111.08	18	1
1	A	90	SER	N-CA-CB	5.05	119.03	110.49	1	1
1	A	108	PHE	CA-CB-CG	5.04	118.84	113.80	15	1
1	A	90	SER	N-CA-C	5.02	116.75	111.28	21	1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	150	ASP	CA-CB-CG	5.00	117.60	112.60	3	1

There are no chirality outliers.

All unique planar outliers are listed below.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	83	PHE	Sidechain	1

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
All	All	18432	18024	18024	-

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is -.

There are no clashes.

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	92/107 (86%)	81±2 (88±2%)	9±2 (10±3%)	2±1 (2±1%)	8	48
All	All	2208/2568 (86%)	1943 (88%)	218 (10%)	47 (2%)	8	48

All 12 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	122	VAL	12
1	A	129	ASN	5
1	A	114	GLY	5
1	A	115	ALA	4
1	A	142	ARG	4
1	A	143	ASN	4
1	A	90	SER	3
1	A	63	GLY	3
1	A	116	GLY	3
1	A	60	TRP	2
1	A	89	GLU	1
1	A	121	TRP	1

6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	82/95 (86%)	79±1 (97±1%)	3±1 (3±1%)	35	84
All	All	1968/2280 (86%)	1906 (97%)	62 (3%)	35	84

All 22 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	77	GLN	19
1	A	65	ILE	7
1	A	124	LYS	7
1	A	146	ILE	4
1	A	120	LEU	4
1	A	132	VAL	2
1	A	140	VAL	2
1	A	151	ILE	2
1	A	117	LYS	2
1	A	89	GLU	1
1	A	111	LEU	1
1	A	76	LYS	1
1	A	100	LYS	1
1	A	72	GLU	1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	148	LEU	1
1	A	150	ASP	1
1	A	118	TYR	1
1	A	125	PHE	1
1	A	130	GLU	1
1	A	145	GLN	1
1	A	74	LEU	1
1	A	87	GLU	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided