



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 18, 2026 – 06:49 AM UTC

PDB ID : 4GH8 / pdb_00004gh8
Title : Crystal structure of a 'humanized' E. coli dihydrofolate reductase
Authors : French, J.B.; Liu, C.T.; Hanoian, P.; Pringle, T.H.; Hammes-Schiffer, S.; Benkovic, S.J.
Deposited on : 2012-08-07
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3428 (1.86-1.86)
Clashscore	190562	3579 (1.86-1.86)
Ramachandran outliers	187476	3553 (1.86-1.86)
Sidechain outliers	187428	3553 (1.86-1.86)
RSRZ outliers	180081	3429 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	162	
1	B	162	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 2878 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Dihydrofolate reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	162	Total	C	N	O	S	0	0	0
			1270	812	214	237	7			
1	A	162	Total	C	N	O	S	0	0	0
			1274	815	217	235	7			

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	23	PRO	ASN	SEE REMARK 999	UNP P0ABQ4
B	24	PRO	ASN	SEE REMARK 999	UNP P0ABQ4
B	52	PRO	GLY	SEE REMARK 999	UNP P0ABQ4
B	53	GLU	GLY	SEE REMARK 999	UNP P0ABQ4
B	54	LYS	GLY	SEE REMARK 999	UNP P0ABQ4
B	55	ASN	GLY	SEE REMARK 999	UNP P0ABQ4
A	23	PRO	ASN	SEE REMARK 999	UNP P0ABQ4
A	24	PRO	ASN	SEE REMARK 999	UNP P0ABQ4
A	52	PRO	GLY	SEE REMARK 999	UNP P0ABQ4
A	53	GLU	GLY	SEE REMARK 999	UNP P0ABQ4
A	54	LYS	GLy	SEE REMARK 999	UNP P0ABQ4
A	55	ASN	GLY	SEE REMARK 999	UNP P0ABQ4

- Molecule 2 is METHOTREXATE (CCD ID: MTX) (formula: $C_{20}H_{22}N_8O_5$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	B	1	Total 33	C 20	N 8	O 5	0	0
2	A	1	Total 33	C 20	N 8	O 5	0	0

- Molecule 3 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (CCD ID: NDP) (formula: $\text{C}_{21}\text{H}_{30}\text{N}_7\text{O}_{17}\text{P}_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	B	1	Total 48	C 21	N 7	O 17	P 3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 4 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Ca	0	0
			1	1		
4	A	2	Total	Ca	0	0
			2	2		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	74	Total	O	0	0
			74	74		
5	A	95	Total	O	0	0
			95	95		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Dihydrofolate reductase



- Molecule 1: Dihydrofolate reductase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	52.24Å 63.77Å 62.44Å 90.00° 106.82° 90.00°	Depositor
Resolution (Å)	50.00 – 1.85 50.00 – 1.85	Depositor EDS
% Data completeness (in resolution range)	95.1 (50.00-1.85) 95.1 (50.00-1.85)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.25 (at 1.85Å)	Xtriage
Refinement program	REFMAC 5.5.0110	Depositor
R, R_{free}	0.203 , 0.245 0.207 , 0.246	Depositor DCC
R_{free} test set	1631 reflections (4.87%)	wwPDB-VP
Wilson B-factor (Å ²)	16.9	Xtriage
Anisotropy	1.090	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 41.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2878	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 50.05 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.8394e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NDP, MTX, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.02	2/1311 (0.2%)	0.98	1/1789 (0.1%)
1	B	1.02	0/1307	0.98	0/1785
All	All	1.02	2/2618 (0.1%)	0.98	1/3574 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	5	ILE	CA-CB	5.48	1.61	1.54
1	A	10	VAL	CA-CB	5.26	1.59	1.54

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	18	ASN	N-CA-C	5.36	119.10	112.24

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1274	0	1226	8	0
1	B	1270	0	1215	12	0
2	A	33	0	20	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	33	0	20	1	0
3	A	48	0	26	0	0
3	B	48	0	26	2	0
4	A	2	0	0	0	0
4	B	1	0	0	0	0
5	A	95	0	0	1	0
5	B	74	0	0	0	0
All	All	2878	0	2533	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:20:MET:HE3	1:A:22:TRP:CZ2	2.25	0.72
1:A:20:MET:HE3	1:A:22:TRP:HZ2	1.64	0.63
1:B:25:LEU:HD12	1:B:119:ILE:HD12	1.82	0.61
1:B:16:MET:HE3	1:B:17:GLU:HG2	1.85	0.56
1:B:149:ALA:HB3	1:B:150:GLN:HE22	1.72	0.55
1:A:20:MET:CE	1:A:22:TRP:CZ2	2.94	0.50
1:A:142:SER:HA	1:A:157:PHE:O	2.14	0.48
1:A:8:LEU:HD23	1:A:12:ARG:HG2	1.95	0.48
1:B:142:SER:HA	1:B:157:PHE:O	2.15	0.47
1:B:25:LEU:CD1	1:B:119:ILE:HD12	2.45	0.47
1:A:112:GLN:NE2	5:A:337:HOH:O	2.48	0.47
2:A:201:MTX:H13	2:A:201:MTX:HM1	1.61	0.47
1:B:150:GLN:N	1:B:150:GLN:CD	2.73	0.46
2:B:201:MTX:H13	2:B:201:MTX:HM1	1.52	0.45
1:B:25:LEU:HD22	1:B:151:ASN:HB3	1.99	0.44
1:B:18:ASN:HA	3:B:202:NDP:H3D	1.99	0.43
1:B:133:GLU:N	1:B:134:PRO:HD2	2.33	0.42
1:B:149:ALA:HB3	1:B:150:GLN:NE2	2.34	0.42
1:A:23:PRO:O	1:A:24:PRO:C	2.63	0.41
1:B:4:LEU:HD13	1:B:111:ALA:HB2	2.03	0.41
1:B:14:ILE:O	3:B:202:NDP:H2N	2.20	0.41
1:A:148:ASP:OD1	1:A:151:ASN:N	2.51	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	160/162 (99%)	158 (99%)	2 (1%)	0	100	100
1	B	160/162 (99%)	156 (98%)	4 (2%)	0	100	100
All	All	320/324 (99%)	314 (98%)	6 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	135/140 (96%)	133 (98%)	2 (2%)	57	47
1	B	135/140 (96%)	134 (99%)	1 (1%)	76	70
All	All	270/280 (96%)	267 (99%)	3 (1%)	65	57

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	63	ASN
1	A	63	ASN
1	A	105	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	150	GLN
1	A	55	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 3 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	MTX	B	201	-	35,35,35	1.75	4 (11%)	47,49,49	2.21	17 (36%)
2	MTX	A	201	-	35,35,35	1.77	6 (17%)	47,49,49	1.89	12 (25%)
3	NDP	A	202	-	51,52,52	1.60	13 (25%)	71,80,80	1.91	14 (19%)
3	NDP	B	202	-	51,52,52	1.60	9 (17%)	71,80,80	1.74	12 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MTX	B	201	-	-	6/25/25/25	0/3/3/3
2	MTX	A	201	-	-	5/25/25/25	0/3/3/3
3	NDP	A	202	-	-	4/34/77/77	0/5/5/5
3	NDP	B	202	-	-	3/34/77/77	0/5/5/5

All (32) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	201	MTX	C7-N8	6.33	1.42	1.31
2	A	201	MTX	C7-N8	5.67	1.41	1.31
2	B	201	MTX	C9-C6	-4.73	1.43	1.51
2	A	201	MTX	C9-C6	-4.66	1.43	1.51
3	A	202	NDP	PA-O3	4.47	1.64	1.59
2	B	201	MTX	C11-C	-4.04	1.41	1.50
2	A	201	MTX	C11-C	-3.84	1.41	1.50
3	B	202	NDP	C2A-N1A	3.76	1.40	1.33
3	B	202	NDP	C4N-C3N	-3.73	1.43	1.50
3	B	202	NDP	O2D-C2D	3.55	1.51	1.43
3	A	202	NDP	C4N-C3N	-3.15	1.44	1.50
3	A	202	NDP	P2B-O3X	-3.13	1.43	1.54
3	A	202	NDP	C5A-N7A	-3.07	1.33	1.39
3	B	202	NDP	C5A-N7A	-2.81	1.34	1.39
3	A	202	NDP	C2A-N1A	2.74	1.38	1.33
3	B	202	NDP	C4N-C5N	-2.69	1.42	1.49
3	A	202	NDP	O4B-C4B	-2.68	1.39	1.45
3	B	202	NDP	C7N-C3N	-2.60	1.43	1.48
2	A	201	MTX	C6-N5	2.58	1.36	1.32
3	A	202	NDP	C8A-N9A	-2.39	1.33	1.37
3	A	202	NDP	C6N-C5N	2.38	1.40	1.33
3	B	202	NDP	C1B-N9A	2.38	1.52	1.46
3	A	202	NDP	C4N-C5N	-2.30	1.43	1.49
3	B	202	NDP	P2B-O3X	-2.26	1.46	1.54
2	B	201	MTX	C6-N5	2.24	1.36	1.32
3	A	202	NDP	C6N-N1N	2.19	1.42	1.37
3	A	202	NDP	C2A-N3A	2.16	1.37	1.33
2	A	201	MTX	C16-C11	2.12	1.42	1.39
3	B	202	NDP	P2B-O2X	-2.11	1.47	1.54
3	A	202	NDP	PN-O5D	2.06	1.67	1.59
3	A	202	NDP	C7N-C3N	-2.00	1.44	1.48
2	A	201	MTX	C13-C12	2.00	1.42	1.38

All (55) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	202	NDP	N3A-C2A-N1A	-6.53	118.69	128.58
2	B	201	MTX	C2-N1-C8A	6.25	122.22	115.48
3	A	202	NDP	N3A-C2A-N1A	-5.75	119.88	128.58
3	A	202	NDP	C4A-C5A-N7A	-5.21	104.63	110.58
3	A	202	NDP	C5A-C4A-N3A	-5.17	119.59	126.72
2	B	201	MTX	CB-CG-CD	-5.03	99.08	112.49
2	A	201	MTX	C2-N1-C8A	5.00	120.87	115.48
3	A	202	NDP	C5A-N7A-C8A	4.70	110.84	103.45
2	B	201	MTX	N1-C2-N3	-4.50	121.49	127.21
3	B	202	NDP	N9A-C8A-N7A	-4.44	107.63	113.94
3	A	202	NDP	N9A-C8A-N7A	-4.24	107.93	113.94
3	B	202	NDP	C5A-N7A-C8A	4.13	109.94	103.45
3	B	202	NDP	C4A-C5A-N7A	-4.11	105.88	110.58
2	B	201	MTX	C9-N10-C14	4.10	125.14	120.17
2	B	201	MTX	N8-C8A-N1	4.03	120.16	115.77
3	B	202	NDP	C5A-C4A-N3A	-3.95	121.28	126.72
2	A	201	MTX	N1-C2-N3	-3.84	122.33	127.21
3	A	202	NDP	C5A-C4A-N9A	3.77	109.92	105.81
2	A	201	MTX	CG-CB-CA	-3.72	106.31	113.16
2	A	201	MTX	C9-N10-C14	3.54	124.46	120.17
2	A	201	MTX	CB-CG-CD	-3.42	103.37	112.49
3	A	202	NDP	C2A-N3A-C4A	3.37	120.06	111.83
2	A	201	MTX	C6-C7-N8	-3.36	119.92	123.14
3	A	202	NDP	O3X-P2B-O2X	3.25	120.00	107.80
3	A	202	NDP	O2B-P2B-O1X	-3.25	97.74	109.33
2	A	201	MTX	N8-C8A-N1	3.22	119.27	115.77
3	B	202	NDP	C5A-C4A-N9A	3.13	109.22	105.81
2	B	201	MTX	C6-C7-N8	-3.08	120.18	123.14
3	B	202	NDP	C2A-N3A-C4A	2.91	118.94	111.83
2	B	201	MTX	CG-CB-CA	-2.88	107.85	113.16
2	A	201	MTX	C7-C6-N5	-2.85	119.02	120.87
2	B	201	MTX	OE1-CD-CG	-2.83	114.10	123.09
3	A	202	NDP	C6A-C5A-C4A	2.81	121.01	117.18
2	B	201	MTX	C4-C4A-N5	2.77	122.46	120.33
3	A	202	NDP	O2A-PA-O3	2.72	114.63	107.27
2	A	201	MTX	C4-C4A-N5	2.65	122.37	120.33
2	B	201	MTX	CB-CA-N	-2.50	105.96	110.91
3	A	202	NDP	O4B-C4B-C5B	-2.48	101.39	109.33
2	B	201	MTX	C4A-C8A-N1	-2.47	117.91	121.74
2	A	201	MTX	C8A-C4A-N5	-2.45	119.62	122.35
2	B	201	MTX	C8A-C4A-N5	-2.42	119.65	122.35
2	B	201	MTX	C7-C6-N5	-2.40	119.31	120.87
3	B	202	NDP	C6A-C5A-C4A	2.38	120.43	117.18

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	202	NDP	C1D-N1N-C2N	-2.34	117.28	121.14
3	B	202	NDP	C3D-C2D-C1D	-2.33	97.05	101.46
2	B	201	MTX	CA-N-C	2.32	127.14	121.56
3	B	202	NDP	O2A-PA-O3	2.29	113.46	107.27
3	A	202	NDP	O4D-C1D-N1N	-2.18	103.93	108.08
2	B	201	MTX	C6-N5-C4A	2.17	121.25	118.17
2	B	201	MTX	C13-C14-N10	-2.16	118.58	121.59
3	B	202	NDP	C1D-N1N-C2N	-2.14	117.61	121.14
2	A	201	MTX	C6-N5-C4A	2.14	121.19	118.17
3	B	202	NDP	O3X-P2B-O2X	2.10	115.68	107.80
2	A	201	MTX	O1-CT-CA	-2.06	115.60	122.26
2	B	201	MTX	OE2-CD-OE1	2.03	128.56	123.33

There are no chirality outliers.

All (18) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	201	MTX	N-CA-CB-CG
2	A	201	MTX	CT-CA-CB-CG
2	B	201	MTX	CT-CA-CB-CG
2	B	201	MTX	CA-CB-CG-CD
2	B	201	MTX	N-CA-CB-CG
2	B	201	MTX	C6-C9-N10-CM
2	A	201	MTX	C6-C9-N10-CM
3	A	202	NDP	O4D-C1D-N1N-C2N
3	B	202	NDP	O4D-C1D-N1N-C2N
2	B	201	MTX	OE1-CD-CG-CB
2	A	201	MTX	OE1-CD-CG-CB
2	A	201	MTX	OE2-CD-CG-CB
3	B	202	NDP	C2N-C3N-C7N-N7N
3	A	202	NDP	C2N-C3N-C7N-N7N
2	B	201	MTX	OE2-CD-CG-CB
3	B	202	NDP	C2B-O2B-P2B-O3X
3	A	202	NDP	C2B-O2B-P2B-O3X
3	A	202	NDP	C2D-C1D-N1N-C2N

There are no ring outliers.

3 monomers are involved in 4 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	201	MTX	1	0


Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	201	MTX	1	0
3	B	202	NDP	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	162/162 (100%)	0.02	3 (1%) 66 70	10, 20, 31, 41	0
1	B	162/162 (100%)	0.17	3 (1%) 66 70	12, 21, 35, 42	0
All	All	324/324 (100%)	0.10	6 (1%) 66 70	10, 21, 35, 42	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	135	ASP	3.2
1	B	149	ALA	3.1
1	B	120	ASP	2.9
1	A	134	PRO	2.6
1	B	150	GLN	2.2
1	A	133	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	MTX	A	201	33/33	0.93	0.08	11,19,35,39	0
2	MTX	B	201	33/33	0.94	0.08	12,21,35,41	0
3	NDP	B	202	48/48	0.97	0.07	13,19,24,26	0
3	NDP	A	202	48/48	0.97	0.07	12,17,22,25	0
4	CA	B	203	1/1	0.98	0.05	27,27,27,27	0
4	CA	A	203	1/1	0.99	0.07	26,26,26,26	0
4	CA	A	204	1/1	0.99	0.03	18,18,18,18	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around MTX B 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NDP B 202:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around NDP A 202:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.