



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 10:59 AM UTC

PDB ID : 2GIJ / pdb_00002gij
Title : Q138F HincII bound to cognate DNA GTTAAC and Ca²⁺
Authors : Horton, N.C.; Joshi, H.K.
Deposited on : 2006-03-28
Resolution : 1.93 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.93 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1452 (1.94-1.94)
Clashscore	190562	1494 (1.94-1.94)
Sidechain outliers	187428	1479 (1.94-1.94)
RSRZ outliers	180081	1453 (1.94-1.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	E	14	71% 21% 7%
1	F	14	79% 14% 7%
2	A	257	5% 75% 18% • 5%
2	B	257	14% 62% 28% • 8%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4766 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called 5'-D(*GP*CP*CP*GP*GP*TP*TP*AP*AP*CP*CP*GP*GP*C)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	E	14	Total	C	N	O	P	0	0	0
			284	135	54	82	13			
1	F	14	Total	C	N	O	P	0	0	0
			284	135	54	82	13			

- Molecule 2 is a protein called Type II restriction enzyme HincII.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	245	Total	C	N	O	S	0	0	0
			1958	1274	306	372	6			
2	B	236	Total	C	N	O	S	0	0	0
			1907	1244	301	356	6			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	130	THR	ARG	conflict	UNP P17743
A	138	PHE	GLN	engineered mutation	UNP P17743
A	173	TRP	SER	conflict	UNP P17743
B	130	THR	ARG	conflict	UNP P17743
B	138	PHE	GLN	engineered mutation	UNP P17743
B	173	TRP	SER	conflict	UNP P17743

- Molecule 3 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	F	1	Total	Na	0	0
			1	1		

- Molecule 4 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total 1	Ca 1	0	0
4	B	1	Total 1	Ca 1	0	0

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	E	52	Total 52	O 52	0	0
5	F	29	Total 29	O 29	0	0
5	A	157	Total 157	O 157	0	0
5	B	92	Total 92	O 92	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 5'-D(*GP*CP*CP*GP*GP*TP*TP*AP*AP*CP*CP*GP*GP*C)-3'

Chain E: 




- Molecule 1: 5'-D(*GP*CP*CP*GP*GP*TP*TP*AP*AP*CP*CP*GP*GP*C)-3'

Chain F: 




- Molecule 2: Type II restriction enzyme HincII

Chain A: 



- Molecule 2: Type II restriction enzyme HincII

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	50.46Å 91.77Å 66.54Å 90.00° 104.46° 90.00°	Depositor
Resolution (Å)	50.00 – 1.93 50.00 – 1.93	Depositor EDS
% Data completeness (in resolution range)	99.7 (50.00-1.93) 99.0 (50.00-1.93)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.88 (at 1.88Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.233 , 0.263 0.234 , 0.263	Depositor DCC
R_{free} test set	1977 reflections (4.20%)	wwPDB-VP
Wilson B-factor (Å ²)	35.3	Xtriage
Anisotropy	0.441	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 54.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4766	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.47% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	E	0.29	0/318	0.79	0/489
1	F	0.28	0/318	0.80	1/489 (0.2%)
2	A	0.39	0/2004	0.88	6/2724 (0.2%)
2	B	0.36	0/1951	0.84	5/2643 (0.2%)
All	All	0.36	0/4591	0.85	12/6345 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	E	0	1

There are no bond length outliers.

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	55	LYS	N-CA-C	-9.87	97.23	110.55
2	B	55	LYS	N-CA-C	-9.26	96.82	110.52
2	A	208	ILE	N-CA-C	-6.52	101.17	109.58
2	A	89	LEU	N-CA-C	6.20	121.50	113.88
2	B	208	ILE	N-CA-C	-6.08	100.94	109.21
2	B	211	HIS	N-CA-C	-5.57	100.61	109.96
1	F	8	DA	N9-C1'-C2'	-5.44	105.34	113.50
2	B	89	LEU	N-CA-C	5.40	120.88	113.97
2	B	40	LEU	N-CA-C	-5.40	106.70	113.72
2	A	123	TYR	N-CA-C	5.29	118.03	109.40
2	A	90	SER	N-CA-C	5.14	117.31	110.53
2	A	41	VAL	N-CA-C	-5.01	105.61	110.42

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	E	9	DA	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	284	0	158	6	0
1	F	284	0	158	2	0
2	A	1958	0	1835	43	0
2	B	1907	0	1820	66	0
3	F	1	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	157	0	0	0	0
5	B	92	0	0	2	0
5	E	52	0	0	1	0
5	F	29	0	0	0	0
All	All	4766	0	3971	111	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (111) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:35:GLU:HB2	2:A:36:PRO:HD3	1.54	0.87
2:A:55:LYS:HB2	2:A:58:GLU:HG3	1.58	0.85
1:E:8:DA:H1'	1:E:9:DA:H5'	1.58	0.84
2:A:66:LYS:NZ	2:A:67:ASN:HD21	1.79	0.81
2:A:11:ILE:HG12	2:A:44:PHE:CE2	2.16	0.80
2:B:4:ILE:CD1	2:B:126:LEU:HD22	2.11	0.80
2:B:251:LYS:C	2:B:251:LYS:HD3	2.12	0.74
2:B:56:GLN:NE2	2:B:113:ALA:H	1.90	0.69
2:A:66:LYS:HZ3	2:A:67:ASN:HD21	1.39	0.69

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:70:ILE:HD12	2:B:79:LEU:HD21	1.73	0.69
2:B:55:LYS:HB2	2:B:58:GLU:HG3	1.74	0.69
2:A:254:LYS:O	2:A:258:LEU:HG	1.93	0.69
2:B:170:GLU:HB2	2:B:212:VAL:HB	1.74	0.68
2:B:195:PRO:O	2:B:198:LEU:HD23	1.94	0.68
2:A:132:ASN:C	2:A:132:ASN:HD22	2.04	0.65
2:B:11:ILE:HG12	2:B:44:PHE:CE2	2.32	0.64
2:A:35:GLU:CB	2:A:36:PRO:HD3	2.27	0.63
1:E:9:DA:H2''	1:E:10:DC:OP2	1.97	0.62
2:B:193:SER:HB3	2:B:215:LEU:HD11	1.81	0.61
2:A:52:LEU:O	2:A:117:LEU:HA	2.01	0.61
2:A:203:ALA:O	2:B:205:ALA:HA	2.01	0.61
2:B:215:LEU:HD13	2:B:216:ASP:O	2.01	0.60
2:A:2:SER:HB3	2:A:5:LYS:HB2	1.83	0.59
2:A:101:ILE:HG22	2:A:102:GLU:OE2	2.03	0.58
2:A:180:LEU:HD12	2:A:180:LEU:N	2.19	0.58
2:B:100:SER:HB2	5:B:489:HOH:O	2.02	0.58
2:A:20:VAL:HG22	2:A:180:LEU:HB2	1.86	0.57
2:A:20:VAL:CG2	2:A:180:LEU:HB2	2.34	0.57
2:A:132:ASN:ND2	2:A:134:SER:H	2.02	0.57
2:B:65:MET:O	2:B:68:PRO:HD3	2.06	0.56
2:B:70:ILE:CD1	2:B:79:LEU:HD21	2.35	0.56
2:B:41:VAL:HG21	2:B:128:VAL:HG21	1.87	0.56
2:B:141:ASN:ND2	2:B:209:GLN:HE21	2.04	0.56
2:A:35:GLU:CD	2:A:35:GLU:N	2.63	0.55
2:A:195:PRO:HG3	2:A:226:TRP:CE2	2.41	0.55
2:A:5:LYS:N	2:A:6:PRO:HD2	2.23	0.54
2:B:133:ILE:HD11	2:B:172:ASP:OD1	2.07	0.54
2:A:250:VAL:HG13	2:B:235:VAL:HG13	1.90	0.54
2:B:5:LYS:HB3	2:B:6:PRO:HD3	1.89	0.53
2:B:56:GLN:HE22	2:B:113:ALA:H	1.57	0.53
2:A:66:LYS:HG3	2:A:67:ASN:ND2	2.24	0.53
2:B:71:ILE:HG22	2:B:72:GLY:H	1.73	0.53
2:A:35:GLU:HB2	2:A:36:PRO:CD	2.32	0.53
2:A:11:ILE:HG12	2:A:44:PHE:HE2	1.73	0.52
2:A:132:ASN:HD22	2:A:134:SER:H	1.56	0.52
2:B:193:SER:HB3	2:B:215:LEU:CD1	2.40	0.52
2:A:256:TYR:HB2	2:B:231:LEU:HD23	1.91	0.52
1:E:8:DA:C1'	1:E:9:DA:H5'	2.35	0.51
1:E:8:DA:H2''	1:E:9:DA:O5'	2.09	0.51
2:B:195:PRO:HA	2:B:198:LEU:HD23	1.93	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:251:LYS:HD3	2:B:251:LYS:O	2.11	0.51
2:A:66:LYS:HZ2	2:A:67:ASN:HD21	1.54	0.51
1:E:9:DA:H62	2:B:141:ASN:ND2	2.09	0.50
2:A:205:ALA:HA	2:B:203:ALA:O	2.11	0.50
2:B:108:LYS:HG2	2:B:109:GLN:OE1	2.12	0.50
2:B:133:ILE:HG22	2:B:133:ILE:O	2.12	0.50
2:B:100:SER:HB3	2:B:103:ASN:HB3	1.94	0.50
2:B:14:ILE:C	2:B:14:ILE:HD12	2.36	0.49
2:B:108:LYS:HB3	2:B:111:ASP:CG	2.36	0.49
2:B:132:ASN:HB2	2:B:173:TRP:CZ2	2.47	0.49
2:B:198:LEU:N	2:B:198:LEU:HD22	2.28	0.49
2:B:4:ILE:HD11	2:B:126:LEU:HD22	1.90	0.49
2:B:42:TYR:CD2	2:B:55:LYS:HE3	2.47	0.49
2:A:7:ILE:O	2:A:11:ILE:HG13	2.12	0.49
2:A:256:TYR:CB	2:B:231:LEU:HD23	2.42	0.49
2:A:20:VAL:HG23	2:A:180:LEU:HD13	1.95	0.48
2:A:169:LEU:C	2:A:169:LEU:HD13	2.38	0.48
2:A:109:GLN:HG3	2:A:110:ASN:N	2.29	0.48
2:B:65:MET:HE3	2:B:101:ILE:HG23	1.96	0.47
2:A:200:ILE:HG12	2:A:208:ILE:HD13	1.95	0.47
1:F:7:DT:H2''	1:F:8:DA:O5'	2.15	0.47
2:B:14:ILE:HD12	2:B:15:LEU:HG	1.96	0.47
1:E:7:DT:H2''	1:E:8:DA:H5'	1.96	0.46
2:A:38:GLU:HG2	2:A:128:VAL:HG21	1.98	0.46
2:B:42:TYR:CG	2:B:55:LYS:HE3	2.51	0.46
2:A:132:ASN:C	2:A:132:ASN:ND2	2.72	0.46
2:B:15:LEU:O	2:B:182:CYS:HB3	2.15	0.46
2:B:131:ARG:HH12	2:B:213:ARG:HD2	1.81	0.46
2:B:4:ILE:HD13	2:B:126:LEU:HD22	1.96	0.45
2:A:127:ASP:HB3	2:A:168:TYR:CE1	2.52	0.45
2:B:170:GLU:HB3	2:B:186:SER:HB2	1.98	0.45
2:A:170:GLU:HB2	2:A:212:VAL:HB	1.99	0.45
2:A:202:TRP:HA	2:A:206:MET:HA	1.99	0.45
2:B:71:ILE:HG22	2:B:72:GLY:N	2.31	0.45
2:A:210:PHE:C	2:A:210:PHE:CD1	2.95	0.44
2:B:197:GLU:HB3	5:B:411:HOH:O	2.17	0.44
2:B:16:ILE:C	2:B:18:GLN:H	2.25	0.44
2:B:210:PHE:CD1	2:B:210:PHE:C	2.96	0.44
2:B:179:ASP:C	2:B:180:LEU:HD12	2.43	0.43
2:B:70:ILE:HG23	2:B:75:ALA:HB1	1.99	0.43
2:A:132:ASN:HD21	2:A:134:SER:HB3	1.84	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:8:DA:H2''	1:F:9:DA:O4'	2.18	0.42
2:A:19:LYS:HA	2:A:180:LEU:O	2.19	0.42
2:B:141:ASN:HD22	2:B:209:GLN:HE21	1.66	0.42
2:B:37:PHE:CD1	2:B:38:GLU:N	2.87	0.42
2:B:132:ASN:C	2:B:134:SER:H	2.27	0.42
2:B:64:PHE:CE1	2:B:76:ARG:HB3	2.54	0.42
5:E:43:HOH:O	2:B:136:SER:HB3	2.19	0.42
2:B:11:ILE:O	2:B:14:ILE:HG13	2.20	0.42
2:B:76:ARG:NH1	2:B:99:TRP:HB3	2.35	0.41
2:B:161:PHE:CZ	2:B:224:GLU:HG3	2.54	0.41
2:A:10:ASP:O	2:A:14:ILE:HG13	2.20	0.41
2:B:133:ILE:CG1	2:B:172:ASP:HB3	2.50	0.41
2:B:241:ARG:O	2:B:245:MET:HG3	2.20	0.41
2:A:4:ILE:C	2:A:4:ILE:HD12	2.46	0.41
2:B:170:GLU:HB2	2:B:212:VAL:CB	2.46	0.41
2:B:180:LEU:HD12	2:B:180:LEU:N	2.36	0.41
2:B:71:ILE:O	2:B:75:ALA:HB3	2.21	0.40
2:B:76:ARG:O	2:B:79:LEU:HB2	2.21	0.40
2:B:123:TYR:CD1	2:B:123:TYR:N	2.88	0.40
2:B:251:LYS:C	2:B:251:LYS:CD	2.89	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein backbone outliers to report in this entry.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
2	A	202/229 (88%)	194 (96%)	8 (4%)	28 14

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	199/229 (87%)	192 (96%)	7 (4%)	32	18
All	All	401/458 (88%)	386 (96%)	15 (4%)	30	16

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	A	35	GLU
2	A	109	GLN
2	A	110	ASN
2	A	132	ASN
2	A	133	ILE
2	A	148	LEU
2	A	169	LEU
2	A	231	LEU
2	B	86	LEU
2	B	109	GLN
2	B	116	LEU
2	B	127	ASP
2	B	215	LEU
2	B	231	LEU
2	B	247	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
2	A	12	ASN
2	A	67	ASN
2	A	98	ASN
2	A	103	ASN
2	A	109	GLN
2	A	110	ASN
2	A	132	ASN
2	A	220	ASN
2	B	9	GLN
2	B	56	GLN
2	B	67	ASN
2	B	98	ASN
2	B	110	ASN
2	B	141	ASN
2	B	167	ASN
2	B	211	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å²)	Q<0.9
1	E	14/14 (100%)	0.24	0	100	100	30, 41, 50, 57	0
1	F	14/14 (100%)	0.12	0	100	100	28, 40, 49, 51	0
2	A	245/257 (95%)	0.73	14 (5%)	29	32	27, 41, 58, 70	0
2	B	236/257 (91%)	1.09	36 (15%)	5	5	27, 50, 82, 99	0
All	All	509/542 (93%)	0.86	50 (9%)	13	14	27, 44, 67, 99	0

All (50) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	40	LEU	5.0
2	B	37	PHE	4.7
2	B	14	ILE	4.7
2	A	44	PHE	4.2
2	B	110	ASN	4.0
2	B	133	ILE	4.0
2	A	122	PHE	3.9
2	B	177	GLY	3.9
2	B	122	PHE	3.9
2	B	11	ILE	3.9
2	A	80	PHE	3.5
2	B	180	LEU	3.5
2	B	39	LYS	3.3
2	A	109	GLN	3.2
2	B	44	PHE	3.1
2	A	133	ILE	3.1
2	B	220	ASN	3.0
2	B	70	ILE	3.0
2	B	83	PRO	2.9
2	B	215	LEU	2.9
2	B	173	TRP	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	A	243	ILE	2.8
2	B	181	VAL	2.8
2	B	78	LYS	2.8
2	B	162	ASP	2.8
2	A	50	SER	2.8
2	A	81	ASN	2.8
2	A	175	LEU	2.6
2	B	104	LEU	2.6
2	B	100	SER	2.6
2	A	244	SER	2.6
2	B	16	ILE	2.5
2	B	41	VAL	2.5
2	B	51	ASP	2.5
2	A	7	ILE	2.5
2	B	132	ASN	2.4
2	B	99	TRP	2.4
2	B	105	PHE	2.3
2	A	51	ASP	2.3
2	B	12	ASN	2.3
2	B	43	LYS	2.2
2	A	134	SER	2.2
2	B	257	ILE	2.1
2	B	18	GLN	2.1
2	B	109	GLN	2.1
2	A	22	ARG	2.1
2	B	71	ILE	2.1
2	B	134	SER	2.1
2	B	15	LEU	2.1
2	B	86	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NA	F	403	1/1	0.96	0.05	38,38,38,38	0
4	CA	B	402	1/1	0.97	0.06	52,52,52,52	0
4	CA	A	401	1/1	0.99	0.04	36,36,36,36	0

6.5 Other polymers [i](#)

There are no such residues in this entry.