



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 11:24 AM UTC

PDB ID : 3GIJ / pdb_00003gij
Title : Dpo4 extension ternary complex with oxoG(syn)-A(anti) and oxoG(anti)-A(syn) pairs
Authors : Rechkoblit, O.; Malinina, L.; Patel, D.J.
Deposited on : 2009-03-05
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	341	61% 36%
1	B	341	57% 38%
2	D	13	62% 38%
2	H	13	15% 100%
3	E	18	11% 28% 61% 11%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	J	18	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	2DA	D	814[B]	-	-	X	-
3	8OG	E	906[A]	X	-	-	-
3	8OG	E	906[B]	X	-	-	-
3	8OG	J	1906	X	-	-	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7025 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase IV.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	341	Total	C	N	O	S	0	1	0
			2750	1763	475	506	6			
1	B	341	Total	C	N	O	S	0	0	0
			2740	1757	472	505	6			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	expression tag	UNP Q97W02
B	1001	GLY	-	expression tag	UNP Q97W02

- Molecule 2 is a DNA chain called 5'-D(*GP*TP*TP*GP*GP*AP*TP*GP*GP*TP*AP*GP*(2DA))-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	13	Total	C	N	O	P	0	1	0
			291	140	58	80	13			
2	H	13	Total	C	N	O	P	0	0	0
			271	130	53	76	12			

- Molecule 3 is a DNA chain called 5'-D(*CP*TP*AP*AP*CP*(8OG)P*CP*TP*AP*CP*CP*AP*TP*CP*CP*AP*AP*C)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	18	Total	C	N	O	P	0	1	0
			381	182	70	111	18			
3	J	14	Total	C	N	O	P	0	0	0
			280	133	50	83	14			

- Molecule 4 is 2'-DEOXYGUANOSINE-5'-TRIPHOSPHATE (CCD ID: DGT) (formula: C₁₀H₁₆N₅O₁₃P₃).



- Molecule 5 is CALCIUM ION (CCD ID: CA) (formula: Ca).

- Molecule 6 is water.

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	J	6	Total	O	0	0
			6	6		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: DNA polymerase IV



• Molecule 1: DNA polymerase IV




• Molecule 2: 5'-D(*GP*TP*TP*GP*GP*AP*TP*GP*GP*TP*AP*GP*(2DA))-3'

Chain D:  62% 38%



- Molecule 2: 5'-D(*GP*TP*TP*GP*GP*AP*TP*GP*GP*TP*AP*GP*(2DA))-3'

Chain H:  15% 100%




- Molecule 3: 5'-D(*CP*TP*AP*AP*CP*(8OG)P*CP*TP*AP*CP*CP*AP*TP*CP*CP*AP*A P*C)-3'

Chain E:  11% 28% 61% 11%



- Molecule 3: 5'-D(*CP*TP*AP*AP*CP*(8OG)P*CP*TP*AP*CP*CP*AP*TP*CP*CP*AP*A P*C)-3'

Chain J:  11% 50% 17% 11% 22%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	51.26Å 107.41Å 97.87Å 90.00° 100.54° 90.00°	Depositor
Resolution (Å)	20.00 – 2.40 20.00 – 2.40	Depositor EDS
% Data completeness (in resolution range)	94.8 (20.00-2.40) 94.6 (20.00-2.40)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.85 (at 2.35Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.209 , 0.253 0.205 , 0.250	Depositor DCC
R_{free} test set	2024 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	48.8	Xtriage
Anisotropy	0.482	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 62.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	7025	wwPDB-VP
Average B, all atoms (Å ²)	61.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.38% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, 8OG, DGT, 2DA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.66	0/2790	0.90	1/3746 (0.0%)
1	B	0.46	0/2779	0.80	0/3731
2	D	0.33	0/282	1.01	0/436
2	H	0.22	0/282	0.70	0/436
3	E	0.41	0/373	1.15	1/568 (0.2%)
3	J	0.27	0/285	0.83	1/432 (0.2%)
All	All	0.53	0/6791	0.87	3/9349 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	E	2	0
3	J	2	0
All	All	4	0

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	909	DA	O5'-C5'-C4'	-6.35	101.27	110.80
3	J	1907	DC	O5'-C5'-C4'	-5.79	102.11	110.80
1	A	97	GLU	N-CA-C	-5.08	106.34	112.54

All (4) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	E	906[A]	8OG	C1'

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atom
3	E	906[B]	8OG	C1'
3	J	1906	8OG	C4',C1'

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2750	0	2889	131	0
1	B	2740	0	2880	126	0
2	D	291	0	160	14	0
2	H	271	0	149	0	0
3	E	381	0	207	19	0
3	J	280	0	155	4	0
4	A	31	0	12	4	0
4	B	31	0	12	5	0
5	A	3	0	0	0	0
5	B	3	0	0	0	0
6	A	129	0	0	18	0
6	B	61	0	0	10	0
6	D	20	0	0	2	0
6	E	20	0	0	3	0
6	H	8	0	0	0	0
6	J	6	0	0	0	0
All	All	7025	0	6464	279	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 22.

The worst 5 of 279 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1116:ARG:H	1:B:1116:ARG:HE	1.04	1.00
1:A:14:GLN:HE22	1:A:139:THR:H	1.02	0.94
1:B:1014:GLN:HE22	1:B:1139:THR:H	1.16	0.93

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:130:ASN:O	1:A:134:GLU:HG3	1.71	0.90
1:B:1115:VAL:HG13	1:B:1116:ARG:HH21	1.36	0.89

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	340/341 (100%)	325 (96%)	9 (3%)	6 (2%)	6	9
1	B	339/341 (99%)	312 (92%)	20 (6%)	7 (2%)	5	7
All	All	679/682 (100%)	637 (94%)	29 (4%)	13 (2%)	6	8

5 of 13 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	1036	ARG
1	B	1117	ASP
1	A	303	PRO
1	B	1047	ASN
1	B	1115	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	300/299 (100%)	285 (95%)	15 (5%)	22	38
1	B	299/299 (100%)	287 (96%)	12 (4%)	28	47
All	All	599/598 (100%)	572 (96%)	27 (4%)	24	42

5 of 27 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	323	LEU
1	B	1116	ARG
1	B	1293	LEU
1	B	1023	LEU
1	B	1141	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	1047	ASN
1	B	1082	GLN
1	B	1304	HIS
1	A	161	ASN
1	A	285	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

6 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	2DA	H	1814	3,5,2	19,22,23	0.29	0	25,31,34	0.60	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	2DA	D	814[B]	5,2	19,22,23	0.27	0	25,31,34	0.50	0
3	8OG	J	1906	3,2	22,25,26	3.96	9 (40%)	26,37,40	3.16	15 (57%)
2	2DA	D	814[A]	3,5,2	19,22,23	0.39	0	25,31,34	0.73	0
3	8OG	E	906[B]	3	22,25,26	3.96	10 (45%)	26,37,40	3.24	14 (53%)
3	8OG	E	906[A]	3,2	22,25,26	4.07	7 (31%)	26,37,40	3.09	16 (61%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	2DA	H	1814	3,5,2	-	4/7/18/19	0/3/3/3
2	2DA	D	814[B]	5,2	-	4/7/18/19	0/3/3/3
3	8OG	J	1906	3,2	2/2/4/4	4/7/21/22	0/3/3/3
2	2DA	D	814[A]	3,5,2	-	3/7/18/19	0/3/3/3
3	8OG	E	906[B]	3	1/1/4/4	1/7/21/22	0/3/3/3
3	8OG	E	906[A]	3,2	1/1/4/4	1/7/21/22	0/3/3/3

The worst 5 of 26 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	906[A]	8OG	O3'-C3'	-10.25	1.21	1.43
3	J	1906	8OG	O3'-C3'	-9.94	1.22	1.43
3	E	906[B]	8OG	O3'-C3'	-9.81	1.22	1.43
3	E	906[B]	8OG	C5-C4	8.69	1.49	1.37
3	E	906[A]	8OG	C5-C4	8.69	1.49	1.37

The worst 5 of 45 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	906[B]	8OG	N9-C4-N3	9.85	138.44	126.13
3	J	1906	8OG	N9-C4-N3	8.23	136.41	126.13
3	E	906[A]	8OG	N9-C4-N3	7.74	135.81	126.13
3	E	906[B]	8OG	N7-C8-N9	5.61	112.85	106.61
3	E	906[A]	8OG	N7-C8-N9	5.46	112.69	106.61

All (4) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	E	906[A]	8OG	C1'
3	E	906[B]	8OG	C1'
3	J	1906	8OG	C4'
3	J	1906	8OG	C1'

5 of 17 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	814[A]	2DA	O4'-C1'-N9-C4
2	D	814[B]	2DA	O4'-C4'-C5'-O5'
2	H	1814	2DA	O4'-C4'-C5'-O5'
3	J	1906	8OG	C2'-C1'-N9-C8
3	J	1906	8OG	C2'-C1'-N9-C4

There are no ring outliers.

4 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	814[B]	2DA	7	0
2	D	814[A]	2DA	4	0
3	E	906[B]	8OG	1	0
3	E	906[A]	8OG	2	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 6 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	DGT	B	1414	5	32,33,33	0.79	3 (9%)	48,52,52	2.00	8 (16%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	DGT	A	414	5	32,33,33	0.71	1 (3%)	48,52,52	2.14	10 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	DGT	B	1414	5	-	8/22/34/34	0/3/3/3
4	DGT	A	414	5	-	6/22/34/34	0/3/3/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	1414	DGT	PB-O3A	2.25	1.61	1.59
4	B	1414	DGT	PA-O3A	2.18	1.61	1.59
4	A	414	DGT	PA-O3A	2.15	1.61	1.59
4	B	1414	DGT	PB-O3B	2.04	1.61	1.59

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	414	DGT	O1G-PG-O3G	-6.80	84.34	110.83
4	B	1414	DGT	O2G-PG-O3G	-6.70	84.71	110.83
4	B	1414	DGT	O1G-PG-O3G	-6.18	86.76	110.83
4	A	414	DGT	C2'-C1'-N9	-5.45	100.18	113.81
4	A	414	DGT	O3B-PG-O3G	-5.41	82.59	111.04

There are no chirality outliers.

5 of 14 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	B	1414	DGT	PB-O3B-PG-O2G
4	A	414	DGT	PB-O3B-PG-O3G
4	B	1414	DGT	O4'-C4'-C5'-O5'
4	B	1414	DGT	PB-O3B-PG-O3G
4	B	1414	DGT	PA-O3A-PB-O2B


There are no ring outliers.

2 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	1414	DGT	5	0
4	A	414	DGT	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	341/341 (100%)	0.34	4 (1%) 76 73	25, 58, 67, 75	1 (0%)
1	B	341/341 (100%)	0.59	4 (1%) 76 73	49, 64, 74, 81	0
2	D	12/13 (92%)	0.21	0 100 100	40, 46, 82, 85	0
2	H	12/13 (92%)	1.03	2 (16%) 4 3	54, 62, 99, 102	0
3	E	17/18 (94%)	0.89	2 (11%) 9 6	43, 58, 72, 75	0
3	J	13/18 (72%)	1.00	2 (15%) 5 4	54, 63, 93, 95	0
All	All	736/744 (98%)	0.49	14 (1%) 66 62	25, 60, 73, 102	1 (0%)

The worst 5 of 14 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	E	901	DC	3.1
3	E	902	DT	2.7
1	B	1141	THR	2.7
1	A	1	GLY	2.6
1	A	60	PRO	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	8OG	J	1906	23/24	0.68	0.19	57,61,65,66	0
2	2DA	H	1814	20/21	0.76	0.18	57,61,69,70	0
2	2DA	D	814[A]	20/21	0.84	0.19	54,60,61,63	20
2	2DA	D	814[B]	20/21	0.84	0.19	49,51,55,56	20

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	8OG	E	906[A]	23/24	0.88	0.14	49,55,60,62	23
3	8OG	E	906[B]	23/24	0.88	0.14	54,57,60,61	23

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	DGT	B	1414	31/31	0.82	0.19	50,53,57,58	0
5	CA	B	1417	1/1	0.85	0.21	69,69,69,69	0
4	DGT	A	414	31/31	0.90	0.14	32,38,40,41	0
5	CA	B	1415	1/1	0.94	0.20	68,68,68,68	0
5	CA	A	417	1/1	0.96	0.13	59,59,59,59	0
5	CA	B	1416	1/1	0.97	0.14	57,57,57,57	0
5	CA	A	415	1/1	0.98	0.23	57,57,57,57	0
5	CA	A	416	1/1	0.98	0.11	42,42,42,42	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around DGT B 1414:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around DGT A 414:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.