



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 12:29 PM UTC

PDB ID : 3GIP / pdb_00003gip
Title : Crystal structure of N-acyl-D-Glutamate Deacylase from Bordetella Bronchiseptica complexed with zinc, acetate and formate ions.
Authors : Fedorov, A.A.; Fedorov, E.V.; Cummings, J.; Raushel, F.M.; Almo, S.C.
Deposited on : 2009-03-05
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4037 (1.50-1.50)
Clashscore	190562	4235 (1.50-1.50)
Ramachandran outliers	187476	4153 (1.50-1.50)
Sidechain outliers	187428	4150 (1.50-1.50)
RSRZ outliers	180081	4039 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	480	3% 84% 14% ..
1	B	480	5% 82% 16% ..

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 7816 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called N-acyl-D-glutamate deacylase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	474	Total	C	N	O	S	0	0	0
			3555	2226	654	657	18			
1	B	474	Total	C	N	O	S	0	0	0
			3555	2226	654	657	18			

- Molecule 2 is ACETIC ACID (CCD ID: ACY) (formula: C₂H₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 3 is FORMIC ACID (CCD ID: FMT) (formula: CH₂O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			3	1	2		
3	A	1	Total	C	O	0	0
			3	1	2		
3	B	1	Total	C	O	0	0
			3	1	2		
3	B	1	Total	C	O	0	0
			3	1	2		

- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Zn	0	0
			1	1		
4	B	1	Total	Zn	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	351	Total	O	0	0
			351	351		
5	B	333	Total	O	0	0
			333	333		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: N-acyl-D-glutamate deacylase



- Molecule 1: N-acyl-D-glutamate deacylase



4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	90.99Å 90.99Å 507.13Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	24.95 – 1.50 24.95 – 1.50	Depositor EDS
% Data completeness (in resolution range)	95.4 (24.95-1.50) 95.5 (24.95-1.50)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.58 (at 1.50Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.205 , 0.215 0.204 , 0.215	Depositor DCC
R_{free} test set	9566 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	16.4	Xtriage
Anisotropy	0.563	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 34.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7816	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.48% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, FMT, ACY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.33	0/3636	0.93	15/4957 (0.3%)
1	B	0.35	0/3636	0.96	20/4957 (0.4%)
All	All	0.34	0/7272	0.94	35/9914 (0.4%)

There are no bond length outliers.

All (35) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	218	HIS	N-CA-C	-8.30	93.39	107.99
1	B	218	HIS	N-CA-C	-8.06	93.81	107.99
1	B	282	TYR	N-CA-C	-7.84	98.31	110.14
1	A	282	TYR	N-CA-C	-7.38	98.99	110.14
1	B	367	LEU	CA-C-N	7.38	127.09	119.56
1	B	367	LEU	C-N-CA	7.38	127.09	119.56
1	B	214	LEU	N-CA-C	6.94	120.80	110.52
1	A	374	HIS	N-CA-C	-6.80	101.19	109.83
1	A	214	LEU	N-CA-C	6.62	120.68	110.42
1	A	217	SER	N-CA-C	6.58	120.33	108.69
1	B	217	SER	N-CA-C	6.56	120.31	108.69
1	B	76	LYS	CA-C-N	6.44	126.94	119.47
1	B	76	LYS	C-N-CA	6.44	126.94	119.47
1	B	162	THR	N-CA-C	-6.36	102.59	110.41
1	A	367	LEU	CA-C-N	6.03	127.38	119.84
1	A	367	LEU	C-N-CA	6.03	127.38	119.84
1	A	17	ILE	N-CA-C	-5.80	97.50	106.72
1	A	154	MET	N-CA-C	5.73	117.86	108.52
1	A	18	ASP	N-CA-C	5.65	120.03	113.20
1	B	462	VAL	N-CA-C	-5.57	107.85	113.47
1	B	374	HIS	N-CA-C	-5.52	101.38	109.50
1	B	380	SER	N-CA-C	5.47	116.94	110.97

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	162	THR	N-CA-C	-5.47	103.48	110.53
1	A	12	THR	N-CA-C	5.46	118.59	110.14
1	A	462	VAL	N-CA-C	-5.33	108.31	113.53
1	B	160	ALA	N-CA-C	-5.33	103.32	109.93
1	B	17	ILE	N-CA-C	-5.28	99.08	106.53
1	B	469	ASP	CB-CA-C	-5.23	109.53	117.07
1	B	191	GLN	N-CA-C	-5.23	102.61	110.24
1	A	427	ASP	N-CA-C	-5.22	100.01	108.41
1	A	160	ALA	N-CA-C	-5.12	103.58	109.93
1	B	358	PRO	N-CA-C	5.11	123.00	112.47
1	B	469	ASP	CA-C-O	5.10	120.90	117.94
1	B	357	HIS	CA-C-N	5.07	126.17	119.84
1	B	357	HIS	C-N-CA	5.07	126.17	119.84

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3555	0	3504	43	0
1	B	3555	0	3504	53	0
2	A	4	0	3	0	0
2	B	4	0	3	0	0
3	A	6	0	2	0	0
3	B	6	0	2	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	351	0	0	4	0
5	B	333	0	0	2	0
All	All	7816	0	7018	94	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (94) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:252:MET:HG2	1:B:346:MET:HE2	1.39	1.04
1:A:16:ILE:HD13	1:A:58:VAL:HB	1.52	0.89
1:A:166:GLN:HE21	1:A:170:GLN:HE21	1.25	0.85
1:B:417:ARG:HH12	1:B:458:ASN:HD21	1.27	0.82
1:A:393:ARG:HH21	1:B:74:VAL:HG22	1.47	0.80
1:B:166:GLN:HE21	1:B:170:GLN:HE21	1.26	0.79
1:B:374:HIS:HD2	1:B:376:ARG:H	1.30	0.79
1:A:374:HIS:HD2	1:A:376:ARG:H	1.30	0.76
1:B:16:ILE:HD13	1:B:58:VAL:HB	1.69	0.74
1:B:139:ASN:HD22	1:B:474:GLN:H	1.36	0.72
1:A:98:SER:H	1:A:148:ASN:HD21	1.41	0.68
1:B:98:SER:H	1:B:148:ASN:HD21	1.42	0.68
1:B:174:GLN:O	1:B:178:GLU:HG3	1.93	0.67
1:A:417:ARG:HH12	1:A:458:ASN:HD21	1.42	0.67
1:A:174:GLN:O	1:A:178:GLU:HG3	1.94	0.67
1:A:271:ARG:HB3	1:A:277:VAL:HG12	1.76	0.67
1:B:345:ALA:C	1:B:346:MET:HE3	2.20	0.66
1:B:261:ARG:HG3	5:B:632:HOH:O	1.96	0.65
1:A:139:ASN:HD22	1:A:474:GLN:H	1.45	0.65
1:B:417:ARG:HH12	1:B:458:ASN:ND2	1.94	0.64
1:B:407:LEU:HB3	1:B:408:PRO:HD3	1.80	0.63
1:A:12:THR:HG23	5:A:804:HOH:O	2.00	0.61
1:A:374:HIS:CD2	1:A:376:ARG:H	2.16	0.59
1:B:253:MET:HE1	1:B:298:THR:HB	1.83	0.59
1:B:286:GLY:O	1:B:374:HIS:HE1	1.85	0.59
1:B:374:HIS:CD2	1:B:376:ARG:H	2.17	0.59
1:A:98:SER:H	1:A:148:ASN:ND2	2.00	0.59
1:B:436:VAL:HA	1:B:451:GLY:HA3	1.86	0.58
1:B:13:GLY:HA2	1:B:26:ARG:NH2	2.19	0.58
1:A:407:LEU:HB3	1:A:408:PRO:HD3	1.85	0.57
1:A:286:GLY:O	1:A:374:HIS:HE1	1.87	0.57
1:B:98:SER:H	1:B:148:ASN:ND2	2.03	0.56
1:A:417:ARG:HH12	1:A:458:ASN:ND2	2.05	0.55
1:B:304:ILE:HG21	1:B:313:CYS:SG	2.47	0.53
1:B:198:ALA:O	1:B:202:GLU:HG3	2.09	0.53
1:B:256:ASN:ND2	1:B:259:ARG:HH11	2.08	0.52
1:A:198:ALA:O	1:A:202:GLU:HG3	2.10	0.52
1:B:345:ALA:HB3	1:B:346:MET:CE	2.42	0.50
1:A:5:GLU:OE2	1:A:32:ARG:HD3	2.11	0.50
1:A:256:ASN:ND2	1:A:259:ARG:HH11	2.09	0.49
1:B:396:THR:OG1	1:B:399:GLN:HG3	2.12	0.49
1:A:168:ALA:O	1:A:172:MET:HG3	2.12	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:60:PRO:HG3	1:B:420:LEU:HG	1.95	0.49
1:B:345:ALA:CA	1:B:346:MET:HE3	2.41	0.49
1:A:393:ARG:HH21	1:B:74:VAL:CG2	2.21	0.49
1:A:253:MET:HE3	1:A:298:THR:O	2.13	0.49
1:A:146:HIS:HE1	1:A:200:GLU:OE2	1.95	0.48
1:B:417:ARG:NH1	1:B:458:ASN:HD21	2.06	0.48
1:A:101:PRO:HG3	1:A:172:MET:HE2	1.95	0.48
1:A:247:SER:O	1:A:248:HIS:C	2.57	0.47
1:A:281:ILE:C	1:A:281:ILE:HD12	2.40	0.47
1:A:13:GLY:HA2	1:A:26:ARG:NH2	2.30	0.46
1:B:313:CYS:SG	1:B:336:LEU:HD22	2.56	0.45
1:B:252:MET:O	1:B:253:MET:HB2	2.17	0.45
1:B:281:ILE:C	1:B:281:ILE:HD12	2.41	0.45
1:B:45:HIS:HD2	1:B:46:PRO:HD2	1.81	0.45
1:B:212:ARG:HH11	1:B:212:ARG:HG3	1.81	0.45
1:B:247:SER:O	1:B:248:HIS:C	2.59	0.45
1:B:29:LEU:C	1:B:29:LEU:HD23	2.42	0.44
1:B:269:ARG:O	1:B:272:GLU:HB2	2.18	0.44
1:A:393:ARG:NH2	5:A:512:HOH:O	2.48	0.44
1:B:128:PHE:HB3	1:B:477:ARG:NH2	2.32	0.44
1:A:29:LEU:HD12	1:A:39:ILE:HG12	1.98	0.44
1:A:253:MET:HA	1:A:254:PRO:HD3	1.91	0.44
1:A:96:GLY:O	1:A:147:ALA:HB3	2.18	0.43
1:B:405:THR:C	1:B:408:PRO:HD2	2.43	0.43
1:A:214:LEU:HD23	1:A:243:ALA:HB3	2.01	0.43
1:B:185:SER:HA	1:B:216:THR:O	2.18	0.43
1:A:152:ALA:CB	1:A:172:MET:HE1	2.49	0.43
1:B:7:LEU:HD23	1:B:32:ARG:NH2	2.34	0.43
1:B:256:ASN:HD22	1:B:259:ARG:HH11	1.67	0.43
1:A:252:MET:O	1:A:253:MET:HB2	2.19	0.42
1:B:383:ARG:O	1:B:387:ARG:HB3	2.19	0.42
1:B:310:HIS:O	1:B:313:CYS:SG	2.72	0.42
1:A:253:MET:HE1	1:A:298:THR:HB	2.00	0.42
1:A:60:PRO:HG3	1:A:420:LEU:HG	2.02	0.42
1:A:436:VAL:HA	1:A:451:GLY:HA3	2.02	0.42
1:B:146:HIS:HE1	1:B:200:GLU:OE2	2.02	0.42
5:A:527:HOH:O	1:B:76:LYS:HG2	2.20	0.42
1:B:253:MET:HE3	1:B:298:THR:O	2.20	0.42
1:B:345:ALA:HB3	1:B:346:MET:HE3	2.01	0.42
1:A:269:ARG:O	1:A:273:GLN:HG3	2.20	0.41
1:A:271:ARG:HB3	1:A:277:VAL:CG1	2.46	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:139:ASN:ND2	1:B:474:GLN:H	2.10	0.41
1:A:25:ARG:NH1	5:A:634:HOH:O	2.52	0.41
1:A:103:PRO:HD3	1:A:120:LEU:HD12	2.02	0.41
1:B:146:HIS:HD2	5:B:507:HOH:O	2.03	0.41
1:B:202:GLU:O	1:B:206:ARG:HG3	2.20	0.41
1:A:152:ALA:HB1	1:A:172:MET:HE1	2.02	0.40
1:B:372:ARG:HG2	1:B:372:ARG:HH11	1.86	0.40
1:B:70:ASP:HB3	1:B:99:ALA:HB2	2.03	0.40
1:A:389:VAL:O	1:A:393:ARG:HD3	2.21	0.40
1:A:75:GLU:CD	1:A:118:THR:HG1	2.29	0.40
1:B:48:ARG:HG3	1:B:48:ARG:HH11	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	472/480 (98%)	457 (97%)	13 (3%)	2 (0%)	30	12
1	B	472/480 (98%)	456 (97%)	14 (3%)	2 (0%)	30	12
All	All	944/960 (98%)	913 (97%)	27 (3%)	4 (0%)	30	12

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	253	MET
1	B	253	MET
1	A	248	HIS
1	B	248	HIS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	352/355 (99%)	351 (100%)	1 (0%)	86	75
1	B	352/355 (99%)	350 (99%)	2 (1%)	78	62
All	All	704/710 (99%)	701 (100%)	3 (0%)	84	71

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	9	PHE
1	B	117	GLU
1	B	346	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (21) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	134	GLN
1	A	139	ASN
1	A	146	HIS
1	A	148	ASN
1	A	158	GLN
1	A	170	GLN
1	A	197	GLN
1	A	256	ASN
1	A	374	HIS
1	A	458	ASN
1	A	465	GLN
1	B	49	HIS
1	B	134	GLN
1	B	139	ASN
1	B	146	HIS
1	B	148	ASN
1	B	170	GLN
1	B	256	ASN
1	B	374	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	421	GLN
1	B	458	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 8 ligands modelled in this entry, 2 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	FMT	A	482	-	2,2,2	0.76	0	1,1,1	0.64	0
3	FMT	B	483	-	2,2,2	0.78	0	1,1,1	0.64	0
2	ACY	B	481	4	3,3,3	0.45	0	3,3,3	0.83	0
2	ACY	A	481	4	3,3,3	0.43	0	3,3,3	0.82	0
3	FMT	A	483	-	2,2,2	0.79	0	1,1,1	0.65	0
3	FMT	B	482	-	2,2,2	0.77	0	1,1,1	0.63	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	474/480 (98%)	0.15	14 (2%) 52 56	12, 17, 28, 41	0
1	B	474/480 (98%)	0.19	24 (5%) 33 36	11, 17, 31, 50	0
All	All	948/960 (98%)	0.17	38 (4%) 42 46	11, 17, 30, 50	0

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	313	CYS	7.1
1	B	469	ASP	4.9
1	B	5	GLU	4.9
1	A	45	HIS	4.7
1	B	44	ALA	4.3
1	A	12	THR	4.3
1	A	74	VAL	3.9
1	B	46	PRO	3.6
1	A	369	ASN	3.6
1	A	469	ASP	3.6
1	A	273	GLN	3.6
1	B	45	HIS	3.5
1	B	358	PRO	3.2
1	B	410	ARG	3.2
1	B	357	HIS	3.0
1	B	41	GLU	2.9
1	B	393	ARG	2.9
1	B	15	TRP	2.8
1	A	261	ARG	2.8
1	B	47	ALA	2.7
1	A	108	THR	2.6
1	B	8	ASP	2.6
1	A	116	GLY	2.5
1	A	155	ARG	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	7	LEU	2.4
1	B	26	ARG	2.3
1	A	468	ALA	2.3
1	A	393	ARG	2.3
1	A	5	GLU	2.3
1	B	6	LYS	2.2
1	B	117	GLU	2.2
1	B	42	LEU	2.2
1	B	48	ARG	2.2
1	B	212	ARG	2.2
1	B	108	THR	2.2
1	B	465	GLN	2.2
1	B	261	ARG	2.1
1	A	197	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	ACY	A	481	4/4	0.96	0.08	19,20,20,21	0
3	FMT	A	483	3/3	0.96	0.06	17,17,18,18	0
3	FMT	B	483	3/3	0.96	0.06	15,15,15,16	0
2	ACY	B	481	4/4	0.97	0.07	15,15,16,18	0
3	FMT	B	482	3/3	0.98	0.04	13,13,13,15	0
3	FMT	A	482	3/3	0.98	0.04	13,13,14,14	0
4	ZN	A	484	1/1	0.99	0.02	14,14,14,14	0
4	ZN	B	484	1/1	0.99	0.02	13,13,13,13	0

6.5 Other polymers [i](#)

There are no such residues in this entry.