



Full wwPDB NMR Structure Validation Report ⓘ

Mar 10, 2026 – 11:01 AM UTC

PDB ID : 1GJ2 / pdb_00001gj2
Title : CO(III)-BLEOMYCIN-OOH BOUND TO AN OLIGONUCLEOTIDE CONTAINING A PHOSPHOGLYCOLATE LESION
Authors : Hoehn, S.T.; Junker, H.-D.; Bunt, R.C.; Turner, C.J.; Stubbe, J.
Deposited on : 2000-11-01

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR



The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	6	 100%
2	B	6	 100%
3	C	13	 54% 15% 31%

2 Ensemble composition and analysis ⓘ

This entry contains 1 models. Identification of well-defined residues and clustering analysis are not possible.

3 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 988 atoms, of which 371 are hydrogens and 0 are deuteriums.

- Molecule 1 is a DNA chain called 5'-D(*CP*CP*AP*AP*AP*G)-3'.

Mol	Chain	Residues	Atoms						Trace
1	A	6	Total	C	H	N	O	P	0
			187	58	67	26	31	5	


- Molecule 2 is a DNA chain called 5'-D(P*AP*CP*TP*GP*GP*G)-3'.

Mol	Chain	Residues	Atoms						Trace
2	B	6	Total	C	H	N	O	P	0
			194	59	68	25	36	6	

- Molecule 3 is a DNA chain called 5'-D(*CP*CP*CP*AP*GP*TP*AP*CP*TP*TP*TP*GP*G)-3'.

Mol	Chain	Residues	Atoms						Trace
3	C	13	Total	C	H	N	O	P	0
			410	126	149	45	78	12	

- Molecule 4 is 2-PHOSPHOGLYCOLIC ACID (CCD ID: PGA) (formula: C₂H₅O₆P).



Mol	Chain	Residues	Atoms				
4	A	1	Total	C	H	O	P
			10	2	2	5	1


- Molecule 5 is COBALT (III) ION (CCD ID: 3CO) (formula: Co).

Mol	Chain	Residues	Atoms	
5	B	1	Total 1	Co 1

- Molecule 6 is OXYGEN ATOM (CCD ID: O) (formula: O).


Mol	Chain	Residues	Atoms
6	B	1	Total O 1 1

- Molecule 7 is BLEOMYCIN B2 (CCD ID: BLB) (formula: $C_{55}H_{85}N_{20}O_{21}S_2$).



Mol	Chain	Residues	Atoms					
7	C	1	Total	C	H	N	O	S
			182	55	84	20	21	2

- Molecule 8 is HYDROGEN PEROXIDE (CCD ID: PEO) (formula: H_2O_2).



Mol	Chain	Residues	Atoms		
			Total	H	O
8	C	1	3	1	2

4 Residue-property plots

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: 5'-D(*CP*CP*AP*AP*AP*G)-3'

Chain A:  100%

There are no outlier residues in this chain.


- Molecule 2: 5'-D(P*AP*CP*TP*GP*GP*G)-3'

Chain B:  100%

There are no outlier residues in this chain.

- Molecule 3: 5'-D(*CP*CP*CP*AP*GP*TP*AP*CP*TP*TP*TP*GP*G)-3'

Chain C:  54% 15% 31%



5 Refinement protocol and experimental data overview ⓘ

The models were refined using the following method: *molecular dynamics*.

Of the ? calculated structures, 1 were deposited, based on the following criterion: ?.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR	structure solution	3.851
X-PLOR	refinement	3.851

No chemical shift data was provided.

6 Model quality

6.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: O, 3CO, BLB, PGA, PEO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.64	0/135 (0.0%)	1.36	0/206 (0.0%)
2	B	0.69	0/141 (0.0%)	1.41	0/216 (0.0%)
3	C	0.67	0/291 (0.0%)	1.88	9/447 (2.0%)
All	All	0.66	0/567 (0.0%)	1.66	9/869 (1.0%)

There are no bond-length outliers.

All angle outliers are listed below. They are sorted according to the Z-score.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	24	DT	C6-C5-C7	-9.81	109.29	124.00
3	C	15	DC	P-O5'-C5'	9.25	133.88	120.00
3	C	22	DT	C6-C5-C7	-9.14	110.29	124.00
3	C	22	DT	P-O3'-C3'	7.65	131.67	120.20
3	C	21	DC	P-O3'-C3'	6.31	129.67	120.20
3	C	23	DT	C6-C5-C7	-5.47	115.79	124.00
3	C	24	DT	C4-C5-C7	5.30	130.35	122.40
3	C	23	DT	N1-C1'-C2'	5.29	121.44	113.50
3	C	24	DT	P-O3'-C3'	5.28	128.12	120.20

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
3	C	261	149	149	3
All	All	617	371	367	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)
3:C:22:DT:H2'	3:C:23:DT:H72	0.46	1.88
3:C:23:DT:C2'	3:C:24:DT:H72	0.42	2.44
3:C:15:DC:C2	3:C:16:DC:C5	0.42	3.07

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

6.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
7	BLB	C	1	5	101,103,103	1.12	6 (5%)
4	PGA	A	7	1	4,7,8	1.02	0 (0%)
8	PEO	C	11	5	1,1,1	0.35	0 (0%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
7	BLB	C	1	5	126,145,145	2.42	36 (28%)
4	PGA	A	7	1	4,8,11	1.44	1 (25%)
8	PEO	C	11	5	-	-	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	BLB	C	1	5	-	0,103,144,144	0,6,6,6
4	PGA	A	7	1	-	0,3,5,6	-

All bond outliers are listed below. They are sorted according to the Z-score.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	C	1	BLB	C48-C49	5.42	1.38	1.48
7	C	1	BLB	C43-NN	3.44	1.35	1.31
7	C	1	BLB	C10-C12	3.24	1.45	1.50
7	C	1	BLB	C27-NJ	3.14	1.33	1.39

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	C	1	BLB	C47-C48	2.76	1.41	1.36
7	C	1	BLB	C44-C45	2.52	1.40	1.36

All angle outliers are listed below. They are sorted according to the Z-score.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	C	1	BLB	C63-OH1-C14	10.26	102.14	115.23
7	C	1	BLB	O70-C70-NQ	6.55	115.25	125.58
7	C	1	BLB	CB-C31-NK	6.42	98.67	109.70
7	C	1	BLB	O68-C70-NQ	5.71	119.76	110.92
7	C	1	BLB	C34-C36-NL	5.69	107.93	116.35
7	C	1	BLB	CC-C34-C36	5.47	119.60	108.77
7	C	1	BLB	C7-C6-NC	4.81	103.92	112.14
7	C	1	BLB	C5-C6-NC	4.78	118.26	109.25
7	C	1	BLB	O12-C12-C10	4.58	128.05	120.75
7	C	1	BLB	O62-C60-C59	4.28	117.42	109.70
7	C	1	BLB	C12-C10-NG	4.27	111.83	116.02
7	C	1	BLB	C61-C60-C59	4.27	123.50	113.02
7	C	1	BLB	C33-C34-C36	4.27	103.12	109.53
7	C	1	BLB	OH3-C38-C37	3.77	116.66	109.07
7	C	1	BLB	OH2-C33-C34	3.73	117.56	108.59
7	C	1	BLB	C30-C13-NH	3.61	120.14	110.32
7	C	1	BLB	C64-O56-C57	3.60	109.45	117.98
7	C	1	BLB	C13-NH-C12	3.59	114.28	121.53
7	C	1	BLB	C63-C57-C58	3.53	116.14	110.37
7	C	1	BLB	O56-C57-C58	3.24	115.48	107.23
7	C	1	BLB	C13-C30-NK	3.15	109.59	116.39
7	C	1	BLB	C58-C59-C60	3.12	115.89	110.23
7	C	1	BLB	O62-C63-C57	3.07	103.53	109.49
7	C	1	BLB	C47-C48-NO	3.01	111.29	115.43
7	C	1	BLB	C47-C48-C49	2.95	128.83	123.92
7	C	1	BLB	C9-C8-NF	2.91	117.92	122.22
7	C	1	BLB	C48-C49-NP	2.69	111.60	115.56
7	C	1	BLB	C45-C44-S43	2.67	107.15	110.24
4	A	7	PGA	O1P-C2-C1	2.66	114.55	110.54
7	C	1	BLB	C31-NK-C30	2.54	118.30	122.67
7	C	1	BLB	O36-C36-NL	2.50	127.43	122.96
7	C	1	BLB	O30-C30-NK	2.43	127.31	122.96
7	C	1	BLB	C44-S43-C43	2.41	92.21	89.59
7	C	1	BLB	C14-C13-NH	2.40	112.78	110.20
7	C	1	BLB	C42-C41-NM	2.28	107.14	112.00
7	C	1	BLB	O68-C68-C67	2.15	112.71	107.76

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	C	1	BLB	C50-NP-C49	2.15	127.17	122.47

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided