



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 10:58 AM UTC

PDB ID : 2GJK / pdb_00002gjk
Title : Structural and functional insights into the human Upf1 helicase core
Authors : Cheng, Z.; Muhlrاد, D.; Parker, R.; Song, H.
Deposited on : 2006-03-31
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	624	 6% 83% 14% ..

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4986 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Regulator of nonsense transcripts 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	614	Total	C	N	O	S	0	0	0
			4844	3075	849	897	23			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	291	PRO	-	cloning artifact	UNP Q92900
A	292	LEU	-	cloning artifact	UNP Q92900
A	293	GLY	-	cloning artifact	UNP Q92900
A	294	SER	-	cloning artifact	UNP Q92900

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (CCD ID: ANP) (formula: C₁₀H₁₇N₆O₁₂P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	31	10	6	12	3	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	110	Total	O	0	0
			110	110		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Regulator of nonsense transcripts 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	188.98Å 188.98Å 44.87Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.60 20.00 – 2.60	Depositor EDS
% Data completeness (in resolution range)	98.8 (20.00-2.60) 98.5 (20.00-2.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.37 (at 2.59Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.245 , 0.271 0.242 , 0.256	Depositor DCC
R_{free} test set	1227 reflections (4.84%)	wwPDB-VP
Wilson B-factor (Å ²)	45.5	Xtriage
Anisotropy	0.102	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 37.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	4986	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.34% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ANP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.47	1/4939 (0.0%)	0.78	0/6682

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	648	VAL	CA-CB	5.05	1.57	1.54

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	371	ALA	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4844	0	4910	37	0
2	A	1	0	0	0	0
3	A	31	0	13	0	0
4	A	110	0	0	1	0
All	All	4986	0	4923	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:796:TYR:OH	1:A:843:ARG:HD2	1.50	1.11
1:A:843:ARG:HG3	1:A:853:LEU:CD1	2.12	0.80
1:A:843:ARG:HG3	1:A:853:LEU:HD13	1.71	0.73
1:A:616:THR:HG22	1:A:619:GLY:H	1.54	0.72
1:A:796:TYR:OH	1:A:843:ARG:CD	2.35	0.70
1:A:796:TYR:HH	1:A:843:ARG:HD2	1.61	0.64
1:A:744:LYS:HE2	1:A:895:GLN:HE21	1.64	0.62
1:A:625:LEU:HA	1:A:628:MET:HE3	1.84	0.59
1:A:558:PHE:CE1	1:A:559:LEU:HD13	2.38	0.58
1:A:482:VAL:HG13	1:A:658:ILE:HD12	1.87	0.56
1:A:523:SER:OG	1:A:830:GLN:NE2	2.38	0.55
1:A:708:LEU:HD22	1:A:870:VAL:HG12	1.88	0.55
1:A:524:ASN:HD22	1:A:543:ARG:HH22	1.54	0.55
1:A:489:LEU:HG	1:A:693:ILE:HD11	1.88	0.55
1:A:571:MET:N	1:A:572:PRO:HD3	2.23	0.54
1:A:843:ARG:HB2	4:A:26:HOH:O	2.07	0.54
1:A:507:TYR:HA	1:A:539:LEU:HD11	1.92	0.52
1:A:719:GLY:O	1:A:722:GLN:NE2	2.41	0.52
1:A:579:GLN:O	1:A:583:GLU:HG2	2.10	0.51
1:A:772:ASN:O	1:A:776:ILE:HG12	2.13	0.49
1:A:755:GLU:OE1	1:A:843:ARG:CZ	2.61	0.48
1:A:438:SER:HB3	1:A:655:LYS:HA	1.96	0.47
1:A:507:TYR:O	1:A:511:ARG:HG2	2.15	0.47
1:A:725:VAL:HG13	1:A:729:ASP:HB2	1.97	0.46
1:A:371:ALA:C	1:A:373:LEU:H	2.25	0.45
1:A:629:GLN:HG3	1:A:653:GLY:HA3	1.97	0.45
1:A:521:ALA:HB3	1:A:527:VAL:HG23	1.99	0.44
1:A:633:ILE:HD12	1:A:654:ALA:HB2	2.00	0.44
1:A:764:TYR:CD2	1:A:843:ARG:NH1	2.69	0.43
1:A:374:TRP:CG	1:A:375:LYS:N	2.86	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:440:TYR:CE2	1:A:486:PRO:HG2	2.54	0.43
1:A:731:VAL:HG13	1:A:731:VAL:O	2.19	0.43
1:A:454:ILE:HD12	1:A:454:ILE:N	2.34	0.42
1:A:607:LEU:HB3	1:A:628:MET:HE1	2.01	0.42
1:A:554:SER:H	1:A:557:SER:HB3	1.85	0.42
1:A:711:PHE:HB3	1:A:712:PRO:HD3	2.01	0.40
1:A:489:LEU:HD22	1:A:659:LEU:HB2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	610/624 (98%)	563 (92%)	40 (7%)	7 (1%)	11	25

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	370	LEU
1	A	399	VAL
1	A	407	HIS
1	A	573	GLU
1	A	405	VAL
1	A	882	GLN
1	A	906	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	531/540 (98%)	485 (91%)	46 (9%)	9 21

All (46) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	309	LEU
1	A	313	GLU
1	A	319	LYS
1	A	328	ASN
1	A	329	ILE
1	A	337	LEU
1	A	338	ASN
1	A	340	LYS
1	A	349	LYS
1	A	367	LYS
1	A	374	TRP
1	A	375	LYS
1	A	395	LEU
1	A	399	VAL
1	A	407	HIS
1	A	408	ASN
1	A	432	VAL
1	A	450	VAL
1	A	451	GLU
1	A	466	GLN
1	A	482	VAL
1	A	545	CYS
1	A	559	LEU
1	A	565	ILE
1	A	580	LEU
1	A	592	GLU
1	A	598	LEU
1	A	607	LEU
1	A	616	THR
1	A	629	GLN
1	A	631	ARG
1	A	637	GLU
1	A	658	ILE
1	A	669	VAL
1	A	673	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	684	LEU
1	A	693	ILE
1	A	725	VAL
1	A	774	GLU
1	A	816	LYS
1	A	837	ILE
1	A	841	CYS
1	A	881	LYS
1	A	882	GLN
1	A	886	ASN
1	A	904	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	301	GLN
1	A	304	ASN
1	A	326	GLN
1	A	386	ASN
1	A	475	GLN
1	A	536	GLN
1	A	629	GLN
1	A	665	GLN
1	A	738	GLN
1	A	830	GLN
1	A	848	GLN
1	A	854	ASN
1	A	860	ASN
1	A	882	GLN
1	A	886	ASN
1	A	895	GLN
1	A	904	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ANP	A	990	2	33,33,33	2.71	11 (33%)	45,52,52	1.86	11 (24%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ANP	A	990	2	-	5/18/38/38	0/3/3/3

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	990	ANP	PG-O1G	8.36	1.58	1.46
3	A	990	ANP	PB-O1B	7.71	1.57	1.46
3	A	990	ANP	C4-N3	5.35	1.44	1.34
3	A	990	ANP	PB-N3B	3.90	1.73	1.63
3	A	990	ANP	PG-N3B	3.66	1.72	1.63
3	A	990	ANP	C5-C4	3.47	1.45	1.39
3	A	990	ANP	C8-N7	3.13	1.37	1.31
3	A	990	ANP	C5-C6	2.81	1.48	1.41
3	A	990	ANP	C4-N9	-2.51	1.32	1.37
3	A	990	ANP	C5-N7	-2.43	1.34	1.39
3	A	990	ANP	PA-O1A	2.24	1.58	1.50

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	990	ANP	C4-C5-N7	-4.31	105.65	110.58
3	A	990	ANP	C5-C4-N9	4.03	110.21	105.81
3	A	990	ANP	N3-C2-N1	-4.01	122.51	128.58
3	A	990	ANP	C6-C5-N7	3.98	139.77	132.09
3	A	990	ANP	C5-C4-N3	-3.74	121.56	126.72
3	A	990	ANP	C2-N1-C6	3.04	123.72	118.73
3	A	990	ANP	O1G-PG-N3B	-2.71	107.78	111.77
3	A	990	ANP	C2-N3-C4	2.67	118.35	111.83
3	A	990	ANP	O2B-PB-O3A	2.61	113.36	104.64
3	A	990	ANP	C3'-C2'-C1'	2.28	105.77	101.46
3	A	990	ANP	C5-N7-C8	2.06	106.68	103.45

There are no chirality outliers.


All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	990	ANP	PB-N3B-PG-O1G
3	A	990	ANP	PG-N3B-PB-O1B
3	A	990	ANP	PG-N3B-PB-O3A
3	A	990	ANP	PA-O3A-PB-O2B
3	A	990	ANP	PA-O3A-PB-O1B

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	614/624 (98%)	0.34	39 (6%) 25 20	15, 40, 69, 77	0

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	337	LEU	5.7
1	A	843	ARG	5.1
1	A	405	VAL	4.9
1	A	373	LEU	3.8
1	A	335	LEU	3.7
1	A	371	ALA	3.7
1	A	406	THR	3.6
1	A	905	ASN	3.6
1	A	903	LEU	3.4
1	A	366	TYR	3.2
1	A	333	TRP	3.2
1	A	513	GLY	3.2
1	A	547	LYS	3.2
1	A	908	GLU	3.1
1	A	334	ASP	3.0
1	A	342	ILE	3.0
1	A	370	LEU	3.0
1	A	906	LEU	3.0
1	A	551	ALA	2.8
1	A	364	LEU	2.8
1	A	403	VAL	2.7
1	A	352	SER	2.7
1	A	809	PHE	2.7
1	A	408	ASN	2.6
1	A	842	VAL	2.5
1	A	472	ASN	2.4
1	A	399	VAL	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	737	PHE	2.4
1	A	434	GLU	2.3
1	A	372	PRO	2.3
1	A	902	PRO	2.3
1	A	844	ALA	2.3
1	A	577	LEU	2.3
1	A	580	LEU	2.3
1	A	291	PRO	2.2
1	A	743	ASP	2.2
1	A	331	VAL	2.2
1	A	742	PRO	2.1
1	A	413	PHE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	MG	A	1000	1/1	0.89	0.22	50,50,50,50	0
3	ANP	A	990	31/31	0.91	0.14	38,49,51,52	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around ANP A 990:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.