



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 5, 2026 – 12:35 PM UTC

PDB ID : 3GJ2 / pdb_00003gj2
Title : Photoactivated state of PA-GFP
Authors : Henderson, J.N.; Gepshtein, R.; Heenan, J.R.; Kallio, K.; Huppert, D.; Remington, S.J.
Deposited on : 2009-03-07
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	229	2% 90% 8% .
1	B	229	% 92% 8%
1	C	229	5% 86% 12% ..
1	D	229	6% 90% 8% ..

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 7614 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Green fluorescent protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	227	Total	C	N	O	S	0	4	0
			1802	1146	308	343	5			
1	B	229	Total	C	N	O	S	0	7	0
			1848	1176	307	359	6			
1	C	225	Total	C	N	O	S	0	2	0
			1733	1102	294	332	5			
1	D	226	Total	C	N	O	S	0	1	0
			1737	1101	292	339	5			

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP P42212
A	?	-	SER	chromophore	UNP P42212
A	80	ARG	GLN	engineered mutation	UNP P42212
A	99	SER	PHE	engineered mutation	UNP P42212
A	153	THR	MET	engineered mutation	UNP P42212
A	163	ALA	VAL	engineered mutation	UNP P42212
A	203	HIS	THR	engineered mutation	UNP P42212
B	0	SER	-	expression tag	UNP P42212
B	?	-	SER	chromophore	UNP P42212
B	80	ARG	GLN	engineered mutation	UNP P42212
B	99	SER	PHE	engineered mutation	UNP P42212
B	153	THR	MET	engineered mutation	UNP P42212
B	163	ALA	VAL	engineered mutation	UNP P42212
B	203	HIS	THR	engineered mutation	UNP P42212
C	0	SER	-	expression tag	UNP P42212
C	?	-	SER	chromophore	UNP P42212
C	80	ARG	GLN	engineered mutation	UNP P42212
C	99	SER	PHE	engineered mutation	UNP P42212
C	153	THR	MET	engineered mutation	UNP P42212
C	163	ALA	VAL	engineered mutation	UNP P42212
C	203	HIS	THR	engineered mutation	UNP P42212

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	0	SER	-	expression tag	UNP P42212
D	?	-	SER	chromophore	UNP P42212
D	80	ARG	GLN	engineered mutation	UNP P42212
D	99	SER	PHE	engineered mutation	UNP P42212
D	153	THR	MET	engineered mutation	UNP P42212
D	163	ALA	VAL	engineered mutation	UNP P42212
D	203	HIS	THR	engineered mutation	UNP P42212

- Molecule 2 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Cl 1 1	0	0
2	B	1	Total Cl 1 1	0	0
2	C	1	Total Cl 1 1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	152	Total O 152 152	0	3
3	B	165	Total O 165 165	0	4
3	C	77	Total O 77 77	0	0
3	D	97	Total O 97 97	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Green fluorescent protein





- Molecule 1: Green fluorescent protein



- Molecule 1: Green fluorescent protein



- Molecule 1: Green fluorescent protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	84.71Å 87.07Å 144.86Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.66 – 1.90 40.66 – 1.90	Depositor EDS
% Data completeness (in resolution range)	97.7 (40.66-1.90) 97.9 (40.66-1.90)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.08 (at 1.89Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.205 , 0.248 0.202 , 0.247	Depositor DCC
R_{free} test set	4183 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	24.5	Xtriage
Anisotropy	0.065	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 38.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.008 for k,h,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7614	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.38% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, CRO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.36	3/1834 (0.2%)	1.20	4/2484 (0.2%)
1	B	1.36	2/1889 (0.1%)	1.17	2/2555 (0.1%)
1	C	1.23	1/1759 (0.1%)	1.09	0/2388
1	D	1.24	5/1759 (0.3%)	1.14	1/2391 (0.0%)
All	All	1.30	11/7241 (0.2%)	1.15	7/9818 (0.1%)

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	12	VAL	CA-CB	7.14	1.59	1.54
1	A	54	PRO	CA-C	6.33	1.58	1.52
1	B	51	GLY	N-CA	5.84	1.53	1.45
1	D	99	SER	CA-C	-5.77	1.46	1.53
1	B	169	HIS	CA-C	5.62	1.59	1.52
1	D	12	VAL	CA-C	-5.59	1.48	1.52
1	D	214	LYS	CA-C	-5.34	1.45	1.52
1	D	55	VAL	N-CA	-5.29	1.42	1.46
1	C	32	GLU	CA-C	-5.17	1.46	1.52
1	A	57	TRP	CA-C	5.17	1.59	1.52
1	A	163	ALA	C-O	-5.01	1.17	1.23

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	134	GLY	N-CA-C	-5.99	103.42	112.41
1	B	189	GLY	N-CA-C	-5.38	104.37	112.60
1	A	195	LEU	CA-C-N	-5.27	114.91	120.66
1	A	195	LEU	C-N-CA	-5.27	114.91	120.66
1	A	73	ARG	CG-CD-NE	-5.17	100.63	112.00
1	D	5	GLU	CB-CA-C	-5.02	102.78	110.81

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	184	GLN	CA-CB-CG	5.00	124.11	114.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1802	0	1720	6	0
1	B	1848	0	1770	6	0
1	C	1733	0	1595	22	0
1	D	1737	0	1585	10	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
3	A	152	0	0	0	0
3	B	165	0	0	0	0
3	C	77	0	0	0	0
3	D	97	0	0	0	0
All	All	7614	0	6670	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:65:CRO:C3	1:C:68:VAL:N	1.72	1.48
1:C:65:CRO:C3	1:C:68:VAL:CA	2.58	0.81
1:B:73:ARG:HB3	1:B:225[B]:THR:HG22	1.72	0.72
1:C:65:CRO:CA3	1:C:68:VAL:N	2.54	0.67
1:A:158:LYS:HB3	1:A:186[B]:THR:HG22	1.84	0.59
1:C:98:ILE:HG12	1:C:181:HIS:CD2	2.38	0.58
1:B:53:LEU:HD21	1:B:60:LEU:HD12	1.87	0.56
1:A:45:LYS:HE2	1:A:47:ILE:HD11	1.90	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:22:VAL:O	1:D:25:HIS:HB2	2.09	0.51
1:D:63:THR:CG2	1:D:123:ILE:HG21	2.40	0.51
1:C:56:PRO:HG2	1:C:141:LEU:HD13	1.93	0.49
1:C:167:ILE:CG1	1:C:181:HIS:CE1	2.95	0.49
1:C:65:CRO:HD1	1:C:65:CRO:N2	2.27	0.48
1:C:110:ALA:HA	1:C:122:ARG:O	2.13	0.48
1:C:43[A]:THR:HG23	1:C:221:LEU:HD11	1.96	0.48
1:B:128:ILE:O	1:B:129:ASP:HB2	2.14	0.47
1:C:167:ILE:HG13	1:C:181:HIS:CE1	2.50	0.46
1:C:220:LEU:O	1:C:221:LEU:HD12	2.15	0.46
1:B:120:VAL:HG12	1:B:122:ARG:HG3	1.98	0.46
1:A:209:LYS:O	1:A:211:PRO:HD3	2.16	0.46
1:D:161:ILE:CG1	1:D:185:ASN:HB2	2.46	0.45
1:C:125:LEU:C	1:C:125:LEU:HD23	2.42	0.45
1:C:136:ILE:HD12	1:C:136:ILE:N	2.33	0.44
1:A:63:THR:CG2	1:A:123:ILE:HG21	2.47	0.44
1:A:53:LEU:HD21	1:A:60:LEU:HD12	1.99	0.44
1:B:93:VAL:O	1:B:185:ASN:HA	2.17	0.44
1:D:53:LEU:HD23	1:D:55:VAL:O	2.17	0.44
1:A:56:PRO:HD3	1:A:136:ILE:O	2.18	0.43
1:D:161:ILE:HG13	1:D:185:ASN:HB2	1.99	0.43
1:C:136:ILE:N	1:C:136:ILE:CD1	2.82	0.43
1:D:86:SER:OG	1:D:193:VAL:HB	2.19	0.42
1:D:46:PHE:CZ	1:D:64:PHE:HB3	2.55	0.42
1:B:63:THR:CG2	1:B:123:ILE:HG21	2.50	0.41
1:C:39:TYR:CE1	1:D:211:PRO:CD	3.04	0.41
1:C:53:LEU:HD21	1:C:60:LEU:HD12	2.03	0.41
1:C:14:ILE:HG22	1:C:15:LEU:N	2.35	0.41
1:D:21:ASP:OD1	1:D:21:ASP:C	2.63	0.41
1:C:133:ASP:O	1:C:140:LYS:NZ	2.54	0.40
1:C:31:GLY:HA2	1:C:45:LYS:O	2.21	0.40
1:C:39:TYR:CE1	1:D:211:PRO:HD2	2.56	0.40
1:C:88:MET:HB3	1:C:89:PRO:HA	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	226/229 (99%)	223 (99%)	3 (1%)	0	100	100
1	B	231/229 (101%)	228 (99%)	3 (1%)	0	100	100
1	C	222/229 (97%)	216 (97%)	6 (3%)	0	100	100
1	D	222/229 (97%)	217 (98%)	5 (2%)	0	100	100
All	All	901/916 (98%)	884 (98%)	17 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	190/199 (96%)	188 (99%)	2 (1%)	65	67
1	B	198/199 (100%)	196 (99%)	2 (1%)	68	70
1	C	175/199 (88%)	172 (98%)	3 (2%)	53	52
1	D	176/199 (88%)	173 (98%)	3 (2%)	53	52
All	All	739/796 (93%)	729 (99%)	10 (1%)	59	59

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	211	PRO
1	A	230	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	124	GLU
1	B	230	THR
1	C	123	ILE
1	C	136	ILE
1	C	186	THR
1	D	5	GLU
1	D	15	LEU
1	D	164	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	149	ASN
1	A	164	ASN
1	A	212	ASN
1	B	105	ASN
1	B	149	ASN
1	B	157	GLN
1	C	77	HIS
1	C	81	HIS
1	D	149	ASN
1	D	164	ASN
1	D	198	ASN
1	D	212	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CRO	A	65	1	20,22,24	3.19	4 (20%)	25,30,34	3.21	12 (48%)
1	CRO	B	65	1	20,22,24	2.82	6 (30%)	25,30,34	3.06	10 (40%)
1	CRO	D	65	1	20,22,24	3.12	5 (25%)	25,30,34	2.14	9 (36%)
1	CRO	C	65	1	20,22,24	3.23	5 (25%)	25,30,34	2.46	8 (32%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CRO	A	65	1	-	3/10/29/32	0/2/2/2
1	CRO	B	65	1	-	2/10/29/32	0/2/2/2
1	CRO	D	65	1	-	2/10/29/32	0/2/2/2
1	CRO	C	65	1	-	3/10/29/32	0/2/2/2

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	65	CRO	CB2-CA2	12.28	1.47	1.35
1	C	65	CRO	CB2-CA2	11.26	1.46	1.35
1	D	65	CRO	CB2-CA2	11.05	1.45	1.35
1	B	65	CRO	CB2-CA2	10.02	1.44	1.35
1	D	65	CRO	O2-C2	4.93	1.33	1.23
1	C	65	CRO	CA2-C2	-4.91	1.43	1.48
1	C	65	CRO	C2-N3	-4.51	1.29	1.40
1	A	65	CRO	C2-N3	-4.36	1.30	1.40
1	C	65	CRO	O2-C2	4.19	1.31	1.23
1	B	65	CRO	C2-N3	-3.86	1.31	1.40
1	A	65	CRO	O2-C2	3.81	1.30	1.23
1	D	65	CRO	C2-N3	-3.46	1.32	1.40
1	B	65	CRO	O2-C2	3.37	1.29	1.23
1	D	65	CRO	C1-N2	3.21	1.36	1.32
1	D	65	CRO	CA2-C2	-3.15	1.45	1.48
1	B	65	CRO	C1-N2	3.08	1.36	1.32
1	B	65	CRO	CA2-C2	-2.46	1.45	1.48
1	C	65	CRO	C1-N2	2.34	1.35	1.32
1	B	65	CRO	CE1-CZ	2.27	1.43	1.39
1	A	65	CRO	CE1-CD1	2.03	1.42	1.38

All (39) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	65	CRO	O2-C2-CA2	-12.18	123.25	131.02
1	B	65	CRO	O2-C2-CA2	-11.00	124.00	131.02
1	C	65	CRO	O2-C2-CA2	-5.86	127.28	131.02
1	C	65	CRO	CA2-C2-N3	5.04	107.74	103.50
1	D	65	CRO	C3-CA3-N3	4.91	123.59	112.43
1	D	65	CRO	CB2-CA2-N2	-4.20	123.06	128.76
1	C	65	CRO	CA1-C1-N3	-4.18	119.40	124.84
1	B	65	CRO	C3-CA3-N3	4.14	121.84	112.43
1	B	65	CRO	CD2-CG2-CD1	4.01	123.60	117.65
1	C	65	CRO	C3-CA3-N3	3.97	121.45	112.43
1	B	65	CRO	CE1-CD1-CG2	-3.95	116.12	121.22
1	B	65	CRO	CA2-N2-C1	-3.77	102.86	105.80
1	A	65	CRO	C3-CA3-N3	3.60	120.63	112.43
1	D	65	CRO	CB2-CA2-C2	3.60	126.72	122.36
1	A	65	CRO	C2-N3-C1	-3.59	106.41	108.07
1	A	65	CRO	CA2-C2-N3	3.43	106.38	103.50
1	B	65	CRO	CA2-C2-N3	3.40	106.36	103.50
1	A	65	CRO	CE1-CD1-CG2	-3.40	116.83	121.22
1	C	65	CRO	CA2-N2-C1	-3.39	103.15	105.80
1	C	65	CRO	C2-N3-C1	-3.28	106.55	108.07
1	A	65	CRO	CD2-CG2-CD1	3.18	122.36	117.65
1	D	65	CRO	CA1-C1-N3	-3.08	120.83	124.84
1	D	65	CRO	C2-N3-C1	3.06	109.48	108.07
1	A	65	CRO	N3-C1-N2	2.92	113.78	111.48
1	A	65	CRO	CA1-C1-N3	-2.72	121.30	124.84
1	B	65	CRO	N3-C1-N2	2.57	113.51	111.48
1	B	65	CRO	O2-C2-N3	2.53	129.59	124.31
1	D	65	CRO	O2-C2-CA2	2.53	132.64	131.02
1	C	65	CRO	CG2-CB2-CA2	-2.53	126.86	129.87
1	D	65	CRO	CA1-C1-N2	2.50	128.45	123.57
1	C	65	CRO	CA1-C1-N2	2.31	128.07	123.57
1	A	65	CRO	CA3-N3-C2	2.31	128.74	123.67
1	A	65	CRO	CD2-CG2-CB2	-2.26	113.47	121.22
1	B	65	CRO	CD2-CE2-CZ	-2.17	117.59	119.88
1	A	65	CRO	CD2-CE2-CZ	-2.10	117.65	119.88
1	D	65	CRO	CA2-C2-N3	2.08	105.25	103.50
1	A	65	CRO	O2-C2-N3	2.06	128.61	124.31
1	D	65	CRO	CD2-CG2-CB2	-2.06	114.15	121.22
1	B	65	CRO	CE2-CZ-CE1	2.02	122.98	119.77

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	C	65	CRO	N1-CA1-CB1-OG1
1	A	65	CRO	C3-CA3-N3-C2
1	C	65	CRO	C1-CA1-CB1-OG1
1	B	65	CRO	C1-CA1-CB1-OG1
1	D	65	CRO	C3-CA3-N3-C2
1	A	65	CRO	C1-CA1-CB1-OG1
1	C	65	CRO	C3-CA3-N3-C2
1	A	65	CRO	N1-CA1-CB1-OG1
1	B	65	CRO	N1-CA1-CB1-OG1
1	D	65	CRO	N1-CA1-CB1-OG1

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	C	65	CRO	4	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	C	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	C	65:CRO	C3	68:VAL	N	1.72

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	226/229 (98%)	-0.19	5 (2%) 62 66	9, 21, 36, 46	4 (1%)
1	B	228/229 (99%)	-0.38	2 (0%) 81 83	9, 19, 32, 43	7 (3%)
1	C	224/229 (97%)	0.64	12 (5%) 31 33	13, 32, 52, 65	2 (0%)
1	D	225/229 (98%)	0.47	13 (5%) 29 30	14, 30, 45, 56	1 (0%)
All	All	903/916 (98%)	0.13	32 (3%) 47 50	9, 25, 44, 65	14 (1%)

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2	SER	3.3
1	C	182	TYR	3.2
1	C	4	GLY	3.2
1	D	171	ILE	3.0
1	C	192	PRO	3.0
1	C	190	ASP	2.9
1	C	191	GLY	2.9
1	A	211	PRO	2.8
1	C	193	VAL	2.8
1	D	97	THR	2.7
1	C	171	ILE	2.7
1	D	22	VAL	2.7
1	D	192	PRO	2.6
1	D	174	GLY	2.6
1	D	182	TYR	2.5
1	A	172	GLU	2.5
1	D	132	GLU	2.5
1	D	173	ASP	2.4
1	A	230	THR	2.4
1	B	230	THR	2.3
1	D	176	VAL	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	151	TYR	2.3
1	C	229	ILE	2.3
1	C	3	LYS	2.3
1	B	133	ASP	2.2
1	C	132	GLU	2.2
1	C	195	LEU	2.2
1	D	229	ILE	2.1
1	C	172	GLU	2.1
1	D	128	ILE	2.1
1	D	157	GLN	2.1
1	A	212	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CRO	C	65	21/23	0.95	0.07	14,20,24,26	0
1	CRO	A	65	21/23	0.97	0.04	8,11,16,17	0
1	CRO	D	65	21/23	0.97	0.05	17,20,23,24	0
1	CRO	B	65	21/23	0.98	0.04	9,13,15,18	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CL	C	240	1/1	0.98	0.06	36,36,36,36	0
2	CL	B	231	1/1	0.99	0.09	30,30,30,30	0
2	CL	A	240	1/1	0.99	0.03	24,24,24,24	0

6.5 Other polymers ⓘ

There are no such residues in this entry.