



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 2, 2026 – 04:11 PM UTC

PDB ID : 3GJ5 / pdb_00003gj5
Title : Crystal structure of human RanGDP-Nup153ZnF4 complex
Authors : Partridge, J.R.; Schwartz, T.U.
Deposited on : 2009-03-07
Resolution : 1.79 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.79 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	221	18% 79% 12% 9%
1	C	221	15% 79% 12% 9%
2	B	34	62% 68% 9% 24%
2	D	34	44% 62% 15% 24%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 4112 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GTP-binding nuclear protein Ran.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	201	Total	C	N	O	S	0	0	0
			1607	1037	279	285	6			
1	C	201	Total	C	N	O	S	0	0	0
			1607	1037	279	285	6			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	GLY	-	expression tag	UNP P62826
A	-3	PRO	-	expression tag	UNP P62826
A	-2	HIS	-	expression tag	UNP P62826
A	-1	MET	-	expression tag	UNP P62826
A	0	ALA	-	expression tag	UNP P62826
A	1	SER	-	expression tag	UNP P62826
A	35	SER	PHE	engineered mutation	UNP P62826
C	-4	GLY	-	expression tag	UNP P62826
C	-3	PRO	-	expression tag	UNP P62826
C	-2	HIS	-	expression tag	UNP P62826
C	-1	MET	-	expression tag	UNP P62826
C	0	ALA	-	expression tag	UNP P62826
C	1	SER	-	expression tag	UNP P62826
C	35	SER	PHE	engineered mutation	UNP P62826

- Molecule 2 is a protein called Nuclear pore complex protein Nup153.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	26	Total	C	N	O	S	0	0	0
			190	114	32	40	4			
2	D	26	Total	C	N	O	S	0	0	0
			190	114	32	40	4			


There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	843	GLY	-	expression tag	UNP P49791
B	844	PRO	-	expression tag	UNP P49791
B	845	LEU	-	expression tag	UNP P49791
B	846	GLY	-	expression tag	UNP P49791
B	847	SER	-	expression tag	UNP P49791
D	843	GLY	-	expression tag	UNP P49791
D	844	PRO	-	expression tag	UNP P49791
D	845	LEU	-	expression tag	UNP P49791
D	846	GLY	-	expression tag	UNP P49791
D	847	SER	-	expression tag	UNP P49791

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Mg 1 1	0	0
3	C	1	Total Mg 1 1	0	0

- Molecule 4 is GUANOSINE-5'-DIPHOSPHATE (CCD ID: GDP) (formula: C₁₀H₁₅N₅O₁₁P₂).



- Molecule 5 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total 1	Zn 1	0	0
5	D	1	Total 1	Zn 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	230	Total 230	O 230	0	0
6	B	7	Total 7	O 7	0	0
6	C	211	Total 211	O 211	0	0
6	D	10	Total 10	O 10	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: GTP-binding nuclear protein Ran




• Molecule 1: GTP-binding nuclear protein Ran



• Molecule 2: Nuclear pore complex protein Nup153



• Molecule 2: Nuclear pore complex protein Nup153



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	70.44Å 61.34Å 74.03Å 90.00° 112.55° 90.00°	Depositor
Resolution (Å)	45.66 – 1.79 45.66 – 1.79	Depositor EDS
% Data completeness (in resolution range)	97.7 (45.66-1.79) 97.8 (45.66-1.79)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.91 (at 1.79Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.199 , 0.233 0.196 , 0.230	Depositor DCC
R_{free} test set	2730 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	23.2	Xtriage
Anisotropy	0.468	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 54.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4112	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.70% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GDP, MG, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.46	0/1646	0.79	1/2231 (0.0%)
1	C	0.46	0/1646	0.74	0/2231
2	B	0.35	0/191	0.64	0/256
2	D	0.32	0/191	0.63	0/256
All	All	0.45	0/3674	0.75	1/4974 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	113	GLU	CB-CA-C	-7.82	106.60	117.23

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1607	0	1623	17	0
1	C	1607	0	1623	21	0
2	B	190	0	178	3	0
2	D	190	0	178	3	0
3	A	1	0	0	0	0
3	C	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	28	0	12	0	0
4	C	28	0	12	0	0
5	B	1	0	0	0	0
5	D	1	0	0	0	0
6	A	230	0	0	2	0
6	B	7	0	0	0	0
6	C	211	0	0	3	0
6	D	10	0	0	1	0
All	All	4112	0	3626	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:850:TRP:CD1	2:B:873:LYS:HZ3	1.94	0.86
1:C:159:LYS:HB2	1:C:160:PRO:HD3	1.77	0.65
1:A:159:LYS:HB2	1:A:160:PRO:HD3	1.77	0.65
1:A:55:ASN:OD1	1:A:174:LEU:HD12	1.98	0.64
1:A:81:ILE:HG21	2:B:868:ALA:HA	1.79	0.63
1:C:53:HIS:HD2	6:C:261:HOH:O	1.81	0.62
1:A:53:HIS:HD2	6:A:377:HOH:O	1.83	0.61
1:C:142:LYS:O	1:C:143:ASN:HB3	2.03	0.59
1:A:32:THR:OG1	1:C:53:HIS:HE1	1.85	0.58
1:A:24:THR:CG2	1:A:28:LYS:HE2	2.33	0.57
1:A:53:HIS:HE1	1:C:32:THR:OG1	1.89	0.55
1:A:53:HIS:O	1:A:177:VAL:HG22	2.11	0.50
1:C:53:HIS:CD2	6:C:261:HOH:O	2.62	0.49
1:C:55:ASN:OD1	1:C:174:LEU:HD12	2.12	0.49
1:C:153:SER:O	1:C:154:ASN:HB2	2.12	0.49
1:C:137:VAL:HG12	1:C:141:LYS:HE3	1.94	0.49
1:C:81:ILE:HG21	2:D:868:ALA:HA	1.95	0.49
1:A:105:HIS:CE1	1:A:142:LYS:HE2	2.47	0.48
1:A:153:SER:O	1:A:154:ASN:HB2	2.13	0.47
1:C:114:ASN:O	1:C:114:ASN:CG	2.58	0.47
2:D:850:TRP:CG	2:D:873:LYS:HZ2	2.33	0.46
1:C:30:HIS:CG	1:C:50:LEU:HG	2.51	0.45
1:A:81:ILE:CG2	2:B:868:ALA:HA	2.44	0.45
1:A:81:ILE:HG13	1:A:82:GLN:HG2	1.98	0.45
1:A:30:HIS:CG	1:A:50:LEU:HG	2.52	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:77:ASP:HB2	6:C:311:HOH:O	2.15	0.45
1:C:103:ASN:ND2	1:C:106:ARG:HH21	2.15	0.45
2:D:860:LYS:HE3	6:D:404:HOH:O	2.17	0.44
1:C:142:LYS:O	1:C:143:ASN:CB	2.65	0.44
1:A:124:VAL:HG22	1:A:149:ILE:O	2.18	0.43
1:C:112:CYS:O	1:C:113:GLU:HB2	2.19	0.43
1:C:39:TYR:CE2	1:C:41:ALA:HA	2.54	0.43
1:C:137:VAL:CG1	1:C:141:LYS:HE3	2.48	0.42
1:C:13:LEU:C	1:C:13:LEU:HD23	2.44	0.42
1:C:28:LYS:HG2	1:C:34:GLU:OE1	2.20	0.42
1:A:152:LYS:HB3	1:A:187:VAL:HG22	2.02	0.41
1:C:81:ILE:HD12	1:C:81:ILE:HA	1.96	0.41
1:A:48:HIS:HD2	6:A:236:HOH:O	2.03	0.41
1:A:202:GLU:O	1:A:206:THR:HG23	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	199/221 (90%)	193 (97%)	6 (3%)	0	100	100
1	C	199/221 (90%)	194 (98%)	5 (2%)	0	100	100
2	B	24/34 (71%)	22 (92%)	2 (8%)	0	100	100
2	D	24/34 (71%)	22 (92%)	2 (8%)	0	100	100
All	All	446/510 (88%)	431 (97%)	15 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	173/188 (92%)	172 (99%)	1 (1%)	78	77
1	C	173/188 (92%)	173 (100%)	0	100	100
2	B	22/27 (82%)	22 (100%)	0	100	100
2	D	22/27 (82%)	21 (96%)	1 (4%)	24	12
All	All	390/430 (91%)	388 (100%)	2 (0%)	81	80

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	144	LEU
2	D	864	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	48	HIS
1	A	53	HIS
1	A	100	ASN
1	A	103	ASN
1	A	105	HIS
1	A	139	HIS
1	A	199	HIS
1	C	10	GLN
1	C	48	HIS
1	C	53	HIS
1	C	62	ASN
1	C	100	ASN
1	C	103	ASN
1	C	139	HIS
1	C	199	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	GDP	A	302	3	29,30,30	1.17	2 (6%)	45,47,47	1.70	7 (15%)
4	GDP	C	302	3	29,30,30	1.16	2 (6%)	45,47,47	1.66	6 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GDP	A	302	3	-	0/16/32/32	0/3/3/3
4	GDP	C	302	3	-	0/16/32/32	0/3/3/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	302	GDP	C5-C4	3.29	1.47	1.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	302	GDP	C5-C4	3.17	1.47	1.38
4	A	302	GDP	C6-N1	-2.51	1.34	1.38
4	C	302	GDP	C6-N1	-2.28	1.34	1.38

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	302	GDP	C5-C4-N3	-5.62	119.45	128.39
4	C	302	GDP	C5-C4-N3	-5.48	119.67	128.39
4	A	302	GDP	C2-N3-C4	4.69	120.38	112.30
4	C	302	GDP	C2-N3-C4	4.56	120.16	112.30
4	A	302	GDP	N9-C4-N3	4.35	134.65	125.95
4	C	302	GDP	N9-C4-N3	4.26	134.48	125.95
4	A	302	GDP	C6-C5-N7	3.39	136.46	130.29
4	C	302	GDP	C6-C5-N7	3.36	136.40	130.29
4	C	302	GDP	C4-C5-N7	-2.40	106.86	110.67
4	A	302	GDP	C4-C5-N7	-2.39	106.88	110.67
4	A	302	GDP	O2A-PA-O3A	2.20	113.22	107.27
4	A	302	GDP	O6-C6-C5	-2.19	120.75	126.53
4	C	302	GDP	O6-C6-C5	-2.18	120.79	126.53


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	201/221 (90%)	1.41	39 (19%) 3 2	27, 37, 62, 83	0
1	C	201/221 (90%)	1.30	34 (16%) 4 3	27, 37, 63, 81	0
2	B	26/34 (76%)	2.57	21 (80%) 0 0	37, 67, 98, 132	0
2	D	26/34 (76%)	2.27	15 (57%) 0 0	38, 63, 97, 112	0
All	All	454/510 (89%)	1.48	109 (24%) 2 1	27, 38, 77, 132	0

All (109) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	144	LEU	5.5
1	C	143	ASN	4.9
1	C	114	ASN	4.9
1	A	137	VAL	4.7
1	A	208	ALA	4.5
1	A	114	ASN	4.5
2	B	861	ALA	4.4
1	A	193	LEU	4.0
1	C	208	ALA	4.0
1	A	143	ASN	3.9
2	D	848	GLY	3.9
1	C	188	VAL	3.9
1	C	144	LEU	3.8
2	B	848	GLY	3.8
2	D	872	ALA	3.8
2	D	853	GLU	3.7
2	B	871	SER	3.6
1	A	8	GLN	3.6
1	A	188	VAL	3.5
2	B	872	ALA	3.5
1	A	109	VAL	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	137	VAL	3.4
1	A	126	ILE	3.3
2	B	863	SER	3.3
1	C	190	ASP	3.2
1	A	81	ILE	3.2
2	D	861	ALA	3.2
1	A	113	GLU	3.1
2	B	853	GLU	3.1
2	B	867	ILE	3.1
1	A	112	CYS	3.1
2	B	862	ASP	3.1
1	A	99	LYS	3.0
1	C	127	LYS	3.0
1	C	192	ALA	3.0
2	D	864	THR	3.0
2	B	873	LYS	3.0
1	A	14	VAL	3.0
1	C	8	GLN	2.9
1	C	81	ILE	2.9
1	C	59	ILE	2.9
1	C	108	LEU	2.8
1	A	136	ILE	2.8
1	C	14	VAL	2.8
2	D	862	ASP	2.8
2	B	858	GLN	2.8
2	D	867	ILE	2.8
1	A	196	GLN	2.7
1	C	193	LEU	2.7
2	D	850	TRP	2.7
1	A	111	VAL	2.7
1	C	118	VAL	2.7
2	B	854	VAL	2.7
2	B	864	THR	2.7
2	D	863	SER	2.7
2	D	868	ALA	2.7
2	B	850	TRP	2.7
1	A	192	ALA	2.6
1	A	108	LEU	2.6
2	D	856	LEU	2.6
1	C	111	VAL	2.6
1	C	136	ILE	2.6
1	C	206	THR	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	849	SER	2.6
2	D	858	GLN	2.5
1	A	127	LYS	2.5
1	A	141	LYS	2.5
1	A	190	ASP	2.5
1	A	142	LYS	2.5
1	A	177	VAL	2.5
1	C	126	ILE	2.5
1	C	195	ALA	2.5
1	C	141	LYS	2.5
2	B	865	LYS	2.5
2	B	860	LYS	2.4
1	C	204	ALA	2.4
1	C	69	GLN	2.4
1	C	31	LEU	2.4
1	A	97	THR	2.4
2	B	851	ASP	2.4
1	A	96	VAL	2.4
1	C	177	VAL	2.3
2	D	873	LYS	2.3
1	A	31	LEU	2.3
1	A	149	ILE	2.3
1	C	113	GLU	2.3
2	D	860	LYS	2.3
1	A	195	ALA	2.3
1	A	203	VAL	2.2
2	B	870	GLU	2.2
1	C	110	ARG	2.2
2	B	856	LEU	2.2
1	A	39	TYR	2.2
1	C	39	TYR	2.2
1	C	79	TYR	2.2
1	C	179	MET	2.1
1	A	9	VAL	2.1
2	B	857	VAL	2.1
1	A	207	THR	2.1
1	A	169	ILE	2.1
1	C	205	GLN	2.1
1	A	42	THR	2.1
1	A	44	GLY	2.0
1	C	44	GLY	2.0
2	B	868	ALA	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	82	GLN	2.0
2	D	865	LYS	2.0
1	A	79	TYR	2.0
1	C	109	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	GDP	A	302	28/28	0.97	0.08	28,33,38,43	0
4	GDP	C	302	28/28	0.98	0.07	28,33,40,42	0
3	MG	A	301	1/1	0.99	0.05	31,31,31,31	0
3	MG	C	301	1/1	0.99	0.05	30,30,30,30	0
5	ZN	B	300	1/1	1.00	0.03	41,41,41,41	0
5	ZN	D	300	1/1	1.00	0.04	41,41,41,41	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around GDP A 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around GDP C 302:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.