



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 12:33 PM UTC

PDB ID : 5GJ7 / pdb_00005gj7
Title : putative Acyl-CoA dehydrogenase
Authors : Moon, H.; Choe, J.
Deposited on : 2016-06-28
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3494 (1.96-1.96)
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	398	6% 90% 7% .
1	B	398	10% 89% 8% ..

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 6313 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Acyl-CoA dehydrogenase type 2 domain protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	388	Total	C	N	O	S	0	0	0
			3021	1912	539	558	12			
1	B	388	Total	C	N	O	S	0	0	0
			3021	1912	539	558	12			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	expression tag	UNP C8WV06
A	-1	ASN	-	expression tag	UNP C8WV06
A	0	ALA	-	expression tag	UNP C8WV06
B	-2	SER	-	expression tag	UNP C8WV06
B	-1	ASN	-	expression tag	UNP C8WV06
B	0	ALA	-	expression tag	UNP C8WV06


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	161	Total	O	0	0
			161	161		
2	B	110	Total	O	0	0
			110	110		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Acyl-CoA dehydrogenase type 2 domain protein



- Molecule 1: Acyl-CoA dehydrogenase type 2 domain protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	155.69Å 69.38Å 83.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.95 20.00 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.0 (20.00-1.95) 99.0 (20.00-1.95)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.22 (at 1.96Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.196 , 0.224 0.210 , 0.236	Depositor DCC
R_{free} test set	3318 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	25.4	Xtriage
Anisotropy	0.559	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 49.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.54$, $\langle L^2 \rangle = 0.38$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6313	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 38.82 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.4818e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.10	2/3084 (0.1%)	0.99	2/4190 (0.0%)
1	B	1.01	2/3084 (0.1%)	0.98	1/4190 (0.0%)
All	All	1.06	4/6168 (0.1%)	0.98	3/8380 (0.0%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	373	PRO	CA-C	7.06	1.55	1.51
1	B	373	PRO	CA-C	6.02	1.55	1.51
1	B	332	LEU	N-CA	5.20	1.52	1.46
1	A	296	LYS	N-CA	5.14	1.52	1.46

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	293	MET	CG-SD-CE	-5.84	88.04	100.90
1	B	293	MET	CG-SD-CE	-5.58	88.63	100.90
1	A	234	VAL	N-CA-C	5.22	120.20	109.34

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3021	0	3026	13	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	3021	0	3026	16	0
2	A	161	0	0	0	0
2	B	110	0	0	0	0
All	All	6313	0	6052	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:165:THR:O	1:B:209:THR:OG1	2.00	0.79
1:B:211:SER:O	1:B:212:HIS:HD2	1.86	0.59
1:B:234:VAL:HG23	1:B:235:GLN:HA	1.85	0.58
1:A:234:VAL:HG23	1:A:235:GLN:HA	1.89	0.55
1:A:307:LEU:HD12	1:A:329:VAL:CG2	2.38	0.53
1:B:307:LEU:HD12	1:B:329:VAL:CG2	2.41	0.50
1:B:34:GLU:HG3	1:B:37:ARG:HH22	1.77	0.49
1:A:156:ARG:HD2	1:A:218:ASP:OD1	2.13	0.49
1:A:214:ILE:HD11	1:A:216:LEU:HD11	1.95	0.48
1:B:211:SER:O	1:B:212:HIS:CD2	2.66	0.48
1:B:214:ILE:HD11	1:B:216:LEU:HD11	1.96	0.48
1:A:307:LEU:CD1	1:A:329:VAL:CG2	2.93	0.47
1:A:355:ARG:HD3	1:B:202:ASP:OD2	2.17	0.45
1:B:307:LEU:CD1	1:B:329:VAL:CG2	2.95	0.44
1:A:201:TRP:CD2	1:B:356:ALA:HB2	2.53	0.43
1:A:32:ALA:N	1:A:33:PRO:CD	2.82	0.42
1:A:363:TYR:CZ	1:B:367:ARG:HD3	2.55	0.42
1:B:129:CYS:SG	1:B:164:SER:HB3	2.59	0.42
1:B:164:SER:OG	1:B:214:ILE:HG23	2.20	0.42
1:A:198:VAL:CG2	1:A:213:ASP:HB2	2.49	0.42
1:A:367:ARG:HD3	1:B:363:TYR:CZ	2.56	0.41
1:B:172:TRP:HB3	1:B:189:LEU:HD11	2.02	0.41
1:B:242:GLY:O	1:B:323:GLN:HG2	2.20	0.41
1:A:307:LEU:CD1	1:A:329:VAL:HG23	2.50	0.40
1:A:172:TRP:HB3	1:A:189:LEU:HD11	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	384/398 (96%)	381 (99%)	2 (0%)	1 (0%)	36	28
1	B	384/398 (96%)	381 (99%)	2 (0%)	1 (0%)	36	28
All	All	768/796 (96%)	762 (99%)	4 (0%)	2 (0%)	36	28

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	236	ALA
1	A	236	ALA

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	313/319 (98%)	304 (97%)	9 (3%)	37	29
1	B	313/319 (98%)	305 (97%)	8 (3%)	40	33
All	All	626/638 (98%)	609 (97%)	17 (3%)	39	32

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	68	LEU
1	A	145	GLU
1	A	164	SER
1	A	175	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	211	SER
1	A	214	ILE
1	A	234	VAL
1	A	375	MET
1	A	395	GLN
1	B	68	LEU
1	B	145	GLU
1	B	175	VAL
1	B	214	ILE
1	B	234	VAL
1	B	295	LEU
1	B	375	MET
1	B	395	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	77	GLN
1	A	371	HIS
1	B	77	GLN
1	B	212	HIS
1	B	371	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	388/398 (97%)	0.23	22 (5%) 29 34	17, 29, 64, 119	0
1	B	388/398 (97%)	0.55	39 (10%) 12 14	18, 34, 75, 145	0
All	All	776/796 (97%)	0.39	61 (7%) 18 21	17, 32, 72, 145	0

All (61) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	210	GLY	11.0
1	B	211	SER	8.3
1	B	234	VAL	6.8
1	B	236	ALA	6.1
1	B	-2	SER	5.9
1	B	209	THR	5.9
1	B	232	PRO	5.4
1	A	-2	SER	4.8
1	B	208	ALA	4.6
1	B	152	PRO	4.5
1	A	235	GLN	4.3
1	A	164	SER	4.3
1	A	233	GLY	4.3
1	A	232	PRO	4.3
1	A	201	TRP	4.2
1	A	236	ALA	4.0
1	B	165	THR	3.9
1	B	235	GLN	3.8
1	A	234	VAL	3.7
1	B	394	GLY	3.7
1	B	233	GLY	3.6
1	B	180	ALA	3.6
1	B	129	CYS	3.6
1	B	144	PRO	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	-1	ASN	3.4
1	B	181	ASP	3.2
1	B	-1	ASN	3.2
1	A	9	ALA	3.1
1	B	40	ASP	3.0
1	B	38	ALA	2.9
1	B	215	VAL	2.9
1	A	274	ASN	2.8
1	B	200	THR	2.8
1	B	9	ALA	2.8
1	B	133	PRO	2.8
1	B	184	VAL	2.6
1	A	273	PRO	2.6
1	B	237	GLU	2.5
1	A	181	ASP	2.5
1	B	179	VAL	2.5
1	A	372	ASN	2.4
1	B	201	TRP	2.4
1	B	273	PRO	2.4
1	A	200	THR	2.3
1	B	153	GLY	2.3
1	A	144	PRO	2.3
1	B	34	GLU	2.3
1	B	10	LEU	2.3
1	B	151	VAL	2.3
1	A	133	PRO	2.2
1	B	218	ASP	2.2
1	A	8	ALA	2.2
1	B	37	ARG	2.2
1	B	355	ARG	2.2
1	A	394	GLY	2.1
1	B	217	LYS	2.1
1	A	211	SER	2.1
1	B	207	ARG	2.1
1	A	276	LEU	2.1
1	B	164	SER	2.0
1	A	275	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.