



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 02:57 AM UTC

PDB ID : 5GK4 / pdb_00005gk4
Title : Native structure of fructose 1,6-bisphosphate aldolase from Escherichia coli at 2.0 Angstrom resolution
Authors : Tran, T.H.; Huynh, K.H.; Ho, T.H.; Kang, L.W.
Deposited on : 2016-07-03
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	359	
1	B	359	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	PEG	A	405	-	-	X	-

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 5537 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fructose-bisphosphate aldolase class 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	329	Total	C	N	O	S	0	2	0
			2567	1634	434	490	9			
1	B	330	Total	C	N	O	S	0	2	0
			2571	1636	435	491	9			

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Zn	0	0
			2	2		
3	B	2	Total	Zn	0	0
			2	2		

- Molecule 4 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			7	4	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	175	Total	O	0	0
			175	175		
5	B	201	Total	O	0	0
			201	201		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Fructose-bisphosphate aldolase class 2



- Molecule 1: Fructose-bisphosphate aldolase class 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	73.13Å 72.89Å 72.45Å 90.00° 103.20° 90.00°	Depositor
Resolution (Å)	50.00 – 2.00 50.00 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.6 (50.00-2.00) 99.8 (50.00-2.00)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	10.62 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.153 , 0.200 0.165 , 0.206	Depositor DCC
R_{free} test set	2561 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	21.2	Xtriage
Anisotropy	0.058	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 39.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.015 for l,-k,h	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5537	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.79% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, ZN, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.53	28/2624 (1.1%)	1.19	14/3555 (0.4%)
1	B	1.56	26/2628 (1.0%)	1.16	10/3560 (0.3%)
All	All	1.54	54/5252 (1.0%)	1.18	24/7115 (0.3%)

All (54) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	339	THR	C-O	-7.18	1.15	1.24
1	B	282	VAL	C-O	-7.18	1.17	1.24
1	A	207	THR	C-O	-7.12	1.16	1.24
1	B	104	VAL	C-O	-6.92	1.16	1.24
1	A	335	ARG	C-O	-6.80	1.16	1.24
1	A	122	LEU	C-O	-6.67	1.16	1.24
1	A	104	VAL	C-O	-6.50	1.17	1.24
1	B	3	ILE	C-O	-6.45	1.16	1.24
1	B	293	TRP	C-O	-6.45	1.16	1.24
1	A	213	SER	C-O	-6.12	1.16	1.24
1	A	109	ASP	C-O	-6.12	1.16	1.23
1	A	120	ASP	C-O	-6.05	1.17	1.24
1	B	140	SER	C-O	-5.96	1.16	1.23
1	A	119	ILE	C-O	-5.87	1.17	1.24
1	B	212	ILE	C-O	-5.86	1.17	1.23
1	A	54	ALA	C-O	-5.74	1.16	1.24
1	B	90	ALA	C-O	-5.73	1.17	1.24
1	B	290	ASP	C-O	-5.68	1.17	1.24
1	A	68	ILE	C-O	-5.62	1.17	1.24
1	B	124	ASP	C-O	-5.62	1.17	1.24
1	A	137	LEU	C-O	-5.60	1.17	1.24
1	B	72	GLY	C-O	-5.60	1.17	1.24
1	A	62	ASN	C-O	-5.54	1.17	1.24
1	A	97	ALA	C-O	-5.53	1.17	1.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	282	VAL	C-O	-5.53	1.18	1.24
1	B	95	GLN	C-O	-5.52	1.17	1.24
1	A	266	GLY	C-O	-5.49	1.19	1.24
1	A	121	GLY	C-O	-5.48	1.17	1.23
1	A	336	ALA	C-O	-5.48	1.17	1.24
1	A	287	ILE	C-O	5.48	1.29	1.24
1	B	93	VAL	C-O	-5.47	1.18	1.24
1	B	131	ALA	N-CA	5.47	1.52	1.46
1	A	239	THR	C-O	-5.44	1.17	1.24
1	B	71	LYS	C-O	-5.40	1.17	1.24
1	B	98	GLU	C-O	-5.39	1.17	1.24
1	B	211	LYS	C-O	-5.37	1.17	1.24
1	B	239	THR	C-O	-5.37	1.17	1.24
1	A	294	ALA	C-O	-5.35	1.17	1.24
1	A	293	TRP	C-O	-5.35	1.18	1.24
1	B	171	LEU	C-O	-5.32	1.17	1.24
1	A	15	ASP	C-O	-5.30	1.17	1.24
1	B	20	VAL	C-O	-5.28	1.18	1.24
1	B	283	VAL	C-O	-5.28	1.16	1.24
1	A	355	ILE	CA-CB	-5.23	1.47	1.54
1	B	88	SER	C-O	-5.19	1.18	1.24
1	B	279	SER	C-O	-5.19	1.17	1.24
1	B	240	ILE	C-O	-5.19	1.18	1.24
1	A	100	TYR	C-O	-5.17	1.17	1.24
1	B	209	LEU	C-O	-5.16	1.18	1.24
1	A	105	ILE	C-O	-5.13	1.18	1.24
1	A	3	ILE	C-O	-5.08	1.18	1.24
1	A	131	ALA	C-O	-5.06	1.18	1.24
1	A	312	GLY	C-O	5.06	1.28	1.23
1	B	57	ILE	C-O	-5.04	1.18	1.24

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	96	MET	N-CA-C	7.47	122.69	112.90
1	A	96	MET	N-CA-C	7.46	122.67	112.90
1	B	34	VAL	N-CA-C	7.30	118.65	107.99
1	B	269	SER	N-CA-C	-7.29	100.71	110.55
1	B	54	ALA	N-CA-C	7.16	114.67	108.22
1	B	218	ILE	N-CA-C	6.86	117.92	107.77
1	B	107	HIS	N-CA-C	6.49	119.04	109.24
1	A	235	VAL	N-CA-C	-6.21	99.48	108.17

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	107	HIS	N-CA-CB	-6.17	101.62	111.62
1	A	132	ALA	N-CA-C	6.06	117.97	111.36
1	B	107	HIS	N-CA-CB	-5.96	101.32	111.69
1	A	54	ALA	N-CA-C	5.95	113.58	108.22
1	A	140	SER	N-CA-C	5.88	117.18	108.60
1	A	76	ASP	N-CA-C	5.72	119.03	112.57
1	A	107	HIS	N-CA-C	5.64	117.55	109.14
1	B	54	ALA	CA-C-N	-5.64	114.94	120.52
1	B	54	ALA	C-N-CA	-5.64	114.94	120.52
1	A	97	ALA	N-CA-C	5.28	117.45	111.11
1	A	351	GLU	N-CA-C	5.22	117.71	111.71
1	B	100	TYR	N-CA-C	-5.11	107.09	113.38
1	A	212	ILE	N-CA-C	5.07	116.17	111.45
1	A	54	ALA	CA-C-N	-5.06	115.35	120.31
1	A	54	ALA	C-N-CA	-5.06	115.35	120.31
1	A	6	PHE	N-CA-C	5.04	120.29	114.04

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2567	0	2526	23	0
1	B	2571	0	2529	16	0
2	A	12	0	16	0	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
4	A	7	0	10	10	0
5	A	175	0	0	3	0
5	B	201	0	0	1	0
All	All	5537	0	5081	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:73:VAL:C	1:B:74:LYS:HE2	1.75	1.10
1:B:73:VAL:O	1:B:74:LYS:HE2	1.64	0.97
1:A:91:HIS:HD1	4:A:405:PEG:HO4	0.98	0.89
1:B:73:VAL:O	1:B:74:LYS:CE	2.23	0.86
1:A:250:LYS:HE2	5:A:633:HOH:O	1.79	0.82
1:A:28:ASN:ND2	5:A:501:HOH:O	2.12	0.78
4:A:405:PEG:H41	1:B:91:HIS:HB3	1.69	0.73
4:A:405:PEG:H11	1:B:88:SER:HA	1.73	0.70
1:A:88:SER:OG	4:A:405:PEG:H31	1.92	0.69
1:B:73:VAL:O	1:B:74:LYS:NZ	2.30	0.64
1:A:293:TRP:CH2	1:B:312:GLY:HA2	2.44	0.52
1:A:141:HIS:CD2	1:A:169:MET:HE1	2.46	0.50
1:A:142:MET:HE1	1:A:264[B]:HIS:HE1	1.76	0.50
1:A:35:ASN:OD1	1:A:59:GLN:HG3	2.10	0.50
4:A:405:PEG:C1	1:B:88:SER:HA	2.40	0.49
4:A:405:PEG:H12	1:B:88:SER:OG	2.12	0.49
1:A:196:TYR:HB3	1:A:240:ILE:CD1	2.43	0.48
1:A:71:LYS:HE3	5:B:546:HOH:O	2.13	0.47
1:A:250:LYS:CE	5:A:633:HOH:O	2.51	0.46
1:A:39:THR:HG23	1:B:39:THR:HG23	1.98	0.45
1:B:316:ASN:HB2	1:B:317:PRO:HD2	1.98	0.45
1:A:55:PRO:HB3	1:A:103:PRO:HG2	1.98	0.45
1:A:312:GLY:HA2	1:B:293:TRP:CH2	2.51	0.45
1:A:33:ALA:HA	1:A:57:ILE:HB	1.99	0.43
1:A:250:LYS:HG3	1:A:251:LYS:N	2.34	0.43
1:B:65:ALA:O	1:B:85:GLY:HA3	2.19	0.43
1:B:316:ASN:HB2	1:B:317:PRO:CD	2.49	0.43
1:A:142:MET:CE	1:A:264[B]:HIS:HE1	2.31	0.42
1:A:91:HIS:CB	4:A:405:PEG:H22	2.50	0.42
1:A:142:MET:HE1	1:A:264[B]:HIS:CE1	2.54	0.42
4:A:405:PEG:H11	1:B:88:SER:CA	2.45	0.42
1:A:316:ASN:HB2	1:A:317:PRO:CD	2.51	0.41
1:A:3:ILE:HB	1:A:358:LEU:HA	2.02	0.41
1:A:91:HIS:HB2	4:A:405:PEG:H32	2.02	0.41
1:B:330:PRO:HA	1:B:333:TRP:CE2	2.57	0.40
1:A:91:HIS:HB2	4:A:405:PEG:H22	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	325/359 (90%)	317 (98%)	6 (2%)	2 (1%)	21	17
1	B	326/359 (91%)	318 (98%)	6 (2%)	2 (1%)	21	17
All	All	651/718 (91%)	635 (98%)	12 (2%)	4 (1%)	36	17

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	110[A]	HIS
1	A	110[B]	HIS
1	B	110[A]	HIS
1	B	110[B]	HIS

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	277/297 (93%)	267 (96%)	10 (4%)	31	31
1	B	277/297 (93%)	273 (99%)	4 (1%)	59	66
All	All	554/594 (93%)	540 (98%)	14 (2%)	42	45

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	25	LYS
1	A	28	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	31	LEU
1	A	171	LEU
1	A	175	LEU
1	A	234	VAL
1	A	235	VAL
1	A	236	LEU
1	A	240	ILE
1	A	259	LEU
1	B	25	LYS
1	B	240	ILE
1	B	241	LEU
1	B	326	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	99	HIS
1	A	256	HIS
1	B	313	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 4 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	PEG	A	405	-	6,6,6	0.35	0	5,5,5	1.36	0
2	GOL	A	402	-	5,5,5	0.58	0	5,5,5	0.49	0
2	GOL	A	401	-	5,5,5	0.53	0	5,5,5	1.26	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PEG	A	405	-	-	4/4/4/4	-
2	GOL	A	402	-	-	1/4/4/4	-
2	GOL	A	401	-	-	2/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	GOL	C3-C2-C1	-2.32	103.28	111.80

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	GOL	C1-C2-C3-O3
4	A	405	PEG	O2-C3-C4-O4
4	A	405	PEG	O1-C1-C2-O2
2	A	401	GOL	O2-C2-C3-O3
4	A	405	PEG	C4-C3-O2-C2
4	A	405	PEG	C1-C2-O2-C3
2	A	402	GOL	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	405	PEG	10	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	329/359 (91%)	-0.25	4 (1%) 76 76	11, 22, 41, 87	2 (0%)
1	B	330/359 (91%)	-0.30	8 (2%) 59 59	9, 20, 42, 65	2 (0%)
All	All	659/718 (91%)	-0.27	12 (1%) 67 67	9, 21, 42, 87	4 (0%)

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	235	VAL	3.9
1	A	234	VAL	3.7
1	B	225	VAL	3.6
1	A	196	TYR	3.3
1	B	196	TYR	3.0
1	B	74	LYS	2.9
1	B	223	GLY	2.4
1	A	268	GLY	2.3
1	B	320	GLU	2.2
1	B	176	GLY	2.1
1	B	271	ALA	2.1
1	B	319	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	GOL	A	402	6/6	0.88	0.11	41,42,43,45	0
4	PEG	A	405	7/7	0.88	0.21	23,27,33,39	0
2	GOL	A	401	6/6	0.92	0.11	31,34,38,40	0
3	ZN	A	404	1/1	0.99	0.03	21,21,21,21	1
3	ZN	B	401	1/1	0.99	0.02	24,24,24,24	1
3	ZN	A	403	1/1	0.99	0.04	28,28,28,28	1
3	ZN	B	402	1/1	1.00	0.04	19,19,19,19	1

6.5 Other polymers [i](#)

There are no such residues in this entry.