



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 03:38 PM UTC

PDB ID : 6GKL / pdb_00006gkl
Title : Translation initiation factor 4E in complex with beta-phosphorothioate trinucleotide mRNA 5' cap diastereomer 2 (m7GppSpApG D2)
Authors : Warminski, M.; Nowak, E.; Kowalska, J.; Jemielity, J.; Nowotny, M.
Deposited on : 2018-05-21
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	190	2% 89% 7% .
1	B	190	3% 87% 9% ..
1	C	190	25% 91% 5% .
1	D	190	19% 81% 14% 5%

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 5908 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Eukaryotic translation initiation factor 4E.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	183	Total	C	N	O	S	0	3	0
			1478	948	253	271	6			
1	B	184	Total	C	N	O	S	0	1	0
			1478	945	256	271	6			
1	C	180	Total	C	N	O	S	0	0	0
			1292	819	227	241	5			
1	D	180	Total	C	N	O	S	0	0	0
			1295	819	230	241	5			

- Molecule 2 is [((2 {R},3 {S},4 {R},5 {R})-5-(6-aminopurin-9-yl)-3-(((2 {R},3 {S},4 {R},5 {R})-5-(6-azany1-4,5-dihydropurin-9-yl)-3,4-bis(oxidanyl)oxolan-2-yl)methoxy-oxidanyl-phosphoryl]oxy-4-oxidanyl-oxolan-2-yl)methoxy-oxidanyl-phosphoryl]oxy-sulfanyl-phosphoryl] ((2 {R},3 {S},4 {R},5 {R})-5-(2-azany1-7-methyl-6-oxidanylidene-1 {H}-purin-9-yl)-3,4-bis(oxidanyl)oxolan-2-yl)methyl hydrogen phosphate (CCD ID: G0Z) (formula: C₃₁H₄₅N₁₅O₂₂P₄S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	S	0	0
			64	26	11	22	4	1		
2	B	1	Total	C	N	O	P	S	0	0
			33	11	5	13	3	1		
2	C	1	Total	C	N	O	P	S	0	0
			33	11	5	13	3	1		
2	D	1	Total	C	N	O	P	S	0	0
			33	11	5	13	3	1		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		



- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	76	Total	O	0	0
			76	76		
4	B	76	Total	O	0	0
			76	76		
4	C	16	Total	O	0	0
			16	16		
4	D	22	Total	O	0	0
			22	22		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Eukaryotic translation initiation factor 4E





- Molecule 1: Eukaryotic translation initiation factor 4E



- Molecule 1: Eukaryotic translation initiation factor 4E



- Molecule 1: Eukaryotic translation initiation factor 4E



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	38.17Å 38.18Å 147.02Å 88.80° 84.67° 76.13°	Depositor
Resolution (Å)	48.80 – 2.20 48.80 – 2.20	Depositor EDS
% Data completeness (in resolution range)	96.1 (48.80-2.20) 96.1 (48.80-2.20)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.83 (at 2.20Å)	Xtriage
Refinement program	PHENIX (1.10_2155: ???)	Depositor
R, R_{free}	0.218 , 0.257 0.222 , 0.258	Depositor DCC
R_{free} test set	1445 reflections (3.69%)	wwPDB-VP
Wilson B-factor (Å ²)	33.4	Xtriage
Anisotropy	0.242	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 62.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.036 for -h,-k,-h+l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	5908	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.79% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, G0Z

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.28	0/1525	0.52	0/2069
1	B	0.32	1/1519 (0.1%)	0.48	0/2065
1	C	0.22	0/1323	0.41	0/1812
1	D	0.22	0/1324	0.42	0/1812
All	All	0.27	1/5691 (0.0%)	0.46	0/7758

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	99	GLU	C-N	6.54	1.49	1.33

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1478	0	1400	7	0
1	B	1478	0	1392	9	0
1	C	1292	0	1077	7	0
1	D	1295	0	1082	17	0
2	A	64	0	0	0	0
2	B	33	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	33	0	0	3	0
2	D	33	0	0	1	0
3	A	6	0	8	0	0
3	B	6	0	8	2	0
4	A	76	0	0	0	0
4	B	76	0	0	0	0
4	C	16	0	0	0	0
4	D	22	0	0	0	0
All	All	5908	0	4967	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:122:ARG:NH2	1:B:157:ARG:O	2.16	0.78
1:D:42:ARG:HD3	1:D:67:ASP:HB3	1.70	0.73
1:D:157:ARG:HD3	1:D:158:ALA:H	1.59	0.66
1:A:80:GLN:NE2	1:A:84[B]:ASN:HD22	1.95	0.65
1:D:42:ARG:NH1	1:D:67:ASP:OD2	2.30	0.65
1:D:47:PHE:HB2	1:D:63:ILE:HD11	1.81	0.63
2:C:300:G0Z:S1	2:C:300:G0Z:O11	2.58	0.62
1:D:157:ARG:HG3	1:D:159:LYS:H	1.65	0.62
1:C:103:GLU:N	1:C:103:GLU:OE1	2.34	0.60
1:A:47:PHE:HB2	1:A:63:ILE:HD11	1.85	0.57
1:C:157:ARG:NH1	2:C:300:G0Z:O8	2.35	0.57
1:C:157:ARG:NH1	2:C:300:G0Z:S1	2.81	0.54
1:D:46:TRP:CG	1:D:100:PRO:HG3	2.43	0.54
1:B:195:ILE:HG22	3:B:302:GOL:H31	1.89	0.54
2:B:301:G0Z:S1	2:B:301:G0Z:O12	2.66	0.53
1:A:31:PRO:HA	1:A:34:TYR:CE1	2.44	0.52
1:D:42:ARG:HH11	1:D:67:ASP:CG	2.19	0.51
1:D:141:SER:HB2	1:D:186:ARG:NH1	2.25	0.51
1:B:48:PHE:HB3	1:B:90:ASP:HB2	1.92	0.50
1:D:141:SER:HB2	1:D:186:ARG:HH12	1.76	0.50
1:D:85:LEU:HB2	1:D:156:VAL:HG11	1.96	0.48
1:A:46:TRP:HB2	1:A:92:SER:HB2	1.96	0.47
1:D:150:CYS:SG	1:D:168:THR:HB	2.55	0.47
2:D:300:G0Z:S1	2:D:300:G0Z:O12	2.73	0.46
1:A:93:LEU:HD23	1:A:137:LEU:HD12	1.96	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:37:HIS:HD2	1:D:70:GLU:OE2	1.99	0.46
1:D:122:ARG:HG3	1:D:126:LEU:HD22	1.97	0.46
1:D:117:LEU:HD23	1:D:121:GLN:HB3	1.98	0.45
1:B:169:GLU:HB3	1:B:172:ASN:ND2	2.33	0.45
1:B:54:LYS:HE3	1:B:59:ASN:OD1	2.18	0.44
1:C:85:LEU:HB2	1:C:156:VAL:HG11	1.99	0.44
1:B:51:ASP:OD1	1:B:53:SER:OG	2.36	0.43
1:D:114:LEU:HD12	1:D:164:ALA:HB2	2.00	0.42
1:A:137:LEU:HD22	1:A:149:VAL:HG11	2.01	0.42
1:B:121:GLN:HE22	3:B:302:GOL:C1	2.33	0.42
1:D:92:SER:HB3	1:D:94:PHE:CE2	2.54	0.42
1:A:37:HIS:HD2	1:A:70:GLU:OE1	2.02	0.42
1:D:182:VAL:O	1:D:186:ARG:HG2	2.19	0.42
1:C:89:CYS:O	1:C:156:VAL:HG12	2.20	0.42
1:C:91:TYR:HB2	1:C:154:VAL:CG1	2.50	0.41
1:C:85:LEU:HD11	1:C:91:TYR:CE2	2.55	0.41
1:B:126:LEU:HD12	1:B:163:ILE:HD11	2.02	0.41
1:B:35:ILE:HG22	1:B:36:LYS:O	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	182/190 (96%)	176 (97%)	6 (3%)	0	100	100
1	B	181/190 (95%)	178 (98%)	3 (2%)	0	100	100
1	C	176/190 (93%)	170 (97%)	6 (3%)	0	100	100
1	D	176/190 (93%)	170 (97%)	6 (3%)	0	100	100
All	All	715/760 (94%)	694 (97%)	21 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	153/169 (90%)	152 (99%)	1 (1%)	76	87
1	B	152/169 (90%)	150 (99%)	2 (1%)	61	76
1	C	108/169 (64%)	108 (100%)	0	100	100
1	D	108/169 (64%)	108 (100%)	0	100	100
All	All	521/676 (77%)	518 (99%)	3 (1%)	78	89

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	148	ASP
1	B	53	SER
1	B	148	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	37	HIS
1	A	41	ASN
1	A	78	HIS
1	A	80	GLN
1	A	120	GLN
1	B	57	GLN
1	B	78	HIS
1	B	120	GLN
1	B	121	GLN
1	B	178	HIS
1	C	155	ASN
1	D	37	HIS
1	D	80	GLN
1	D	107	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	B	302	-	5,5,5	0.33	0	5,5,5	0.47	0
2	G0Z	C	300	-	31,35,81	3.96	5 (16%)	40,56,127	2.03	10 (25%)
2	G0Z	A	301	-	66,70,81	2.90	14 (21%)	89,110,127	1.92	20 (22%)
3	GOL	A	302	-	5,5,5	0.43	0	5,5,5	0.36	0
2	G0Z	B	301	-	31,35,81	3.98	6 (19%)	40,56,127	2.05	11 (27%)
2	G0Z	D	300	-	31,35,81	3.97	5 (16%)	40,56,127	1.99	10 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	302	-	-	4/4/4/4	-
2	G0Z	C	300	-	-	1/20/50/130	0/3/3/9
2	G0Z	A	301	-	-	0/41/103/130	0/7/7/9

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	302	-	-	4/4/4/4	-
2	G0Z	B	301	-	-	3/20/50/130	0/3/3/9
2	G0Z	D	300	-	-	2/20/50/130	0/3/3/9

All (30) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	300	G0Z	C6-N5	-19.54	1.33	1.45
2	C	300	G0Z	C6-N5	-19.38	1.33	1.45
2	A	301	G0Z	C6-N5	-19.32	1.33	1.45
2	B	301	G0Z	C6-N5	-19.02	1.33	1.45
2	C	300	G0Z	O1-C1	6.93	1.36	1.23
2	B	301	G0Z	O1-C1	6.90	1.36	1.23
2	D	300	G0Z	O1-C1	6.82	1.36	1.23
2	A	301	G0Z	O1-C1	6.45	1.35	1.23
2	B	301	G0Z	C3-N5	-4.28	1.32	1.37
2	A	301	G0Z	C3-N5	-4.01	1.32	1.37
2	D	300	G0Z	C3-N5	-3.96	1.33	1.37
2	B	301	G0Z	P1-O8	-3.73	1.55	1.59
2	D	300	G0Z	C4-C1	-3.72	1.34	1.43
2	C	300	G0Z	C4-C1	-3.65	1.34	1.43
2	B	301	G0Z	C4-C1	-3.61	1.34	1.43
2	C	300	G0Z	C3-N5	-3.50	1.33	1.37
2	A	301	G0Z	C20-N8	3.49	1.40	1.33
2	A	301	G0Z	C20-N9	3.47	1.40	1.33
2	A	301	G0Z	C4-C1	-3.44	1.34	1.43
2	A	301	G0Z	C18-N7	-3.00	1.33	1.39
2	A	301	G0Z	C18-C19	-2.93	1.33	1.39
2	A	301	G0Z	C18-C21	-2.62	1.33	1.41
2	B	301	G0Z	C1-N1	-2.57	1.34	1.38
2	A	301	G0Z	C17-N6	-2.39	1.33	1.37
2	C	300	G0Z	C1-N1	-2.37	1.34	1.38
2	D	300	G0Z	C1-N1	-2.36	1.34	1.38
2	A	301	G0Z	C1-N1	-2.24	1.34	1.38
2	A	301	G0Z	P1-O8	2.19	1.61	1.59
2	A	301	G0Z	C19-N6	-2.08	1.33	1.37
2	A	301	G0Z	C7-N5	2.04	1.50	1.46

All (51) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301	G0Z	N9-C20-N8	-6.85	118.21	128.58
2	A	301	G0Z	C4-C1-N1	6.16	121.77	110.94
2	B	301	G0Z	C4-C1-N1	5.93	121.37	110.94
2	C	300	G0Z	C4-C1-N1	5.90	121.31	110.94
2	D	300	G0Z	C4-C1-N1	5.86	121.26	110.94
2	A	301	G0Z	C2-N3-C3	4.96	120.85	112.30
2	C	300	G0Z	C2-N3-C3	4.62	120.26	112.30
2	D	300	G0Z	C2-N3-C3	4.60	120.22	112.30
2	B	301	G0Z	C2-N3-C3	4.51	120.06	112.30
2	A	301	G0Z	C18-C19-N8	-4.29	120.81	126.72
2	A	301	G0Z	N6-C17-N7	-4.28	107.86	113.94
2	A	301	G0Z	C4-C3-N3	-4.07	120.49	128.13
2	D	300	G0Z	O1-C1-C4	-4.05	117.68	127.62
2	C	300	G0Z	C4-C3-N3	-4.04	120.54	128.13
2	C	300	G0Z	O1-C1-C4	-3.93	117.98	127.62
2	A	301	G0Z	O1-C1-C4	-3.86	118.15	127.62
2	D	300	G0Z	C4-C3-N3	-3.83	120.94	128.13
2	B	301	G0Z	C4-C3-N3	-3.82	120.96	128.13
2	C	300	G0Z	N5-C3-N3	3.70	130.88	125.46
2	B	301	G0Z	O1-C1-C4	-3.64	118.69	127.62
2	D	300	G0Z	N5-C3-N3	3.59	130.71	125.46
2	C	300	G0Z	N5-C6-N4	3.47	108.28	103.37
2	A	301	G0Z	C2-N1-C1	-3.37	119.00	125.11
2	A	301	G0Z	N5-C3-N3	3.36	130.39	125.46
2	B	301	G0Z	N5-C3-N3	3.28	130.26	125.46
2	A	301	G0Z	C20-N8-C19	3.19	119.62	111.83
2	B	301	G0Z	C2-N1-C1	-3.18	119.35	125.11
2	D	300	G0Z	N5-C6-N4	3.17	107.86	103.37
2	B	301	G0Z	N5-C6-N4	3.11	107.77	103.37
2	C	300	G0Z	C2-N1-C1	-3.05	119.58	125.11
2	D	300	G0Z	C2-N1-C1	-2.97	119.72	125.11
2	A	301	G0Z	N5-C6-N4	2.97	107.58	103.37
2	A	301	G0Z	C18-N7-C17	2.95	108.09	103.45
2	A	301	G0Z	C18-C21-N10	-2.81	116.34	123.29
2	A	301	G0Z	C21-C18-C19	2.56	120.68	117.18
2	A	301	G0Z	N8-C19-N6	2.51	131.44	127.17
2	A	301	G0Z	C19-N6-C17	2.46	108.33	105.74
2	A	301	G0Z	C1-C4-C3	-2.29	118.37	122.40
2	A	301	G0Z	C19-C18-N7	-2.28	107.97	110.58
2	C	300	G0Z	O11-P3-O12	2.22	119.49	110.83
2	B	301	G0Z	C1-C4-C3	-2.22	118.50	122.40
2	D	300	G0Z	C1-C4-C3	-2.21	118.52	122.40
2	D	300	G0Z	O11-P3-O12	2.18	119.34	110.83

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	300	G0Z	N1-C2-N3	-2.18	119.34	123.32
2	A	301	G0Z	C4-C3-N5	2.17	109.11	106.33
2	B	301	G0Z	O5-C11-C10	2.15	116.31	108.99
2	B	301	G0Z	O11-P3-O12	2.14	119.18	110.83
2	C	300	G0Z	C1-C4-C3	-2.11	118.70	122.40
2	A	301	G0Z	N1-C2-N3	-2.08	119.51	123.32
2	B	301	G0Z	O13-P3-O12	-2.08	102.75	110.83
2	C	300	G0Z	N1-C2-N3	-2.03	119.60	123.32

There are no chirality outliers.

All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	301	G0Z	P1-O8-P2-O10
2	B	301	G0Z	P2-O10-P3-O13
3	B	302	GOL	O1-C1-C2-C3
3	A	302	GOL	O1-C1-C2-O2
3	B	302	GOL	O1-C1-C2-O2
3	A	302	GOL	O1-C1-C2-C3
3	A	302	GOL	C1-C2-C3-O3
3	B	302	GOL	C1-C2-C3-O3
3	A	302	GOL	O2-C2-C3-O3
3	B	302	GOL	O2-C2-C3-O3
2	C	300	G0Z	C10-C11-O5-P1
2	D	300	G0Z	C10-C11-O5-P1
2	B	301	G0Z	P3-O10-P2-O9
2	D	300	G0Z	P3-O10-P2-O9


There are no ring outliers.


4 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	302	GOL	2	0
2	C	300	G0Z	3	0
2	B	301	G0Z	1	0
2	D	300	G0Z	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.

Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	183/190 (96%)	0.18	4 (2%) 62 59	23, 36, 60, 84	3 (1%)
1	B	184/190 (96%)	0.27	6 (3%) 49 46	23, 37, 65, 91	1 (0%)
1	C	180/190 (94%)	1.29	47 (26%) 1 1	32, 69, 108, 122	0
1	D	180/190 (94%)	1.10	36 (20%) 3 2	32, 66, 96, 104	0
All	All	727/760 (95%)	0.71	93 (12%) 7 5	23, 47, 96, 122	4 (0%)

All (93) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	142	PHE	4.3
1	C	172	ASN	3.9
1	A	31	PRO	3.7
1	C	194	VAL	3.4
1	C	33	HIS	3.4
1	C	201	ALA	3.4
1	C	196	GLY	3.3
1	C	55	THR	3.3
1	C	197	TYR	3.3
1	D	113	TRP	3.3
1	C	114	LEU	3.2
1	D	145	TYR	3.2
1	C	173	ARG	3.1
1	C	117	LEU	3.1
1	A	205	THR	3.1
1	C	195	ILE	3.1
1	D	215	PHE	3.1
1	B	30	ASN	3.1
1	B	158	ALA	3.0
1	D	197	TYR	3.0
1	D	102	TRP	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	179	ILE	2.9
1	A	211	THR	2.9
1	C	31	PRO	2.9
1	C	215	PHE	2.8
1	D	32	GLU	2.8
1	C	193	ILE	2.8
1	D	55	THR	2.8
1	D	115	ILE	2.8
1	C	171	GLU	2.8
1	D	196	GLY	2.7
1	C	145	TYR	2.7
1	D	48	PHE	2.7
1	C	217	VAL	2.7
1	D	158	ALA	2.7
1	C	216	VAL	2.7
1	D	179	ILE	2.7
1	D	199	SER	2.7
1	C	170	CYS	2.7
1	C	202	ASP	2.6
1	D	60	LEU	2.6
1	C	176	VAL	2.6
1	C	141	SER	2.5
1	C	148	ASP	2.5
1	D	33	HIS	2.5
1	D	56	TRP	2.5
1	D	114	LEU	2.5
1	B	205	THR	2.5
1	C	113	TRP	2.5
1	C	175	ALA	2.5
1	B	160	GLY	2.5
1	C	203	THR	2.4
1	D	202	ASP	2.4
1	C	146	SER	2.4
1	A	193	ILE	2.4
1	D	187	LEU	2.3
1	D	160	GLY	2.3
1	C	32	GLU	2.3
1	C	180	GLY	2.3
1	C	102	TRP	2.3
1	D	168	THR	2.3
1	D	203	THR	2.3
1	B	33	HIS	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	53	SER	2.3
1	D	201	ALA	2.2
1	D	103	GLU	2.2
1	C	105	GLU	2.2
1	C	168	THR	2.2
1	D	211	THR	2.2
1	C	191	PRO	2.2
1	C	160	GLY	2.2
1	C	204	ALA	2.2
1	D	214	ARG	2.1
1	C	159	LYS	2.1
1	C	212	LYS	2.1
1	D	100	PRO	2.1
1	D	58	ALA	2.1
1	D	175	ALA	2.1
1	D	117	LEU	2.1
1	D	189	LEU	2.1
1	C	101	MET	2.1
1	D	186	ARG	2.1
1	C	177	THR	2.1
1	C	85	LEU	2.1
1	C	189	LEU	2.1
1	C	143	ASP	2.1
1	B	32	GLU	2.0
1	D	143	ASP	2.0
1	C	100	PRO	2.0
1	D	190	PRO	2.0
1	C	56	TRP	2.0
1	D	191	PRO	2.0
1	D	195	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	G0Z	D	300	33/73	0.62	0.17	97,99,104,137	0
2	G0Z	C	300	33/73	0.72	0.15	99,103,104,108	0
3	GOL	A	302	6/6	0.84	0.14	51,52,52,53	0
3	GOL	B	302	6/6	0.89	0.10	57,57,57,58	0
2	G0Z	B	301	33/73	0.90	0.10	33,36,63,63	0
2	G0Z	A	301	64/73	0.94	0.09	29,60,75,75	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around G0Z D 300:

2mF_o-DF_c (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
 and green (positive)






Electron density around G0Z C 300:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around G0Z B 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around G0Z A 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.