



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 11:18 AM UTC

PDB ID : 1GLE / pdb_00001gle
Title : CATION PROMOTED ASSOCIATION (CPA) OF A REGULATORY AND
TARGET PROTEIN IS CONTROLLED BY PHOSPHORYLATION
Authors : Feese, M.D.; Meadow, N.D.; Roseman, S.; Pettigrew, D.W.; Remington, S.J.
Deposited on : 1994-03-07
Resolution : 2.94 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.94 Å.



Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	1184 (2.96-2.92)
Ramachandran outliers	187476	1131 (2.96-2.92)
Sidechain outliers	187428	1131 (2.96-2.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	F	168	
2	G	501	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	G3H	G	503	X	-	X	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4996 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GLUCOSE-SPECIFIC PROTEIN III_{Glc}.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	F	161	Total	C	N	O	S	0	0	0
			1157	741	181	233	2			


- Molecule 2 is a protein called GLYCEROL KINASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	G	489	Total	C	N	O	S	0	0	0
			3752	2379	644	710	19			

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	F	1	Total	Zn	0	0
			1	1		
3	G	1	Total	Zn	0	0
			1	1		

- Molecule 4 is GLYCERALDEHYDE-3-PHOSPHATE (CCD ID: G3H) (formula: C₃H₇O₆P).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	G	1	Total	C	O	P	0	0
			10	3	6	1		

- Molecule 5 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	G	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	F	7	Total 7	O 7	0	0
6	G	41	Total 41	O 41	0	0

W454	Q455	M456	L457	D458	E459	L460	Q461	K463	A464	V465	L466	E467	R468	E469	F470	R471	P472	G473	I474	E475	T476	T477	E478	R479	M480	Y481	R482		G485	W486	K487	K488	A489	V490	K491	R492	A493	M494	A495	W496	E497	E498	H499	ASP	GLU
------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	--	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	-----	-----

4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	124.03Å 125.11Å 133.33Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	21.80 – 2.94	Depositor
% Data completeness (in resolution range)	(Not available) (21.80-2.94)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	TNT	Depositor
R, R_{free}	0.149 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	4996	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ADP, G3H, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	F	1.76	19/1172 (1.6%)	2.28	64/1592 (4.0%)
2	G	1.63	39/3831 (1.0%)	2.17	165/5211 (3.2%)
All	All	1.66	58/5003 (1.2%)	2.20	229/6803 (3.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	G	2	0

The worst 5 of 58 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	G	39	TYR	N-CA	-7.25	1.37	1.46
2	G	38	ILE	CA-C	-7.12	1.44	1.52
1	F	104	LYS	CA-C	6.96	1.62	1.52
1	F	91	PHE	CA-C	-6.76	1.44	1.52
2	G	101	ILE	CA-CB	-6.64	1.45	1.54

The worst 5 of 229 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	G	336	VAL	N-CA-C	14.68	127.29	113.20
2	G	185	ASN	CA-CB-CG	-12.27	100.33	112.60
1	F	36	VAL	N-CA-CB	10.74	120.07	110.08
2	G	404	HIS	CA-CB-CG	-10.21	103.59	113.80
2	G	107	ARG	N-CA-C	9.56	121.30	111.07

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	G	255	CYS	CA
2	G	480	ASN	CA

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	1157	0	1133	148	0
2	G	3752	0	3605	408	0
3	F	1	0	0	0	0
3	G	1	0	0	0	0
4	G	10	0	5	4	0
5	G	27	0	12	0	0
6	F	7	0	0	1	0
6	G	41	0	0	5	0
All	All	4996	0	4755	555	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 57.

The worst 5 of 555 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:40:VAL:HA	1:F:45:ILE:HD12	1.18	1.10
2:G:7:VAL:HG22	2:G:20:VAL:HG22	1.32	1.09
2:G:419:MET:HE2	2:G:419:MET:HA	1.38	1.04
1:F:30:ILE:HD11	1:F:163:VAL:HG12	1.38	1.03
2:G:152:ARG:HH21	2:G:208:ASP:HB3	1.32	0.95

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	F	157/168 (94%)	133 (85%)	18 (12%)	6 (4%)	2	5
2	G	485/501 (97%)	410 (84%)	58 (12%)	17 (4%)	3	7
All	All	642/669 (96%)	543 (85%)	76 (12%)	23 (4%)	2	6

5 of 23 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	F	109	GLU
1	F	124	LEU
2	G	42	PRO
2	G	175	GLN
2	G	255	CYS

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	F	123/145 (85%)	77 (63%)	46 (37%)	0	0
2	G	378/412 (92%)	274 (72%)	104 (28%)	0	0
All	All	501/557 (90%)	351 (70%)	150 (30%)	0	0

5 of 150 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	G	264	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	G	475	GLU
2	G	295	THR
2	G	403	LEU
2	G	6	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
2	G	456	ASN
2	G	396	GLN
2	G	185	ASN
2	G	387	GLN
2	G	81	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	ADP	G	504	-	28,29,29	1.16	1 (3%)	43,45,45	2.83	7 (16%)
4	G3H	G	503	-	8,9,9	2.35	1 (12%)	7,12,12	1.06	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	ADP	G	504	-	-	6/16/32/32	0/3/3/3
4	G3H	G	503	-	1/1/2/3	2/7/8/8	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	G	503	G3H	O1-C1	6.32	1.44	1.20
5	G	504	ADP	PA-O3A	4.94	1.64	1.59

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	G	504	ADP	O2A-PA-O3A	11.21	137.56	107.27
5	G	504	ADP	O3A-PA-O1A	-7.69	87.58	110.70
5	G	504	ADP	O2'-C2'-C1'	6.51	132.51	110.10
5	G	504	ADP	C1'-N9-C8	6.25	140.97	127.09
5	G	504	ADP	C4-N9-C1'	-6.19	112.16	126.63

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	G	503	G3H	C2

5 of 8 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	G	503	G3H	C1-C2-C3-O1P
4	G	503	G3H	O2-C2-C3-O1P
5	G	504	ADP	C3'-C4'-C5'-O5'
5	G	504	ADP	O4'-C4'-C5'-O5'
5	G	504	ADP	PA-O3A-PB-O2B

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	G	503	G3H	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.