



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 08:56 AM UTC

PDB ID : 2GLQ / pdb_00002glq
Title : X-ray structure of human alkaline phosphatase in complex with strontium
Authors : Llinas, P.; Masella, M.; Stigbrand, T.; Menez, A.; Stura, E.A.; Le Du, M.H.
Deposited on : 2006-04-05
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4673 (1.60-1.60)
Clashscore	190562	4931 (1.60-1.60)
Ramachandran outliers	187476	4831 (1.60-1.60)
Sidechain outliers	187428	4830 (1.60-1.60)
RSRZ outliers	180081	4672 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	484	2% 83% 13% ...

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 4289 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alkaline phosphatase, placental type.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
1	A	479	3660	2288	652	701	1	18	0	0	0

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	14	8	1	5	0	0
2	A	1	14	8	1	5	0	0

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Zn	0	0
			2	2		

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mg	0	0
			1	1		

- Molecule 5 is STRONTIUM ION (CCD ID: SR) (formula: Sr).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Sr	0	0
			1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	597	Total	O	0	0
			597	597		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Alkaline phosphatase, placental type



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	89.09Å 115.30Å 107.32Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	9.99 – 1.60 9.99 – 1.60	Depositor EDS
% Data completeness (in resolution range)	99.0 (9.99-1.60) 98.5 (9.99-1.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.96 (at 1.60Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.148 , 0.188 (Not available) , 0.193	Depositor DCC
R_{free} test set	3637 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	14.9	Xtriage
Anisotropy	0.150	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.44 , 64.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4289	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.69% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SR, NAG, ZN, MG, SEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.97	3/3730 (0.1%)	1.04	17/5059 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	462	MET	SD-CE	13.91	2.14	1.79
1	A	469	GLU	CA-C	-9.32	1.41	1.52
1	A	469	GLU	CA-CB	-7.22	1.42	1.53

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	474	CYS	N-CA-C	-8.57	99.27	110.33
1	A	472	THR	N-CA-C	-8.22	98.90	110.50
1	A	431	THR	N-CA-CB	-7.70	98.08	110.63
1	A	469	GLU	CA-C-N	-7.12	113.57	120.83
1	A	469	GLU	C-N-CA	-7.12	113.57	120.83
1	A	474	CYS	CA-C-N	-6.93	111.45	122.95
1	A	474	CYS	C-N-CA	-6.93	111.45	122.95
1	A	260	PRO	N-CA-C	-6.67	101.40	111.41
1	A	26	LEU	CB-CA-C	6.09	119.82	109.53
1	A	60	LYS	CA-C-N	-5.69	113.39	122.66

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	60	LYS	C-N-CA	-5.69	113.39	122.66
1	A	263	THR	N-CA-C	5.68	122.91	110.80
1	A	62	LYS	N-CA-C	5.68	117.86	109.69
1	A	472	THR	CA-C-N	-5.42	114.93	122.74
1	A	472	THR	C-N-CA	-5.42	114.93	122.74
1	A	292	THR	N-CA-CB	5.31	118.02	110.16
1	A	469	GLU	CA-C-O	-5.24	112.98	120.16

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	468	LEU	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3660	0	3548	77	0
2	A	28	0	26	0	0
3	A	2	0	0	0	0
4	A	1	0	0	0	0
5	A	1	0	0	0	0
6	A	597	0	0	29	1
All	All	4289	0	3574	77	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (77) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:462:MET:CE	1:A:462:MET:SD	2.14	1.34
1:A:251:THR:HB	6:A:2585:HOH:O	1.35	1.27
1:A:392:LEU:HG	6:A:2580:HOH:O	1.09	1.26

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:243:GLY:O	1:A:262:VAL:O	1.64	1.15
1:A:28:PRO:HB3	1:A:447:HIS:ND1	1.80	0.96
1:A:254:MET:HG3	6:A:2179:HOH:O	1.67	0.94
1:A:392:LEU:CD2	6:A:2580:HOH:O	2.14	0.84
1:A:392:LEU:CG	6:A:2580:HOH:O	1.79	0.82
1:A:453:GLN:HG3	6:A:2361:HOH:O	1.78	0.82
1:A:162:HIS:H	1:A:189:GLN:HE22	1.27	0.81
1:A:472:THR:O	1:A:473:ALA:HB3	1.83	0.79
1:A:391:LEU:C	1:A:392:LEU:HD22	2.12	0.74
1:A:389:THR:O	1:A:392:LEU:HD21	1.89	0.72
1:A:131:SER:O	1:A:135:ARG:HD3	1.89	0.72
1:A:469:GLU:CA	6:A:2562:HOH:O	2.36	0.71
1:A:26:LEU:C	1:A:26:LEU:HD23	2.16	0.71
1:A:28:PRO:CB	1:A:447:HIS:ND1	2.54	0.70
1:A:301:ARG:NE	6:A:2581:HOH:O	2.03	0.70
1:A:422:GLN:NE2	6:A:2600:HOH:O	2.26	0.68
1:A:472:THR:O	6:A:2584:HOH:O	2.15	0.63
1:A:26:LEU:HD21	6:A:2231:HOH:O	1.98	0.63
1:A:28:PRO:CB	1:A:447:HIS:CE1	2.82	0.62
1:A:260:PRO:O	1:A:261:SER:CB	2.46	0.62
1:A:469:GLU:O	6:A:2562:HOH:O	2.16	0.62
1:A:469:GLU:HA	6:A:2562:HOH:O	2.00	0.62
1:A:391:LEU:O	1:A:392:LEU:HD22	1.99	0.62
1:A:28:PRO:HB3	1:A:447:HIS:CE1	2.36	0.61
1:A:469:GLU:CB	1:A:470:PRO:CD	2.78	0.61
1:A:431:THR:HG23	1:A:432:HIS:O	2.01	0.60
1:A:409:VAL:HB	6:A:2600:HOH:O	2.01	0.60
1:A:26:LEU:HD23	1:A:26:LEU:O	2.02	0.60
1:A:255:GLN:NE2	6:A:2387:HOH:O	2.36	0.58
1:A:389:THR:O	1:A:392:LEU:CD2	2.51	0.58
1:A:90:PRO:O	1:A:432:HIS:HD2	1.87	0.57
1:A:469:GLU:C	6:A:2562:HOH:O	2.49	0.54
1:A:302:ASN:ND2	1:A:304:ARG:H	2.04	0.54
1:A:472:THR:O	1:A:473:ALA:CB	2.46	0.54
1:A:111:GLY:O	1:A:162:HIS:HD2	1.90	0.54
1:A:135:ARG:HH21	1:A:469:GLU:CB	2.21	0.53
1:A:259:ASP:OD2	1:A:260:PRO:O	2.27	0.53
1:A:28:PRO:HB2	1:A:447:HIS:CE1	2.44	0.53
1:A:320:HIS:CD2	1:A:360:HIS:HE1	2.26	0.53
1:A:285:ASP:CG	6:A:2517:HOH:O	2.53	0.52
1:A:95:THR:HG22	1:A:455:GLN:HE22	1.75	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:26:LEU:HD21	6:A:2379:HOH:O	2.10	0.51
1:A:447:HIS:HD2	1:A:448:LEU:N	2.09	0.50
1:A:235:GLN:HG3	6:A:2281:HOH:O	2.11	0.50
1:A:240:LYS:HD3	6:A:2277:HOH:O	2.12	0.50
1:A:472:THR:HG22	6:A:2127:HOH:O	2.12	0.49
1:A:26:LEU:CD2	6:A:2231:HOH:O	2.59	0.48
1:A:453:GLN:NE2	6:A:2598:HOH:O	2.47	0.48
1:A:353:LEU:HD12	1:A:458:ILE:HG23	1.96	0.48
1:A:393:TYR:O	1:A:423:SER:HA	2.13	0.48
1:A:172:ALA:HA	1:A:228:LEU:HD22	1.95	0.47
1:A:251:THR:HG23	6:A:2080:HOH:O	2.15	0.46
1:A:288:LEU:O	1:A:292:THR:HG23	2.16	0.46
1:A:472:THR:HG23	6:A:2390:HOH:O	2.16	0.45
1:A:450:HIS:HE1	6:A:2083:HOH:O	1.99	0.45
1:A:240:LYS:CE	6:A:2277:HOH:O	2.65	0.44
1:A:153:HIS:CD2	1:A:154:ALA:H	2.35	0.44
1:A:8:ASN:C	1:A:8:ASN:HD22	2.26	0.44
1:A:302:ASN:HD22	1:A:302:ASN:C	2.26	0.43
1:A:473:ALA:C	1:A:474:CYS:O	2.55	0.43
1:A:162:HIS:HE1	6:A:2153:HOH:O	2.02	0.43
1:A:270:GLU:HG3	1:A:274:MET:HA	2.01	0.43
1:A:302:ASN:HD22	1:A:303:PRO:N	2.17	0.43
1:A:8:ASN:HD22	1:A:9:PRO:N	2.17	0.43
1:A:84:ASN:HD21	1:A:91:ASP:H	1.65	0.43
1:A:398:GLY:HA3	1:A:421:GLN:O	2.19	0.43
1:A:447:HIS:CD2	1:A:447:HIS:C	2.97	0.43
1:A:8:ASN:HD22	1:A:9:PRO:HD2	1.84	0.42
1:A:26:LEU:C	1:A:26:LEU:CD2	2.91	0.42
1:A:162:HIS:H	1:A:189:GLN:NE2	2.05	0.42
1:A:458:ILE:HG13	6:A:2558:HOH:O	2.20	0.41
1:A:260:PRO:O	1:A:261:SER:HB3	2.18	0.40
1:A:447:HIS:CD2	1:A:448:LEU:N	2.90	0.40
1:A:302:ASN:HD22	1:A:304:ARG:H	1.67	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:2131:HOH:O	6:A:2601:HOH:O[4_555]	1.84	0.36

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	476/484 (98%)	464 (98%)	10 (2%)	2 (0%)	30	14

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	469	GLU
1	A	361	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	377/386 (98%)	361 (96%)	16 (4%)	26	7

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	26	LEU
1	A	91	ASP
1	A	132	VAL
1	A	228	LEU
1	A	241	ARG
1	A	242	GLN
1	A	245	ARG
1	A	261	SER
1	A	268	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	288	LEU
1	A	292	THR
1	A	296	LEU
1	A	402	LYS
1	A	431	THR
1	A	475	ASP
1	A	476	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	8	ASN
1	A	84	ASN
1	A	127	ASN
1	A	153	HIS
1	A	162	HIS
1	A	167	ASN
1	A	180	GLN
1	A	189	GLN
1	A	193	ASN
1	A	264	HIS
1	A	302	ASN
1	A	320	HIS
1	A	432	HIS
1	A	445	GLN
1	A	447	HIS
1	A	450	HIS
1	A	455	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	SEP	A	92	3,1	8,9,10	1.47	2 (25%)	7,12,14	3.02	3 (42%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SEP	A	92	3,1	-	1/6/8/10	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	92	SEP	P-OG	2.94	1.69	1.60
1	A	92	SEP	P-O1P	2.21	1.57	1.50

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	92	SEP	OG-P-O1P	-5.63	91.22	106.44
1	A	92	SEP	OG-CB-CA	4.10	112.13	108.14
1	A	92	SEP	O3P-P-O2P	2.43	116.92	107.80

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	92	SEP	CA-CB-OG-P

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	A	1001	1	14,14,15	0.59	0	17,19,21	1.80	4 (23%)
2	NAG	A	1003	1	14,14,15	0.59	0	17,19,21	1.43	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	1001	1	-	2/6/23/26	0/1/1/1
2	NAG	A	1003	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1001	NAG	C1-O5-C5	5.14	119.08	112.19
2	A	1003	NAG	O5-C1-C2	-3.41	106.01	111.29
2	A	1001	NAG	C3-C4-C5	2.51	114.78	110.23
2	A	1001	NAG	C1-C2-N2	2.14	113.81	110.43
2	A	1001	NAG	C6-C5-C4	-2.11	107.85	113.02
2	A	1003	NAG	C3-C4-C5	-2.06	106.50	110.23
2	A	1003	NAG	C4-C3-C2	-2.01	108.07	111.02

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1001	NAG	O5-C5-C6-O6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	1001	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	478/484 (98%)	-0.57	8 (1%) 69 72	11, 17, 31, 44	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	475	ASP	3.6
1	A	472	THR	3.2
1	A	262	VAL	3.1
1	A	61	ASP	2.6
1	A	260	PRO	2.4
1	A	473	ALA	2.3
1	A	403	ASP	2.1
1	A	447	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	SEP	A	92	10/11	0.99	0.03	11,13,14,14	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	NAG	A	1001	14/15	0.79	0.10	30,38,44,45	0
2	NAG	A	1003	14/15	0.91	0.07	23,28,33,34	0
4	MG	A	2003	1/1	0.99	0.02	12,12,12,12	0
3	ZN	A	2002	1/1	1.00	0.01	11,11,11,11	0
3	ZN	A	2001	1/1	1.00	0.01	12,12,12,12	0
5	SR	A	2004	1/1	1.00	0.04	22,22,22,22	0

6.5 Other polymers [i](#)

There are no such residues in this entry.